- 5 Brusasco V, Pellegrino R, Rodarte JR. Vital capacities in acute and chronic airway obstruction: dependence on flow and volume histories *Eur Respir J* 1997; 10: 1316–1320.
- 6 Paoletti P, Pistelli G, Fazzi P, *et al.* Reference values for vital capacity and flow–volume curves from a general

population study *Bull Eur Physiopathol Respir* 1986; 22: 451–459.

DOI: 10.1183/09031936.00087507

Antibiotics for chronic bronchitis: a meta-analysis that speaks only four languages

To the Editors:

We have read with interest the meta-analysis by SIEMPOS *et al.* [1] evaluating the use of different antibiotics for exacerbations of chronic bronchitis. We are puzzled by the criteria used to select the studies that were included in the analysis. The authors utilised a well-established methodology and quality criteria for the selection of randomised clinical trials, but only studies written in English, French, German or Italian were included in the analysis [1]. English is now accepted as the most common language in scientific literature; however, there is an increasing number of peer-reviewed scientific journals written in languages other than English that have published clinical studies with data relevant to the issue of this meta-analysis. In fact, among the top 25 respiratory journals ranked by impact factor there is one published in Spanish with an impact factor of 1.401 [2].

It is not clear to us why SIEMPOS *et al.* [1] have reviewed articles written in French, German and Italian and excluded the others. Most of the indexed non-English journals include English abstracts with information that may help verify the quality of the study. If the study proved to be scientifically important, there are individuals, organisations and companies that provide translation services that would help make the overall goals more accurate.

In the article by SIEMPOS *et al.* [1], five studies were excluded due to language selection. Considering that in one of the comparisons, conclusions were drawn based on only four studies, the inclusion of one or more of the excluded studies could have impacted on the final results. Furthermore, the authors did not provide references for the publications that were excluded. Provided with this information, the interested reader could compare the results of the excluded works with those of the studies included in the meta-analysis and verify whether this arbitrary exclusion could have potentially biased the results.

Therefore, we believe that the *European Respiratory Journal* should not accept exclusion criteria beyond those that are strictly based on science. The exclusion of studies based on the language of publication is simply unacceptable.

M. Miravittles*, B. Celli[#] and A. Anzueto[†]

*Dept of Pneumology, Institut Clínic del Tòrax (IDIBAPS), Hospital Clínic, Barcelona, Spain. [#]St Elizabeth's Medical

Center, Brighton, MA. [†]Health Science Center, University of Texas, San Antonio, TX, USA.

STATEMENT OF INTEREST

None declared.

REFERENCES

- 1 Siempos II, Dimopoulos G, Korbila IP, Manta K, Falagas ME. Macrolides, quinolones and amoxicillin/clavulanate for chronic bronchitis: a meta-analysis *Eur Respir J* 2007; 29: 1127–1137.
- 2 ISI Web of Knowledge. Journal citation reports. <http://portal.isiknowledge.com/portal.cgi>. Date last accessed: June 12, 2007.

DOI: 10.1183/09031936.00071607

From the authors:

We would like to thank M. Miravittles and co-workers for their interest in our meta-analysis [1]. They commented on our choice not to include randomised controlled trials (RCTs) in the meta-analysis that were written in languages other than English, French, German or Italian. In addition, they criticised the fact that the references of the excluded RCTs were not available. We welcome the letter by our respectable colleagues and would like to respond to the points they raised.

We would like to emphasise that our research team's policy regarding the reporting of meta-analyses is to provide the references of all excluded trials to secure transparency and reproducibility of our work. For instance, in another meta-analysis performed by our team that has been recently published in the *European Respiratory Journal* (ERJ), we have done so [2]. However, for the contribution under discussion, we would have to have provided 139 additional references (*i.e.* the number of the studies that were omitted for various reasons as explained in the relevant figure of our article); we considered that this number was probably excessive, given the space limitations of the ERJ.

We carefully reviewed the abstracts of the five RCTs that were excluded due to language restrictions [3–7]. None of them provided data relevant to the subject of our meta-analysis (namely the comparison of macrolides, quinolones and amoxicillin/clavulanate for the treatment of patients with