

## Detecting uncontrolled asthma in children

Comparing GINA criteria with the Childhood Asthma Control Test and Asthma Control Test

B.B. Koolen<sup>1</sup>, M.W.H. Pijnenburg<sup>2</sup>, H.J.L. Brackel<sup>3</sup>, A.M. Landstra<sup>4</sup>, N.J. van den Berg<sup>5</sup>,  
P.J.F.M. Merkus<sup>6</sup>, W.C.J. Hop<sup>7</sup> and A.A.P.H. Vaessen-Verberne<sup>1</sup>

### Authors:

<sup>1</sup> Amphia Hospital, Department of Pediatrics, Langendijk 75, 4819 EV Breda; <sup>2</sup> Erasmus University Medical Centre – Sophia Children’s Hospital, Department of Pediatric Pulmonology, Dr. Molewaterplein 60, 3015 GJ Rotterdam; <sup>3</sup> Catharina Hospital, Department of Pediatrics, Michelangelolaan 2, 5623 EJ Eindhoven; <sup>4</sup> Rijnstate Hospital, Department of Pediatrics, Wagnerlaan 55, 6815 AD Arnhem; <sup>5</sup> Flevo Hospital, Department of Pediatrics, Hospitaalweg 1, 1315 RA Almere; <sup>6</sup> Radboud University Nijmegen Medical Centre, Department of Pediatric Pulmonology, Geert Grooteplein-Zuid 10, 6525 GA Nijmegen; <sup>7</sup> Erasmus University Medical Centre, Department of Biostatistics, Dr. Molewaterplein 50, 3015 GE Rotterdam – all from the Netherlands

**Corresponding author:**

A.A.P.H. Vaessen-Verberne

Telephone number: 0031 76 5951538

Fax number: 0031 76 5952389

E-mail: [avaessen-verberne@amphia.nl](mailto:avaessen-verberne@amphia.nl)

**Address for reprints:**

Amphia Hospital – Department of Pediatrics

attn A.A.P.H. Vaessen-Verberne, pediatric pulmonologist

Postbus 90157

4800 RL Breda, the Netherlands

Word count abstract 183, word count full article 2763

Numbers of figures and tables: 5 tables, 2 figures

Conflict of interest: none

Statement: neither the work nor any part of its essential substance, tables or figures have been or will be published or submitted to another scientific journal or is being considered for publication elsewhere

**Keywords:**

Asthma, Asthma control, Child, Asthma Control Test, Childhood Asthma Control Test, GINA

**Running title:** Detecting uncontrolled asthma in children

## **Abstract**

Several tools are useful in detecting uncontrolled asthma in children. The aim of this study was to compare GINA guidelines with the Childhood Asthma Control Test (C-ACT) and the Asthma Control Test (ACT) in detecting uncontrolled asthma in children.

145 children with asthma filled in a web-based daily diary card for 4 weeks on symptoms, use of rescue medication and limitations of activities, followed by either the C-ACT or ACT. For predicting uncontrolled asthma, cut-off points of 19 for C-ACT and ACT were used.

According to GINA guidelines, asthma was uncontrolled in 71 children (51%) and completely controlled in 19 children (14%). The area under the curve (AUC) in the Receiver Operating Characteristic (ROC) curves for C-ACT and ACT versus GINA guidelines were 0.89 and 0.92, respectively. Cut-off points of 19 for C-ACT and ACT resulted in a sensitivity of 33% and 66% in predicting uncontrolled asthma, respectively.

C-ACT and ACT correlate well with GINA criteria in predicting uncontrolled asthma, but commonly used cut-off points for C-ACT and ACT seem to underestimate the proportion of children with uncontrolled asthma as defined by GINA.

## Introduction

Based on recent asthma guidelines, like the Global Initiative for Asthma (GINA) guidelines, asthma treatment focuses on achieving and maintaining asthma control rather than on asthma severity<sup>(1-5)</sup>. A step-up in treatment is advised in order to achieve asthma control in uncontrolled patients. GINA guidelines state that “it is reasonable to expect that in most patients with asthma, control of the disease can and should be maintained”. However, despite the availability of guidelines, a substantial proportion of adults as well as children with asthma is not optimally controlled<sup>(6-10)</sup>.

Several tools have been developed to determine the level of control and to guide treatment<sup>(11-17)</sup>. Amongst them is the Asthma Control Test (ACT), which consists of 5 items and has been shown useful in the detection of poorly controlled asthma in adults and children above the age of 12<sup>(12)</sup>. More recently, a 7-item Childhood ACT (C-ACT) has been validated in children 4 – 11 years of age<sup>(14)</sup>. A cut-off point  $\leq 19$  was selected to indicate uncontrolled asthma for both questionnaires. The Asthma Control Questionnaire (ACQ) is perhaps the most used questionnaire worldwide and consists of seven items on symptoms, use of rescue medication and pulmonary function<sup>(11)</sup>. Moreover, a shortened version with five or six items has been validated<sup>(16)</sup>. Although the ACQ has recently been validated for use in children 6 – 16 years of age<sup>(17)</sup>, there is limited experience with this questionnaire in children.

C-ACT, ACT and ACQ have been used extensively in clinical trials. Despite definition of asthma control in GINA guidelines, the gold standard for determining asthma control in most studies was an estimation of the degree of asthma control by a physician<sup>(13, 14, 18-20)</sup>. In general, this estimation was based on a single outpatient visit<sup>(13, 14, 18, 20)</sup> and expressed in scores ranging from 1 (totally uncontrolled) to 5 (well controlled)<sup>(13, 14)</sup> or three groups of control status (uncontrolled, partly controlled or controlled)<sup>(18, 20)</sup>. However, several studies

have shown that both patients and their doctors tend to overestimate the level of control and the extent of improvement achieved with therapy<sup>(6-8, 10, 21, 22)</sup>. Four recently published studies compared ACT or ACQ with asthma control as assessed by GINA guidelines<sup>(23-26)</sup>. Three of them included children above the age of 12<sup>(23, 25, 26)</sup>. Three studies used the ACQ<sup>(23, 24, 26)</sup> and one used the ACT<sup>(25)</sup>.

Until now, no studies have been published comparing C-ACT and ACT to GINA criteria in children. The aim of this study was to compare the results of the ACT and the C-ACT with GINA guidelines in detecting uncontrolled asthma in children.

## **Methods**

This prospective study is the extension of a study validating a web-based version of the C-ACT and ACT (in press *Pediatric Pulmonology*). After completing three web-based or paper-based questionnaires, participants filled in a web-based daily diary card on symptoms, use of rescue medication and limitations of activities for a period of 4 weeks. At the end of this period they completed either the C-ACT (4 – 11 years old) or the ACT (12 years and older).

### *Study population*

Children with asthma, treated in the outpatient clinics of 4 general hospitals and 2 university hospitals, were eligible for this study. Almost all children were treated with inhaled corticosteroids. Children, and parents of children in the younger age group (4 – 11 years), had to be able to speak the Dutch language and should have Internet access at home. Children with lung diseases other than asthma were excluded from the study.

The study was approved by the Medical Ethical Committees of the participating hospitals. All parents and children from the age of 12 signed an informed consent form before entering the study.

#### *Diary and GINA guidelines*

The web-based diary card was filled in once a day, in the afternoon or evening, during four consecutive weeks and was based on GINA criteria (table 1) <sup>(1)</sup>. In this study we used slightly modified criteria, since we didn't take lung function into account in assessing asthma control. Patients were allowed to fill in diary cards for the preceding 3 days. Diary cards contained questions on daytime symptoms, nocturnal symptoms, limitation of activity and use of rescue medication. Daytime symptoms as coughing, wheezing and shortness of breath were scored on a scale from 0 (no complaints) to 3 (complaints during most of the day). In the same way, nocturnal symptoms as coughing, wheezing and shortness of breath were scored from 0 (no complaints) to 3 (hardly no sleep due to respiratory symptoms). Limitation of activity ranged from 0 (no limitations) to 3 (severe limitation of activities) as well. Children and/or their parents could report the use of rescue medication as extra puffs taken.

Access to the web-based diary was granted by a personal account with username and password. After logging in to the secured Internet page, participants could answer the questions by clicking the appropriate box. After finishing the questions for a single day, results were submitted.

Control status in any week was assessed according to GINA criteria. An overall score was determined after 4 weeks, being the worst score of control status in any week. A minimum of five completed diary days in one week was considered the minimum to consider the weeks' data in the final assessment of asthma control.

### *Childhood Asthma Control Test*

The Childhood Asthma Control Test consists of 7 items, addresses the previous four weeks and is divided into two parts<sup>(14)</sup>. One part is filled in by the child and consists of 4 questions on perception of asthma control, limitation of activities, coughing and awakenings a night. Each question has 4 response options. The second part is filled in by the parent or caregiver and consists of 3 questions (daytime complaints, daytime wheezing and awakenings at night) with 6 response options. The sum of all scores yields the C-ACT score, ranging from 0 (poorest asthma control) to 27 (optimal asthma control). A cut-off point  $\leq 19$  indicates uncontrolled asthma<sup>(14)</sup>.

### *Asthma Control Test*

The Asthma Control Test is a patient-completed questionnaire and consists of 5 items evaluating the preceding 4 weeks (limitation of activities, shortness of breath, awakenings at night, use of reliever medication and patient's perception of asthma control)<sup>(12, 13)</sup>. Each question has 5 response options, resulting in scores of 1 – 5. The sum of all scores yields the total ACT score, ranging from 5 (poorest asthma control) to 25 (optimal asthma control). It has been validated from the age of 12 and a score  $\leq 19$  indicates poorly controlled asthma . In this study we used the Dutch version, translated by the MAPI-research institute in Lyon, France.

### *Study protocol*

During a regular visit at the pediatric outpatient clinic, 234 children were asked to participate. 173 children (74%) were prospectively included in the study, which was carried out between December 2008 and July 2009. After completing three web- or paper-based questionnaires,

145 participants proceeded into an extended study and filled in a web-based daily diary card, followed by the C-ACT or ACT. A flow chart of the study is shown in figure 1.

### *Statistical analysis*

The diagnostic value of the C-ACT and ACT to detect uncontrolled asthma as defined by GINA criteria was quantified by calculation of area under the curve (AUC) of Receiver Operating Characteristic (ROC) curves. Sensitivity, specificity, positive predicting value, negative predicting value and percentage correctly classified were calculated for different cut-off points of C-ACT and ACT when compared to GINA criteria for uncontrolled asthma.

Pearson Chi-Square was used to determine differences in GINA-defined control status between children with missing weeks and children with complete diaries. The Mann-Whitney test was used to compare ACT and C-ACT scores between control status groups according to GINA.  $P = 0.05$  (two-sided) was the limit of statistical significance.

### **Results**

Out of 173 included children, 145 children filled in the diary cards. The main reason for refusal to participate was lack of interest, whereas one patient did not have Internet access.

Baseline characteristics are shown in table 2 . During the study period no medication changes or exacerbations occurred in the study population. Children who failed to fill in their diaries did not differ in baseline characteristics from participants.

145 children completed at least one diary day. As the study period for the diary comprised 28 days, there were 4060 possible diary days. At the end of the study, 3266 diary days (80%) were filled in. Seventy one out of 139 patients (51%) had uncontrolled asthma in at least one

week. Of these 71 children, 25 children were uncontrolled in 1 week, 20 children were uncontrolled in two weeks, 12 children in 3 weeks and 14 children were uncontrolled in all four weeks. Only 19 patients (14%) were classified as controlled, including 7 children with missing weeks. There were no differences in GINA-defined control between children with missing weeks and children with complete diaries ( $p = 0.713$ ).

Median C-ACT score was 23 (Interquartile Range (IQR) 20.5 – 24), median ACT score was 21 (IQR 18.5 – 24). Overall, 17% (13/78) in the younger age group and 33% (19/57) in the older age group had uncontrolled asthma according to a C-ACT or ACT score  $\leq 19$ . Table 3 shows ACT and C-ACT outcomes in relation to asthma control according to GINA criteria.

Comparing C-ACT to GINA criteria for uncontrolled asthma resulted in an AUC of the ROC curve of 0.89 (95% CI 0.82 – 0.96). Using a cut-off point  $\leq 19$  for detecting uncontrolled asthma with the C-ACT, this resulted in a sensitivity of 33% (13/39) and a specificity of 100% (39/39) for the C-ACT. When comparing the ACT to GINA criteria for uncontrolled asthma, we found an AUC of the ROC curve of 0.92 (95% CI 0.84 – 0.99). For detecting uncontrolled asthma with the ACT, a cut-off point  $\leq 19$  resulted in a sensitivity of 66% (19/29) and a specificity of 100% (28/28). ROC curves for C-ACT and ACT versus uncontrolled asthma according to GINA criteria are shown in figure 2. Excluding children with missing weeks resulted in an AUC of 0.89 (95% CI 0.80 – 0.98) and 0.91 (95% CI 0.81 – 1.01) for C-ACT ( $n = 47$ ) and ACT ( $n = 36$ ) in predicting uncontrolled asthma, respectively. Table 4 and 5 show the performance of C-ACT and ACT in predicting uncontrolled asthma for different cut-off points, respectively.

Out of a total of 458 weeks with sufficient diary days, 157 weeks were classified as uncontrolled, 156 weeks as partially controlled and 145 weeks as controlled according to GINA criteria. Of 1032 evaluable days in weeks with uncontrolled asthma, daytime symptoms occurred in 834 days (81%), limitations of activities in 641 days (62%), nocturnal symptoms in 573 days (56%) and rescue medication in 553 days (54%). Of 1015 evaluable days in weeks with partially controlled asthma, 289 days were scored because of daytime symptoms (28%), 285 days because of limitations of activities (28%), 178 days because of the use of rescue medication (18%) and 88 days were scored because of nocturnal symptoms (9%). Children with a week of uncontrolled or partially controlled asthma are failing on the characteristics of daytime symptoms and limitations of activities in particular. Surprisingly, rescue medication was not used as often as the amount of days with complaints.

## **Discussion**

In this prospective study we compared the results of the C-ACT and ACT with GINA criteria in predicting uncontrolled asthma. We found a good predictive value of both the C-ACT and the ACT for detecting uncontrolled asthma as compared to GINA guidelines. Overall control status in this group of children, using questionnaires to determine the level of control, was acceptable. However, when GINA criteria were used to assess asthma control, a large proportion of children had uncontrolled asthma in at least one out of four weeks (71 of 139 patients, 51%) when symptoms were scored. This high percentage of children with at least one week of uncontrolled asthma might possibly be explained by intercurrent infections, in particular as children were included during winter and spring and the majority had uncontrolled asthma in only 1 or 2 weeks.

Comparing the C-ACT and ACT to GINA criteria, we found an excellent predictive value with an AUC of 0.89 and 0.92, respectively. Using a cut-off point  $\leq 19$  for both tests in predicting uncontrolled asthma, specificity for both tests was 100%; however, sensitivity was low. As children with missing weeks were probably misclassified and overrated in their control status (in case missing weeks were uncontrolled weeks), in this study we might even have overestimated the sensitivity. This low sensitivity is remarkable and suggests that overestimation of asthma control by patients and/or their parents is considerable, as mentioned in previous publications <sup>(6-8, 10, 21)</sup>.

On the basis of the findings in this study and presuming that GINA criteria are the gold standard for control status, one could argue that cut-off points for ACT as well as C-ACT should be increased. However, this study was not designed for validation of these questionnaires and the sample of included patients is too low for such an exercise. A limitation in our study is that we failed to systematically rate the control status by the treating paediatrician. This would have been of special interest as a large proportion of children was classified as uncontrolled by GINA criteria. According to GINA guidelines this would have had the implication of augmenting treatment, usually by increasing inhaled corticosteroid doses. It also raises the question, whether the GINA criteria are probably too strict, as from a clinical point of view those children with one week uncontrolled asthma, due to a short increase in symptoms in a viral infection period, will usually not be treated that way. However, in daily practice GINA criteria will usually not be based on diary cards, but on reporting symptoms, so likely be underreported <sup>(7)</sup>. A second limitation of this study is that we included no follow-up period, so we cannot report on the value of (C)-ACT as a predictor of future exacerbations, which might be an additional aspect of control status.

Most studies use doctor's rating as the gold standard for assessing asthma control and only a few studies assessed asthma control according to GINA criteria. The first study using GINA criteria was published in 2006 by Juniper et al, comparing ACQ with GINA criteria in patients 12 – 80 years old <sup>(23)</sup>. Modified GINA criteria were used to determine the optimal cut-off point for the ACQ and a correlation of 0.76 (no p-value) for ACQ and GINA criteria was found. Second, van den Nieuwenhof et al investigated the correlation of ACQ and GINA criteria in adults <sup>(24)</sup>. Out of 108 asthma patients, 20% was defined as poorly controlled and 53% as very poorly controlled according to GINA criteria in a 4-week registration period. An optimal sum score cut-off value of 4.00 points for ACQ in discriminating between well/moderately controlled asthma and poorly/very poorly controlled asthma was found. A third study using GINA criteria by Thomas et al compared ACT to GINA criteria in almost 3000 patients  $\geq 12$  years of age <sup>(25)</sup>. Asthma control was defined according to modified GINA criteria from single patient and physician response forms with reference to the previous 4 weeks. 68% of patients had uncontrolled asthma according to GINA criteria. The AUC of the ROC curve for ACT was 0.84. Using a cut-off point  $\leq 19$  in predicting uncontrolled or partly controlled asthma, they found a sensitivity of 60% and a specificity of 92%, comparable to our data. Recently, O'Byrne et al compared three classification systems on asthma control (GINA criteria, GOAL criteria and ACQ) in a large group of patients aged 12 years and older (n = 8188) <sup>(26)</sup>. GINA and GOAL criteria agreed very well and GINA controlled/partly controlled and GOAL totally controlled/well controlled correlated with an ACQ-5 score  $< 1.00$ .

In conclusion, C-ACT and ACT compared to GINA guidelines demonstrate good agreement. However, applying GINA criteria resulted in a large proportion of children with uncontrolled asthma for at least one out of four weeks. With commonly used cut-off points of C-ACT and

ACT ( $\leq 19$ ), a fairly large proportion of children is considered controlled, although GINA criteria suggest that they are uncontrolled for at least one out of four weeks. Specificity in detecting uncontrolled asthma was high, but sensitivity was low. Therefore, C-ACT and ACT using generally used cut-off points might underestimate the proportion of children with uncontrolled asthma. Further studies in children are necessary to compare different control measurements with doctor's rating of control status and study the predictive value for future exacerbations.

### **Acknowledgements**

We gratefully thank the involved pediatric asthma nurses and medical students for their help and active guidance throughout the study: Dorien Stevens-Blijlevens, Dorine de Hond, Anouk van Limpt (Amphia Hospital), Mirelle Lagendijk (Erasmus University Medical Centre), Christel Linssen-Geven (Catharina Hospital), Linda van Gilst, Henriette Gerrits (Rijnstate Hospital), Ghislaine van der Zande (Flevo Hospital) and Marianne Bobbert (Radboud University Medical Centre).

## References

1. GINA report, Global Strategy for Asthma Management and Prevention; <http://www.ginasthma.com/GuidelinesResources.asp?l1=2&l2=0>. Date last updated: December 2009. Date last accessed: March 20 2010.
2. Graham LM. Classifying asthma. *Chest*. 2006 Jul;130(1 Suppl):13S-20S.
3. Bateman ED, Hurd SS, Barnes PJ, Bousquet J, Drazen JM, FitzGerald M, et al. Global strategy for asthma management and prevention: GINA executive summary. *Eur Respir J*. 2008 Jan;31(1):143-78.
4. Dutch guideline for the diagnosis and treatment of asthma in children: "Asthma in children"; [http://www.nvk.pedinet.nl/pdfs/200810\\_astma\\_bij\\_kinderen.pdf](http://www.nvk.pedinet.nl/pdfs/200810_astma_bij_kinderen.pdf). Date last updated: April 27 2009. Date last accessed: March 20 2010.
5. Expert Panel Report 3: Guidelines for the Diagnosis and Management of Asthma; <http://www.nhlbi.nih.gov/guidelines/asthma/asthgdln.htm>. Date last updated: August 5 2008. Date last accessed: March 20 2010.
6. Rabe KF, Vermeire PA, Soriano JB, Maier WC. Clinical management of asthma in 1999: the Asthma Insights and Reality in Europe (AIRE) study. *Eur Respir J*. 2000 Nov;16(5):802-7.
7. Sawyer SM, Fardy HJ. Bridging the gap between doctors' and patients' expectations of asthma management. *J Asthma*. 2003 Apr;40(2):131-8.
8. Rabe KF, Adachi M, Lai CK, Soriano JB, Vermeire PA, Weiss KB, et al. Worldwide severity and control of asthma in children and adults: the global asthma insights and reality surveys. *J Allergy Clin Immunol*. 2004 Jul;114(1):40-7.
9. Braman SS. The global burden of asthma. *Chest*. 2006 Jul;130(1 Suppl):4S-12S.

10. FitzGerald JM, Boulet LP, McIvor RA, Zimmerman S, Chapman KR. Asthma control in Canada remains suboptimal: the Reality of Asthma Control (TRAC) study. *Can Respir J*. 2006 Jul-Aug;13(5):253-9.
11. Juniper EF, O'Byrne PM, Guyatt GH, Ferrie PJ, King DR. Development and validation of a questionnaire to measure asthma control. *Eur Respir J*. 1999 Oct;14(4):902-7.
12. Nathan RA, Sorkness CA, Kosinski M, Schatz M, Li JT, Marcus P, et al. Development of the asthma control test: a survey for assessing asthma control. *J Allergy Clin Immunol*. 2004 Jan;113(1):59-65.
13. Schatz M, Sorkness CA, Li JT, Marcus P, Murray JJ, Nathan RA, et al. Asthma Control Test: reliability, validity, and responsiveness in patients not previously followed by asthma specialists. *J Allergy Clin Immunol*. 2006 Mar;117(3):549-56.
14. Liu AH, Zeiger R, Sorkness C, Mahr T, Ostrom N, Burgess S, et al. Development and cross-sectional validation of the Childhood Asthma Control Test. *J Allergy Clin Immunol*. 2007 Apr;119(4):817-25.
15. Murphy KR, Zeiger RS, Kosinski M, Chipps B, Mellon M, Schatz M, et al. Test for respiratory and asthma control in kids (TRACK): a caregiver-completed questionnaire for preschool-aged children. *J Allergy Clin Immunol*. 2009 Apr;123(4):833-9 e9.
16. Juniper EF, Svensson K, Mork AC, Stahl E. Measurement properties and interpretation of three shortened versions of the asthma control questionnaire. *Respir Med*. 2005 May;99(5):553-8.
17. Juniper EF, Gruffydd-Jones K, Ward S, Svensson K. Asthma control questionnaire in children validation, measurement properties, interpretation. *Eur Respir J*. 2010 Jun 7.
18. Zhou X, Ding FM, Lin JT, Yin KS. Validity of asthma control test for asthma control assessment in Chinese primary care settings. *Chest*. 2009 Apr;135(4):904-10.

19. Juniper EF, Chauhan A, Neville E, Chatterjee A, Svensson K, Mork AC, et al. Clinicians tend to overestimate improvements in asthma control: an unexpected observation. *Prim Care Respir J*. 2004 Dec;13(4):181-4.
20. Ko FW, Leung TF, Hui DS, Chu HY, Wong GW, Wong E, et al. Asthma Control Test correlates well with the treatment decisions made by asthma specialists. *Respirology*. 2009 May;14(4):559-66.
21. Boulet LP, Phillips R, O'Byrne P, Becker A. Evaluation of asthma control by physicians and patients: comparison with current guidelines. *Can Respir J*. 2002 Nov-Dec;9(6):417-23.
22. Raheison C, Abouelfath A, Le Gros V, Taytard A, Molimard M. Underdiagnosis of nocturnal symptoms in asthma in general practice. *J Asthma*. 2006 Apr;43(3):199-202.
23. Juniper EF, Bousquet J, Abetz L, Bateman ED. Identifying 'well-controlled' and 'not well-controlled' asthma using the Asthma Control Questionnaire. *Respir Med*. 2006 Apr;100(4):616-21.
24. van den Nieuwenhof L, Schermer T, Eysink P, Halet E, van Weel C, Bindels P, et al. Can the Asthma Control Questionnaire be used to differentiate between patients with controlled and uncontrolled asthma symptoms? A pilot study. *Fam Pract*. 2006 Dec;23(6):674-81.
25. Thomas M, Kay S, Pike J, Williams A, Rosenzweig JR, Hillyer EV, et al. The Asthma Control Test (ACT) as a predictor of GINA guideline-defined asthma control: analysis of a multinational cross-sectional survey. *Prim Care Respir J*. 2009 Mar;18(1):41-9.
26. O'Byrne PM, Reddel HK, Eriksson G, Ostlund O, Peterson S, Sears MR, et al. Measuring asthma control: a comparison of three classification systems. *Eur Respir J*. 2010 Aug;36(2):269-276.

<b>Characteristic</b>	<b>Controlled (all of the following)</b>	<b>Partly controlled (any measure present in any week)</b>	<b>Uncontrolled</b>
Daytime symptoms	None (< 3 / wk)	> 2 / wk	3 or more features of partly controlled asthma present in any week
Limitations of activity	None	Any	
Nocturnal symptoms	None	Any	
Need for rescue treatment	None (< 3 / wk)	> 2 / wk	
Lung function (FEV1 or PEF)	Normal	< 80% predicted or personal best	
Exacerbations	None	1 or more / yr	1 in any week

Table 1: control status according to GINA guidelines.

<b>Characteristic</b>	<b>4 – 11 yrs n = 84</b>	<b>12 – 18 yrs n = 61</b>
Age (yrs)	8.5 ± 2.3	13.9 ± 1.5
Gender		
Male	54 (64%)	36 (59%)
Female	30 (36%)	25 (41%)
Duration of asthma (yrs)	4.7 ± 2.9	8.2 ± 4.3
Hospital admissions last year per child		
0	78 (93%)	56 (92%)
1	5 (6%)	3 (5%)
2	1 (1%)	2 (3%)
Courses of oral steroids last year per child		
0	76 (91%)	53 (87%)
1	5 (6%)	3 (5%)
2	3 (4%)	4 (7%)
3	0	1 (2%)
Daily dose of ICS *	400 (200 – 500)	400 (200 – 500)
Use of long acting beta-mimetics		
No	60 (71%)	19 (31%)
Yes	24 (29%)	42 (69%)
Use of leukotriene receptor antagonists		
No	74 (88%)	49 (80%)
Yes	10 (12%)	12 (20%)
FEV1 (% predicted) #	101.3 ± 16.9	96.6 ± 12.9
FeNO (ppb) (n = 113) #	12.0 (7.9 – 18.9)	17.2 (10.1 – 32.0)

Table 2: baseline characteristics of children who completed at least one diary day (n = 145). Data are shown as absolute numbers (%), as median (interquartile range) or as mean ± SD. \*: Daily dose of ICS = daily dose of inhaled corticosteroids as an equivalent of budesonide. #: Measurement of FEV1 (Forced Expiratory Volume in 1 second) and FeNO (Fraction of exhaled Nitric Oxide) within six months prior to the start of the study.

<b>GINA</b>	<b>Age 4-11 years</b>	<b>C-ACT score</b>	<b>Age 12-18 years</b>	<b>ACT score</b>
Uncontrolled	41 (51%)	21 (6-26)	30 (52%)	18 (10-24)
Partially Controlled	26 (32%)	23 (20-27)	23 (40%)	23 (20-25)
Controlled	14 (17%)	27(23-27)	5 (9%)	25 (24-25)
Total group of children	81	23 (6-27) #	58	21 (10-25) \$

Table 3: ACT and C-ACT outcomes in relation to asthma control according to GINA criteria. Data shown are numbers of patients (%) or median (range); # n=78, \$ n=57.

All pairwise differences between the three Gina control categories regarding median values of C-ACT scores were significant (all  $p < 0.002$ ). The same applied to the ACT scores.

Cut-off point	Sensitivity (%)	Specificity (%)	PPV (%)	NPV (%)	% Correctly Classified
≤ 17	18	100	100	55	59
≤ 18	23	100	100	57	62
≤ 19	33	100	100	60	67
≤ 20	49	97	95	66	73
≤ 21	62	95	92	71	78
≤ 22	82	85	84	83	83
≤ 23	85	64	70	81	74
≤ 24	97	44	63	94	71

Table 4: performance of the C-ACT with various cut-off points as compared with GINA criteria for assessing uncontrolled asthma in 78 patients aged 4 – 11 years. PPV: positive predicting value. NPV: negative predicting value.

Cut-off point	Sensitivity (%)	Specificity (%)	PPV (%)	NPV (%)	% Correctly Classified
≤ 17	45	100	100	64	72
≤ 18	55	100	100	68	77
≤ 19	66	100	100	74	82
≤ 20	76	96	96	79	86
≤ 21	86	75	78	84	81
≤ 22	90	64	72	86	77
≤ 23	93	54	68	88	74
≤ 24	100	21	57	100	61

Table 5: performance of the ACT with various cut-off points as compared with GINA criteria for assessing uncontrolled asthma in 57 patients aged 12 – 18 years. PPV: positive predicting value. NPV: negative predicting value.

Figure 1




Figure 2

