



Smoke exposure as a determinant of autoantibody titre in α_1 -antitrypsin deficiency and COPD

A.M. Wood*, P. de Pablo*, C.D. Buckley*, A. Ahmad* and R.A. Stockley#

ABSTRACT: Liberation of elastin peptides from damaged lung may be a mechanism of autoimmune lung disease. Citrullination, and anti-citrullinated protein antibody formation occurs in smokers, but the role of smoking in autoantibody generation relevant to pulmonary disease is unclear.

Anti-elastin, anti-cyclic citrullinated peptide (anti-CCP) and anti-mutated citrullinated vimentin (anti-MCV) antibodies were measured in 257 subjects with α_1 -antitrypsin deficiency (AATD), 113 subjects with usual chronic obstructive pulmonary disease (COPD) and 22 healthy nonsmokers. Levels were compared between groups, against phenotypic features and against smoke exposure.

Anti-elastin antibodies were higher in controls relative to AATD ($p=0.008$) and usual COPD ($p<0.00001$), and in AATD relative to usual COPD ($p<0.00001$). Anti-elastin levels showed a threshold at 10 pack-yrs, being higher in those who had smoked less ($p=0.004$). No relationships between antibody levels and clinical phenotype were seen after adjustment for smoke exposure. Anti-CCP antibodies were higher in usual COPD than AATD ($p=0.002$) but the relationship to smoke exposure was less clear.

Smoke exposure is the main determinant of anti-elastin antibody levels, which fall after 10 pack-yrs. Local antibody complexes may be a better measure of elastin directed autoimmunity than circulating levels.

KEYWORDS: α_1 -Antitrypsin deficiency, autoimmune disease, chronic obstructive pulmonary disease, smoking and health

The protease–antiprotease hypothesis of the pathogenesis of chronic obstructive pulmonary disease (COPD) concerns imbalances between proteases that digest elastin and other components of the extracellular matrix in the lung parenchyma, and antiproteases that protect [1–2]. The origin of this theory comes from the observation that patients with α_1 -antitrypsin deficiency (AATD) develop early onset emphysema [3]. α_1 -Antitrypsin (AAT) is an antiprotease, which acts predominantly to block the action of neutrophil elastase, a protease released by neutrophils. Neutrophil elastase is a serine protease, the first of three classes of protease important in COPD. The remaining two classes are the cysteine proteases, such as cathepsin-B, and the matrix metalloproteases (MMPs) [4]. In general, the serine and cysteine proteases are capable of degrading elastin and some forms of collagen [4], whilst MMPs have more of an effect on collagen, gelatin and laminin [2], all of which are components of the extracellular matrix of the lung.

Breakdown of the extracellular matrix, particularly elastin, in the lung is a key feature of both

COPD and AATD, as shown by the presence of high levels of elastin breakdown products. A desmosine cross-link is unique to elastin and may be used as a marker of its degradation [5]. Desmosine and elastin peptides are elevated in the plasma, urine and sputum of COPD patients [6], whilst urinary levels positively correlate with the annual rate of decline in forced expiratory volume in 1 s (FEV₁) in smokers [7]. In general, elastin breakdown is greater in subjects with AATD at a given level of smoking because of a relative excess of neutrophil elastase activity in the lung [8], confirmed by elevated desmosine levels compared with usual COPD subjects [6]. Elastin breakdown has been proposed as a mechanism for generation of autoantibodies directed against elastin, which could in theory perpetuate and/or aggravate lung destruction [9]. In both AATD and COPD, the main stimulus to elastin breakdown is cigarette smoking, but studies of anti-elastin antibody prevalence in these conditions to date have been too small to assess smoke effects on antibody levels adequately [9–11]. Although the original report of circulating anti-elastin antibodies in COPD was

AFFILIATIONS

*College of Medical and Dental Sciences, University of Birmingham, and

#Lung Investigation Unit, University Hospitals Birmingham, Birmingham, UK.

CORRESPONDENCE

A.M. Wood
College of Medical and Dental Sciences
University of Birmingham
Birmingham
B15 2TT
UK
E-mail: a.m.wood@bham.ac.uk

Received:

March 03 2010

Accepted after revision:

May 04 2010

First published online:

June 04 2010

European Respiratory Journal

Print ISSN 0903-1936

Online ISSN 1399-3003

able to show antibody generation by lung tissue and a correlation between peripheral blood responses to elastin peptides and severity of emphysema on computed tomography (CT) scan [9], the effect of anti-elastin antibodies in the lung remained uncertain.

It is also interesting to note that smoking in and of itself is now of interest as a risk factor for autoimmunity, as discussed in a recent review [12]. In some autoimmune diseases this has been noted at epidemiological level, and there is now biological evidence for direct smoke effects on antibodies diagnostic of rheumatoid arthritis, known as anti-citrullinated protein antibodies (ACPA). Citrullination is a post-translational modification of proteins, which occurs in inflammatory disease [13]. It is recognised to occur in the lung of smokers, and is thought to influence occurrence of antibodies directed against citrullinated proteins [14]. Two main ACPAs are recognised, anti-cyclic citrullinated peptide (anti-CCP) and anti-mutated citrullinated vimentin (anti-MCV) antibodies. Both are markers of disease severity in rheumatoid arthritis [15, 16] and there is some suggestion that they occur more in extra-articular disease [17], particularly lung disease [18]. We reasoned that pulmonary citrullination might be a relevant process generating autoantibodies in COPD, alongside elastin liberation.

We hypothesised that anti-elastin and ACPAs would be higher in subjects who had smoked more, and that anti-elastin antibodies would be higher in AATD than COPD due to excessive elastin breakdown.

METHODS

Subjects and clinical phenotyping

AATD

257 PiZZ subjects from the UK national registry for AATD were studied, this being the number with a baseline serum sample available at the time of the study. The project was approved by the local research ethics committee and all subjects gave informed consent. All AATD subjects had a full clinical history (including smoke exposure), a serum AAT level of $<11 \mu\text{M}$, PiZZ genotype confirmed by specific PCR (Hereditab, Salt Lake City, UT, USA) and underwent full pulmonary function testing, as described previously [19]. None had ever received AAT replacement therapy.

COPD

The West Midlands COPD Collection (WMCC) comprises subjects with COPD, excluding those with AATD, defined as FEV₁/forced vital capacity (FVC) <0.7 and post-bronchodilator FEV₁ $<80\%$ predicted, recruited in the central region of the UK. Studies of clinical phenotype and genetic risk in the WMCC have been approved by the local ethics committee. All subjects in the WMCC have undergone pulmonary function testing and clinical history gathering in the same manner as the AATD group. At the time of data collection there were 116 subjects registered with the WMCC, all of whom had given informed consent.

Healthy nonsmokers

The control group consisted of 22 healthy nonsmokers with normal pulmonary function tests, in whom AATD had been excluded, a stored serum sample available and who had consented to participate in research.

Antibody levels

Anti-elastin antibodies were measured in duplicate in serum using methods published by LEE *et al.* [9] on all AATD and COPD subjects. The original publication measured antibodies in plasma; thus the validity of the assay in serum was confirmed by measurement of antibodies in both serum and plasma samples taken at the same time point from 10 individuals; titres were no different between the two sample types ($p=0.841$). Serial dilutions of a pooled serum sample were used to make a standard curve and $\text{IU}\cdot\text{mL}^{-1}$ for subject samples calculated from this. The inter-assay coefficient of variation over the working range was 8.30% ($n=20$ repeats in a single subject). The spike recovery was 87.34%, and assay specificity for anti-human immunoglobulin (Ig)G was confirmed by a zero reading when control antibody was used.

ACPA antibodies were measured in serum or plasma using commercial ELISA kits (valid for use in both sample types) for anti-CCP and anti-MCV antibodies (Phadia UniCAP system; Phadia Ltd, Milton Keynes, UK and Orgentec Diagnostika GmbH, Mainz, Germany, respectively). These are clinically validated tests with a threshold for positivity of $10 \text{IU}\cdot\text{mL}^{-1}$ and $20 \text{IU}\cdot\text{mL}^{-1}$, respectively.

Statistical analysis

Data were analysed using SPSS version 16.0 (SPSS Inc., Chicago, IL, USA). Comparisons of antibody levels between groups were carried out using the Mann–Whitney and Kruskal–Wallis tests. Comparisons of antibody levels within each group were first analysed as univariate models, using Pearson's and Spearman's rank correlations, as dictated by the distribution of the data, for continuous variables, the unpaired t-test and Mann–Whitney tests for binary outcomes, and the Chi-squared test for frequency variables. Adjustments for smoke exposure were made using coefficients from disease group regression models seeking predictors of antibody levels.

RESULTS

Characteristics of the subjects

Characteristics of the three groups are shown in table 1. COPD subjects were generally older and had smoked more than those with AATD (both $p<0.001$), consistent with known features of the conditions. There were more current smokers ($p<0.001$), gas transfer was lower ($p=0.02$), and chronic bronchitis more common ($p<0.001$) in the usual COPD group than those with AATD. Tests of significance regarding the difference in emphysema frequency between AATD and COPD were considered invalid in view of the differences in proportion scanned.

Anti-elastin antibody levels and associations

Anti-elastin antibodies were detected in all 257 AATD samples, all 22 healthy subjects and 113 out of 116 COPD subjects. The mean level was significantly higher in healthy subjects than those with COPD ($p=0.00001$), and those with AATD ($p=0.008$). Levels in AATD were higher than in usual COPD ($p<0.00001$), illustrated in figure 1. Levels in AATD were higher in those within this group who did not have COPD ($p=0.032$; fig. 2).

Within AATD anti-elastin antibodies showed a negative correlation with pack-yrs smoked ($r=-0.199$, $p=0.001$), although visually the relationship did not appear linear

TABLE 1 Characteristics of the disease cohorts

	AATD	COPD	Healthy
Subjects n	257	116	22
Age yrs	50.71 ± 0.70	68.83 ± 0.93	47.07 ± 1.64
Male sex	146 (56.8)	64 (55.17)	7 (31.82)
Pack-yrs smoked	15.30 ± 0.91	46.31 ± 2.69	0
Current smoker	13 (5.06)	32 (27.58)	0
Never smoked	64 (24.90)	0	22
FEV₁ post BD L	1.93 ± 0.07	1.89 ± 0.86	4.56 ± 0.33
FEV₁ % predicted	57.67 ± 2.11	43.43 ± 1.66	101.40 ± 7.59
FEV₁/FVC post BD	0.46 ± 0.01	0.37 ± 0.20	0.74 ± 0.01
Kco mmol·min⁻¹·kPa·L⁻¹	1.12 ± 0.03	1.09 ± 0.86	1.74 ± 0.09
Kco % predicted	71.35 ± 1.34	56.58 ± 4.21	97.50 ± 0.71
COPD	208 (80.93)	116 (100)	0
Emphysema	158 (61.47)	38 (32.76) [#]	Not scanned
Chronic bronchitis	80 (31.50)	68 (58.62)	0
Anti-elastin antibody IU·mL⁻¹	31.18 ± 5.39	5.90 ± 0.88	68.04 ± 11.98
Anti-CCP antibody IU·mL⁻¹	2.68 ± 0.49	18.01 ± 8.13	1.34 ± 0.76
Anti-MCV antibody IU·mL⁻¹	23.57 ± 5.04	39.12 ± 12.09	15.87 ± 6.92
Positive anti-CCP >10 IU·mL⁻¹	8 ± 3.2	6 ± 5.2	0
Positive anti-MCV >20 IU·mL⁻¹	34 ± 13.4	20 ± 17.7	2 ± 9.1

Data are presented as mean ± SE or n (%), unless otherwise stated. AATD: α₁-antitrypsin deficiency; COPD: chronic obstructive pulmonary disease; FEV₁: forced expiratory volume in 1 s; FVC: forced vital capacity; BD: bronchodilator; Kco: transfer coefficient of the lung for carbon monoxide; CCP: cyclic citrullinated peptide; MCV: mutated citrullinated vimentin. [#]: proportionally fewer subjects in the usual COPD group had undergone a computed tomography scan, when the percentage is calculated as a proportion of those scanned it rises to 88.40%.

(fig. 3). We therefore stratified into categories by pack-yrs smoked, and showed a clear threshold at 10 pack-yrs, above which antibodies were lower (fig. 3, p=0.004). Consistent with this, the mean level in never smokers was significantly higher




FIGURE 1. Mean anti-elastin antibody levels stratified by group. The chart shows the mean ± SE of circulating antibody concentration in each group. Healthy subjects exhibited significantly higher concentrations than those with chronic obstructive pulmonary disease (COPD) (p=0.00001), and α₁-antitrypsin deficiency (AATD) (p=0.008). AATD subjects also exhibited higher concentrations than usual COPD subjects (p<0.00001). The values shown are unadjusted for smoke exposure.




FIGURE 2. Anti-elastin antibody concentrations are higher in those with normal lung function in α₁-antitrypsin deficiency (AATD). The chart shows the relationship between forced expiratory volume in 1 s (FEV₁) and anti-elastin levels in AATD which is significant; those with FEV₁ >80% predicted and no evidence of airflow obstruction had higher levels (p=0.032).

than the smoke exposed (51.14 ± 13.05 versus 27.04 ± 5.21; p=0.04). A concomitant positive correlation with FEV₁ was seen (r=0.189, p=0.005) which was lost after adjustment for pack-yrs smoked (r=0.084, p=0.251). A trend towards higher levels in those without emphysema (51.12 ± 17.33 versus 29.30 ± 6.06; p=0.074) was also lost after similar adjustments (p=0.622). The unadjusted correlations in AATD appeared to be driven by never smokers, and indeed in usual COPD, where all individuals had smoked, no relationship between antibody level and pack-yrs or FEV₁ was seen (both p>0.25). Emphysema analyses were not performed in usual COPD because of the low proportion of CT scans available.

Anti-elastin antibody levels were then compared between groups after adjustment for pack-yrs smoked; the difference between AATD and usual COPD was lost (p=0.258). This lack of relationship remained when including only AATD individuals who had spirometrically defined COPD (p=0.131). All other between group comparisons of antibody levels became nonsignificant after adjustment for smoke exposure (all p>0.6). Never smokers with AATD and no evidence of COPD (n=24) were then compared to healthy never smokers without AATD (n=22) in order to ascertain if AATD itself were a significant determinant of antibody levels after control for smoke exposure and the presence of COPD; there was no difference in their anti-elastin antibody levels (p=0.295).

Current versus past smoking did not show a significant difference in antibody levels in either group (both p>0.5). Years since last cigarette exposure did not relate to antibody levels, nor did age at commencement of smoking (both p>0.5). No sex or age relationships to antibody levels were seen in AATD or usual COPD, and sub-stratification by sex did not change the relationship with pack-yrs seen (r= -0.208, p=0.014 in males and r= -0.255, p=0.018 in females).

ACPA associations

Anti-CCP antibodies were higher in usual COPD than AATD (p=0.002, fig. 4), and exceeded the conventional cut-off for a positive level in eight AATD subjects and six of those with




FIGURE 3. Anti-elastin antibody concentrations relate to smoke exposure in α_1 -antitrypsin deficiency. a) The scatterplot shows the relationship between smoke exposure and anti-elastin antibody titre. Whilst it is statistically significant as a linear relationship ($r = -0.199$, $p = 0.001$), visually it appeared to be driven by low level smoking. b) Smoke exposure was divided into categories according to pack-yr smoked. Anti-elastin antibodies then appeared to exhibit a threshold level at 10 pack-yr. c) Comparisons of antibody levels were made using 10 pack-yr to stratify the population. Antibody levels were higher in those who have smoked <10 pack-yr ($p = 0.004$).

usual COPD. Anti-MCV antibodies were positive in 34 AATD subjects and 20 with usual COPD; the level did not differ between groups ($p = 0.876$, fig. 4). There was no difference in either ACPA between AATD and controls ($p = 0.412$). In AATD, neither ACPA showed any relationship to FEV₁ (anti-CCP: $p = 0.608$ for high levels, $p = 0.868$ as a continuous variable; anti-MCV: $p = 0.710$ and $p = 0.864$, respectively).




FIGURE 4. Anti-citrullinated protein antibody (ACPA) concentrations stratified by group. The chart shows the mean anti-cyclic citrullinated peptide (anti-CCP) and anti-mutated citrullinated vimentin (MCV) concentration in the three groups studied. Anti-CCP titre was significantly higher in usual chronic obstructive pulmonary disease (COPD) than α_1 -antitrypsin deficiency (AATD) ($p = 0.002$) but no other significant differences were seen. ■: anti-CCP; ■: anti-MCV.

Equally, in usual COPD, neither ACPA was independently associated with FEV₁ (anti-CCP: $p = 0.424$ and 0.969 ; anti-MCV: $p = 0.957$ and 0.406). No age or sex associations were seen in either group (all $p > 0.5$). In sex sub-stratified analyses the lack of relationship with FEV₁ persisted (all $p > 0.6$).

Neither anti-CCP nor anti-MCV antibodies differed between those who had never smoked and those who had been smoke exposed in AATD ($p = 0.903$), a test which could not be performed in usual COPD since all subjects had smoked. No linear dose–response relationship (pack-yr *versus* ACPA) existed in AATD (anti-CCP: $r = 0.002$, $p = 0.979$; anti-MCV: $r = -0.069$, $p = 0.301$) or usual COPD (anti-CCP: $r = -0.158$, $p = 0.161$; anti-MCV: $r = -0.024$, $p = 0.864$). However, when split into pack-yr categories in AATD, as before, a threshold was apparent at 10 pack-yr, although less clear than that seen for anti-elastin antibodies (fig. 5). However, the difference in titre between those either side of the 10 pack-yr threshold was not significant (anti-CCP, $p = 0.661$; anti-MCV, $p = 0.110$), and the incidence of positive ACPA did not differ above and below this threshold ($p = 0.761$ and 0.503 , respectively). Given that no relationship with smoke was observed further inter-group comparisons after adjustment for smoke exposure were not performed.

Since ACPA are diagnostic of rheumatoid arthritis, the incidence of this, and other autoimmune diseases, was determined in all those with a positive ACPA. Of the 36 AATD subjects with at least one positive ACPA, none had rheumatoid arthritis either at baseline or during follow-up (mean duration of follow-up since serum collection was 9 yrs). However, four had autoimmune thyroid disease at baseline, one had ulcerative colitis at baseline and a further individual developed ulcerative colitis during follow-up. Amongst the 22 ACPA positive individuals with COPD three had rheumatoid arthritis. To date, none has completed 1 yr of follow-up.

DISCUSSION

The data presented here show a clear relationship between circulating anti-elastin antibody levels and smoke exposure,




FIGURE 5. Anti-citrullinated protein antibody (ACPA) relationships to smoke exposure in α_1 -antitrypsin deficiency. a and b show ACPA levels stratified by smoke exposure categories. Although a threshold was suggested visually at 10 pack-yr this was not significant (anti-cyclic citrullinated peptide (anti-CCP), $p=0.661$ and anti-mutated citrullinated vimentin (anti-MCV), $p=0.110$).

and no relationship between these antibodies and clinical phenotype after smoking is taken into account. Conversely ACPA, which are theoretically generated after smoke exposure, did not show any significant relationship to smoke exposure in our cohorts, and less marked differences between groups.

One study has previously reported high anti-elastin antibody levels in COPD [9], which we were unable to confirm. This study concerned a group with emphysema, thus it could be argued that our analyses should have been restricted to those with CT-proven emphysema if we wished to confirm their findings. We did not exclude subjects who had not been scanned or who did not have emphysema, as our primary objective was to compare levels between groups whose elastin breakdown might be expected to differ [6], and against smoke exposure, not to reproduce the work of LEE *et al.* [9]. However, secondary analyses comparing antibody titre between individuals with and without emphysema in AATD did not reveal a significant difference. It should be noted that differences in titre seen by LEE *et al.* [9] between groups were small, and that this observation was a small part of their study, which examined anti-elastin autoimmunity more specifically, using lung tissue to assess anti-elastin antibody complexes, and T-cell responses to elastin peptides. Our results, in direct contrast

to theirs, showed anti-elastin antibodies were higher in those with better lung function and in healthy individuals. This implies that the presence and level of circulating anti-elastin antibody is not a key feature of COPD, an observation recently reported in another study [11], and may even fall as disease progresses. We hypothesise that this might be due to ongoing, or increasing, antibody consumption by binding in the lung. This is consistent with the increased antibodies seen in lung tissue by LEE *et al.* [9], although longitudinal studies involving both lung tissue and circulating samples would be required to definitively answer this question. Antibody levels do not rise with cessation of smoke exposure, consistent with the perpetuation of inflammation and lung damage in COPD seen after this time, with measurable disease progression in many subjects [20].

None of the studies of anti-elastin antibodies to date has taken effects of smoking into consideration in the analyses pertaining to antibody levels, despite the theory that anti-elastin antibody generation occurs secondary to smoke-induced elastin liberation in the lung [9]. The data presented here show a clear difference in anti-elastin antibody levels between smokers and those who have never smoked in AATD, and a relationship between pack-yr smoked and antibody levels. There was also a threshold level of smoke exposure apparent, a feature identified in another large cohort of AATD subjects recently, which contributed to their predictive model for FEV₁ [21]. Smoke exposure was common in AATD, higher in usual COPD, and absent in controls, thus between-group comparisons of antibody levels, which appeared highly significant initially, required adjustment for this exposure. After such adjustment, not only was the relationship with FEV₁ lost within AATD, but also all inter-group comparisons ceased to be significant. This, together with the recent suggestion that smoking may be a risk factor for autoimmunity [12], is important to note in future studies of the prevalence of other autoantibodies in COPD and control populations. Subjects with AATD may be more sensitive to other environmental influences on lung function, as shown by relationships between outdoor air pollution and lung function in this group [22]. We surmised that differences in elastin breakdown and subsequent anti-elastin generation might still occur between nonsmokers with and without AATD for this reason. This was not confirmed, perhaps because the effect sizes of agents other than cigarette smoke are small in both groups [22, 23].

Although differences in ACPA titre were seen between COPD, AATD and controls, the differences were less marked than with anti-elastin antibodies and less clearly related to smoke exposure. This is the first study, to our knowledge, of these antibodies in lung disease, and our findings will require further validation; nevertheless, it is interesting to consider their significance. The pathogenic role of ACPA in rheumatoid arthritis is thought to involve complement activation [24] and alteration of the T-cell repertoire in the joint [25], with reaction to self as a result. Our data would suggest that this may be organ specific, occurring just in the joint despite the fact that any protein containing citrulline could be a target for ACPA. Alternatively the effect on synovium could depend upon a further genetic or environmental influence, given that ACPA positive individuals here did not always exhibit rheumatoid arthritis (or indeed other autoimmune diseases). Shared

epitope alleles, contained within class II of the human leukocyte antigen system (*HLA-DRB1**04, *01 and *10), are important in determining risk of rheumatoid arthritis [26] and ACPA status [27]. Recent data suggest that smoking is a key feature of determining ACPA-positive disease risk in the presence of these alleles [27, 28], consistent with this hypothesis. ACPA may occur up to 10 yrs before clinical features of rheumatoid arthritis appear [29], the titre relating inversely to the time until diagnosis [30], such that highly positive individuals are likely to be diagnosed soon. It will therefore be of importance to follow up positive individuals in the cohort, particularly since the incidence of positive anti-CCP antibodies exceeded that reported in healthy controls [29] five-fold in AATD and eight-fold in usual COPD. The majority of AATD subjects have, however, been followed up for >8 yrs since their baseline assessment and blood sampling, making it unlikely that many new cases will be found in this group. In the usual COPD cohort, high ACPA levels were more common than in AATD, and more common in both groups than in the general population. Intuitively, the role of smoking in determining pulmonary citrullination [14], and subsequent antibody formation, might have been thought responsible. Again, however, our data suggested that in AATD ACPA levels were lower in heavier smokers, albeit not statistically different. Since there was no significant dose-response effect adjustments of ACPA level for lifetime lack of smoke exposure, analyses could not be carried out to determine if differences between AATD, COPD and healthy individuals persisted after adjusting for this feature. Larger cohorts may be required for further clarification of this issue. Further study of citrullination in smokers is also warranted to clarify the mechanism of antibody generation and relationship to ACPA suggested here.

Autoimmunity is a current topic of debate in COPD pathogenesis, since some characteristic features of autoimmunity are present. First, a lymphocytic infiltrate in the lung occurs, with increasing pulmonary CD4+ and CD8+ T-cell numbers relating to worsening disease [31, 32]. Autoantibodies to pulmonary epithelium have also been demonstrated [33]. Furthermore, oligoclonal T-cell populations have been demonstrated in the lung, suggesting accumulation due to antigenic stimulation [34]. Evidence from animal models also supports a role for autoimmunity in COPD. Specifically, the transfer of pathogenic CD4+ T-cells from rats with emphysema and anti-endothelial cell antibodies results in the development of emphysema in healthy animals [35], and mice injected with elastin peptides develop both anti-elastin antibodies and a bronchoalveolar lavage fluid profile similar to that seen in COPD [36, 37]. The initial report of elevated anti-elastin antibodies by LEE *et al.* [9] has now been refuted by two small studies (in addition to the data presented here), the first in a mixed-disease cohort [10], and the second in pure-disease cohorts, but with small numbers in each [11]. Indeed, in the latter study the authors noted that antibody levels were higher in healthy subjects, consistent with the data we report here. Autoimmune processes are generally thought to be more common in female subjects, hence we undertook some sex-specific analyses but again no differences were seen. Taken together, the results suggest that anti-elastin antibodies are unlikely to be a useful disease marker in COPD and, if autoimmunity exists directed against elastin, circulating anti-elastin

antibody titres may be of less importance than local levels in lung tissue.

Our study is weakened by the absence of a healthy smoking group for comparison to the usual COPD group, although comparisons within AATD of healthy individuals and those with lung disease, controlled for smoking, act as a suitable surrogate for this. We also have not tested for IgM, IgA and other antibody subtypes for anti-elastin. This feature was investigated by a previous group, whose findings concurred with ours in terms of between-group comparisons, such that we felt it was unnecessary to repeat their experiments [11].

In summary, our data show a clear relationship between smoke exposure and anti-elastin antibody levels, and no difference between AATD, COPD and healthy individuals after adjustment for this feature. This suggests that circulating anti-elastin levels cannot be used as a disease marker and that future study of the autoimmunity hypothesis may need to be directed to antibody complexes in lung tissue rather than circulating antibodies.

SUPPORT STATEMENT

We would like to thank Talecris Biotherapeutics, whose unrestricted grant to R.A. Stockley funded this work.

STATEMENT OF INTEREST

A statement of interest for R.A. Stockley and the study itself can be found at www.ers.ersjournals.com/site/misc/statements.xhtml

ACKNOWLEDGEMENTS

We would like to thank all members of the UK α_1 -antitrypsin deficiency registry and West Midlands COPD Collection investigating teams for gathering of clinical data and biological samples, and D. DeMeo (Brigham and Women's Hospital, Boston, MA, USA) for her advice on statistical analysis.

REFERENCES

- 1 Stockley RA. Neutrophils and protease/antiprotease imbalance. *Am J Respir Crit Care Med* 1999; 160: S49–S52.
- 2 Shapiro SD, Senior RM. Matrix metalloproteinases. Matrix degradation and more. *Am J Respir Cell Mol Biol* 1999; 20: 1100–1102.
- 3 Eriksson S. Studies in alpha 1 antitrypsin deficiency. *Acta MedScand Suppl.* 1965; 432: 1–85.
- 4 Barnes PJ, Shapiro SD, Pauwels RA. Chronic obstructive pulmonary disease: molecular and cellular mechanisms. *Eur Respir J* 2003; 22: 672–688.
- 5 Harel S JA, Yu SY, Hurewitz A, *et al.* Desmosine radioimmunoassay for measuring elastin degradation *in vitro*. *Rev Respir Dis* 1980; 122: 769–773.
- 6 Ma S, Lin YY, Turino GM. Measurements of desmosine and isodesmosine by mass spectrometry in COPD. *Chest* 2007; 131: 1363–1371.
- 7 Gottlieb DJ SP, Sparrow D, Gale ME, *et al.* Urinary desmosine excretion in smokers with and without rapid decline of lung function: the normal aging study. *Am J Respir Crit Care Med* 1996; 154: 1290–1295.
- 8 Hill AT, Bayley DL, Campbell EJ, *et al.* Airways inflammation in chronic bronchitis: the effects of smoking and alpha1-antitrypsin deficiency. *Eur Respir J* 2000; 15: 886–890.
- 9 Lee SH, Goswami S, Grudo A, *et al.* Anti-elastin autoimmunity in tobacco smoking-induced emphysema. *Nature Med* 2007; 13: 567–569.

- 10 Cottin V, Fabien N, Khouatra C, *et al.* Anti-elastin autoantibodies are not present in combined pulmonary fibrosis and emphysema. *Eur Respir J* 2009; 33: 219–221.
- 11 Greene CM, Low TB, O'Neill SJ, *et al.* Anti-PGP or anti-elastin autoantibodies are not evident in chronic inflammatory lung disease. *Am J Respir Crit Care Med* 2009; 181: 31–35.
- 12 Arnson Y, Shoenfeld Y, Amital H. Effects of tobacco smoke on immunity, inflammation and autoimmunity. *J Autoimmun* 2010; 35: J258–J265.
- 13 Makrygiannakis D, af Klint E, Lundberg IE, *et al.* Citrullination is an inflammation-dependent process. *Ann Rheum Dis* 2006; 65: 1219–1222.
- 14 Makrygiannakis D, Hermansson M, Ulfgren A-K, *et al.* Smoking increases peptidylarginine deaminase 2 enzyme expression in human lungs and increases citrullination in BAL cells. *Ann Rheum Dis* 2008; 67: 1488–1492.
- 15 Mewar D, Coote A, Moore DJ, *et al.* Independent associations of anti-cyclic citrullinated peptide antibodies and rheumatoid factor with radiographic severity of rheumatoid arthritis. *Arthritis Res Ther* 2006; 8: R128.
- 16 Wagner E, Skoumal M, Bayer PM, *et al.* Antibody against mutated citrullinated vimentin: a new sensitive marker in the diagnosis of rheumatoid arthritis. *Rheumatol Int* 2009; 29: 1315–1321.
- 17 Korkmaz C, Us T, Tasifoglu T, *et al.* Anti cyclic citrullinated peptide (CCP) antibodies in patients with long standing rheumatoid arthritis and their relationship with extra articular manifestations. *Clin Biochem* 2006; 39: 961–965.
- 18 Alexiou I, Germenis A, Koutroumpos A, *et al.* Anti cyclic citrullinated peptide 2 (CCP2) autoantibodies and extra articular manifestations in Greek patients with rheumatoid arthritis. *Clin Rheumatol* 2008; 27: 511–513.
- 19 Dowson LJ, Guest PJ, Hill SL, *et al.* High-resolution computed tomography scanning in alpha1-antitrypsin deficiency: relationship to lung function and health status. *Eur Respir J* 2001; 17: 1097–1104.
- 20 Celli BR, Thomas NE, Anderson JA, *et al.* Effect of pharmacotherapy on rate of decline of lung function in chronic obstructive pulmonary disease: results from the TORCH study. *Am J Respir Crit Care Med* 2008; 178: 332–338.
- 21 Castaldi PJ, DeMeo D, Kent DM, *et al.* Development of predictive models for airflow obstruction in alpha 1 antitrypsin deficiency. *Am J Epidemiol* 2009; 170: 1005–1013.
- 22 Wood AM, Harrison RM, Semple S, *et al.* Outdoor air pollution is associated with disease severity in alpha1-antitrypsin deficiency. *Eur Respir J* 2009; 34: 346–353.
- 23 Downs SH, Schindler C, Liu LJ, *et al.* Reduced exposure to PM10 and attenuated age-related decline in lung function. *N Engl J Med* 2007; 357: 2338–2347.
- 24 Trouw LA, Haisma EM, Levahrt EW, *et al.* Anti cyclic citrullinated antibodies from rheumatoid arthritis patients activate complement via both the classical and alternate pathways. *Arthritis Rheum* 2009; 60: 1923–1931.
- 25 Cantaert T, Brouard S, Thurlings RM, *et al.* Alterations of the synovial T cell repertoire in anti citrullinated protein antibody positive rheumatoid arthritis. *Arthritis Rheum* 2009; 60: 1944–1956.
- 26 Gregersen PK, Silver J, Winchester RJ. The shared epitope hypothesis. An approach to understanding the molecular genetics of susceptibility to rheumatoid arthritis. *Arthritis Rheum* 1987; 30: 1205–1213.
- 27 Lundstrom E, Kallberg H, Alfredsson L, *et al.* Gene–environment interaction between the DRB1 shared epitope and smoking in the risk of anti citrullinated protein antibody positive rheumatoid arthritis. *Arthritis Rheum* 2009; 60: 1597–1603.
- 28 Mahdi H, Fisher BA, Kallberg H, *et al.* Specific interaction between genotype, smoking and autoimmunity to citrullinated alpha enolase in the etiology of rheumatoid arthritis. *Nat Genet* 2009; 41: 1319–1324.
- 29 Nielen MM, van Schaardenburg D, Reesink HW, *et al.* Specific autoantibodies precede the symptoms of rheumatoid arthritis: a study of serial measurements in blood donors. *Arthritis Rheum* 2004; 50: 380–386.
- 30 Chibnik LB, Mandl LA, Costenbader KH, *et al.* Comparison of threshold cut points and continuous measures of anti cyclic citrullinated peptide antibodies in predicting future rheumatoid arthritis. *J Rheumatol* 2009; 36: 706–711.
- 31 Hogg JC, Chu F, Utokaparch S, *et al.* The nature of small-airway obstruction in chronic obstructive pulmonary disease. *N Engl J Med* 2004; 350: 2645–2653.
- 32 Grumelli S, Corry DB, Song LZ, *et al.* An immune basis for lung parenchymal destruction in chronic obstructive pulmonary disease and emphysema. *PLoS Med* 2004; 1: e8.
- 33 Feghali-Bostwick CA, Gadgil AS, Otterbein LE, *et al.* Autoantibodies in patients with chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2008; 177: 156–163.
- 34 Sullivan AK, Simonian PL, Falta MT, *et al.* Oligoclonal CD4+ T cells in the lungs of patients with severe emphysema. *Am J Respir Crit Care Med* 2005; 172: 590–596.
- 35 Taraseviciene-Stewart L, Scerbavicius R, Choe KH, *et al.* An animal model of autoimmune emphysema. *Am J Respir Crit Care Med* 2005; 71: 734–742.
- 36 Goswami S, Barranco W, Lee SH, *et al.* An autoimmune basis in the disease pathogenesis of COPD/emphysema. *J Immunol* 2007; 178: 130.41.
- 37 Finlay GA, Russell KJ, McMahon KJ, *et al.* Elevated levels of matrix metalloproteinases in bronchoalveolar lavage fluid of emphysematous patients. *Thorax* 1997; 52: 502–506.