

# European Respiratory Society Annual Congress 2012

**Abstract Number:** 2479

**Publication Number:** 3105

**Abstract Group:** 6.3. Tobacco, Smoking Control and Health Education

**Keyword 1:** Smoking **Keyword 2:** Adolescents **Keyword 3:** Public health

**Title:** Smoking rate in 12-15 years old Parisian school is sensitive to political decisions

Bertrand 13183 Dautzenberg bdautz@gmail.com MD <sup>1,4,5</sup>, Maryvonne 13184 Noel maryvonne2@free.fr <sup>2</sup>, Pierre 13185 Birkui dirkui@aliceadsl.fr MD <sup>1</sup> and Marie-Dominique 13186 Dautzenberg mddautz@gmail.com MD <sup>1,3,5</sup>. <sup>1</sup> Association, Paris Sans Tabac, Paris, France, 75010 ; <sup>2</sup> Infirmière Conseillère, Rectorat AcAdémie de Paris, France, 75019 Paris ; <sup>3</sup> Consultation Tabacologie, Hôpital Necker-Enfants Malades, Paris, France, 75015 Paris ; <sup>4</sup> Unité Tabac Service de Pneumologie, GHU Pitié-Salpêtrière-Charles Foix, Paris, France, 75013 Paris and <sup>5</sup> Association, OFT, Paris, France, 75006 Paris .

**Body:** The first cancer plan was launched with force, funded and followed with a strong commitment by the head of state. The second cancer plan included only limited measures against smoking and was accompanied by the cessation of funding and lack of control. Methods: The cross-sectional survey conducted annually PST on 2% of students per class randomly and distributed and collected with the help of the Rector of the Academy of Paris gives an image of smoking among college students, we compare the rate of smoking daily among college students (12-15 years) the 2 years preceding each of the plans (2000-2001 and 2006-2007) in the second and third years of each of the two plans (2010-2011 and 2004-2005). Results: Half the leading cancer plan has helped lower smoking rate of 12.0% to 5.8% (13.6% to 5.7% in girls and 10.3% to 5.9% in boys) to 52% of relative decline. For the entire foreground is the relative decline of 73% with a rate of smoking at the end of plan by 3.2%.

The first phase of second cancer plan has had the opposite effect with a rise in smoking rates by 3.7% to 4.5% for females and 2.6 to 3%, 8% for boys is a relative rise 29% and smoking rates by 4.1%.

Conclusions: A few years apart national tobacco policies can make the best and the worst. Abandoning teenagers recently at their smoking policies hamper the future of a generation, and cutting back the significant benefits of the first cancer plan.