## **European Respiratory Society Annual Congress 2012**

**Abstract Number:** 1622

**Publication Number: P449** 

Abstract Group: 4.2. Sleep and Control of Breathing

Keyword 1: Sleep disorders Keyword 2: Sleep studies Keyword 3: Comorbidities

**Title:** Pulmonary diffusion capacity is not associated with severity of Cheyne-Stokes respiration in heart failure patients

Olaf 8999 Oldenburg ooldenburg@hdz-nrw.de MD <sup>1</sup>, Britta 8995 Körber bkoerber@hdz-nrw.de MD <sup>1</sup>, Thomas 8996 Bitter tbitter@hdz-nrw.de MD <sup>1</sup>, Thomas 8997 Fischbach tfischbach@hdz-nrw.de MD <sup>1</sup> and Dieter 8998 Horstkotte akleemeyer@hdz-nrw.de MD <sup>1</sup>. <sup>1</sup> Department of Cardiology, Heart and Diabetes Center North Rhine-Westphalia, Ruhr University Bochum, Bad Oeynhausen, Germany.

**Body:** A recent study in severe heart failure (HF) patients indicates a possible pathophysiological role of an impaired pulmonary diffusion capacity and respiratory disturbance during sleep, especially the degree of Cheyne-Stokes respiration (CSR). Aim of the present study was to verify this hypothesis in a larger cohort of HF patients. In 87 patients (83 male, age  $68 \pm 9$  years) with polysomnography confirmed and untreated CSR due to cardiac failure (NYHA  $\geq$  II, LV-EF  $\leq$ 45%), pulmonary diffusion capacity (TLCO and KCO) were prospectively investigated using the single-breath method. Apnoea-hypopnoea-index (AHI) was  $38.0 \pm 14$ /h, LV-EF  $34 \pm 7$ %, TLCO  $67 \pm 17$ % of normal and alveolar-volume corrected diffusion capacity (KCO)  $85 \pm 20$ % of normal. Correlation analysis using Spearman rank order correlation revealed no significant correlation of AHI and TLCO (r = 0.071, p = 0.516) or AHI and KCO (0.019, p = 0.863).

In our cohort of HF patients with untreated CSR no correlation of respiratory disturbance during sleep and pulmonary diffusion capacity was found. Thus, a significant role of an impaired pulmonary diffusion on CSR genesis seems to be unlikely.