

Physiological effects of vibration in subjects with cystic fibrosis

Bredge McCarren

School of Physiotherapy

The University of Sydney

Jennifer A Alison

School of Physiotherapy

The University of Sydney

Correspondence to:

Bredge McCarren

School of Physiotherapy

Faculty of Health Sciences

The University of Sydney

PO Box 170

Lidcombe NSW 1825

Australia

Email: B.McCarren@fhs.usyd.edu.au

FAX: 0011 612 93519601

Short title: Physiological effects of vibration

Key words: airway clearance interventions, peak expiratory flow rate, physiotherapy

ABSTRACT

The physiological mechanisms by which vibration and other physiotherapy interventions may clear secretions in subjects with cystic fibrosis are unknown. The main aim of the study was to compare the expiratory flow rates and frequencies of airflow oscillation of vibration to those of Acapella ®, Flutter ®, positive expiratory pressure (PEP) and percussion.

Methods: The respiratory flow rates were measured during the interventions, the order of which was randomised. The oscillation of the airflow of the interventions was determined by frequency spectral analysis.

Results: In 18 young adult subjects with cystic fibrosis, the mean peak expiratory flow rate (PEFR) of vibration was greater than Flutter ($p=0.002$), percussion ($p<0.001$), Acapella ($p<0.001$) and PEP ($p<0.001$). The mean (SD) of the oscillation of the airflow of vibration, 8.4 (0.4) Hz, was lower than Acapella, (13.5 (1.7) Hz; $p<0.001$) and Flutter (11.3 (1.5) Hz; $p<0.001$) but similar to percussion 7.3 (0.3) Hz.

Conclusions: Theoretically, the higher PEFR of vibration compared to the other physiotherapy interventions may promote secretion clearance. In addition, the frequency of oscillation of vibration was within the range demonstrated to increase mucus transport [1]. This study has provided some evidence for the physiological rationale for the use of vibration to aid secretion clearance.

(202 words)

INTRODUCTION

Vibration is a traditional physiotherapy intervention that is used in patients with respiratory disorders. Vibration is the manual application of fine oscillatory movements combined with compression of the patient's chest wall and is commonly used by physiotherapists to assist with secretion clearance [2]. The compression and oscillation applied during vibration are proposed to aid secretion clearance by a number of physiological mechanisms. These include:

1. increasing absolute peak expiratory flow rates to move secretions towards the oropharynx [3];
2. increasing the expiratory bias of airflow to increase the annular flow of mucus towards the oropharynx which occurs when the peak expiratory flow rate (PEFR) is 10% greater than peak inspiratory flow rate (PIFR) ie. $PEFR/PIFR$ ratio >1.1 [4];
3. increasing mucus transport by decreasing the viscosity of mucus and increasing expiratory flow due the effects of oscillation of airflow at frequencies ranging from 3 – 17 Hz [1]; and
4. eliciting spontaneous coughs by the mechanical stimulation of the airways [5].

Other physiotherapy interventions that may aid secretion clearance, incorporate positive pressure or oscillation applied either at the mouth or chest wall. These include the manual techniques of percussion, and the mechanical devices such as positive expiratory pressure (PEP) device, Flutter VRP1® and Acapella ®. These physiotherapy interventions are proposed to aid secretion movement to the central airways, from where they can be expectorated by forced expiratory manoeuvres. Although physiotherapy has been shown to be more effective for secretion clearance than no intervention, there appears to be no

difference between the physiotherapy interventions [6, 7]. In addition, the physiological mechanisms by which many of these physiotherapy interventions aid secretion clearance are unknown. An understanding of physiological mechanisms of these interventions in patients with excessive secretions may aid treatment selection.

Once secretions have reached the central airways, the forced expiratory manoeuvres of cough and huff, which rely on high expiratory flow rates, have been demonstrated to be effective for secretion clearance and sputum expectoration [8, 9, 10]. The expiratory flow rates of these forced expiratory manoeuvres in patients with cystic fibrosis have not been previously reported.

Therefore the main aim of this study in subjects with cystic fibrosis was to compare the physiological effects of vibration to other physiotherapy interventions used for airway clearance and relate these effects to the proposed mechanisms of secretion clearance. A secondary aim was to measure the flow rates of the forced expiratory manoeuvres in patients with cystic fibrosis.

MATERIALS AND METHODS

Subjects

Subjects diagnosed with cystic fibrosis were recruited for the study. The exclusion criteria were the presence of two or more of any of the five signs or symptoms of an acute infection within the previous two weeks ie. fever $>37.5^{\circ}\text{C}$, acute increase in secretion production, increase in shortness of breath, feeling unwell and an increased use of antibiotics.

Procedure

All testing was carried out during one session. The subjects were asked to refrain from performing any respiratory physiotherapy treatment and taking hypertonic saline or pulmonzyme for at least four hours prior to the study [11]. In addition, subjects were asked not to take any other medications (eg. bronchodilators etc) that they would routinely take prior to respiratory physiotherapy. These medications were administered 15 minutes prior to the dynamic spirometry (Vitalograph Compact, Vitalograph Ltd, Buckingham) and the testing session.

All interventions were carried out by the same physiotherapist, who instructed the subjects in the various interventions to ensure consistent application of the interventions. Each intervention was implemented to replicate current clinical practice and was applied or performed three times by each subject. The order for all interventions was randomised using a computer generated list.

The physiotherapy interventions were vibration, percussion, PEP device (PEP) (Astra Tech Denmark), Flutter VRP1 valve® (Flutter) (VRP1 Desitin/Scandipharm Flutter VarioRaw SA) and Acapella PEP ® therapy DH (Acapella) (DHD Healthcare, NY). The forced expiratory manoeuvres were voluntary cough and huff from high lung volumes (huff_{HIGH}). The subjects also performed an inspiration to total lung capacity (TLC) followed by passive expiration (TLC_{relax}) to act as a control manoeuvre to account for the effects of lung recoil on expiratory flow.

Vibration was applied manually to the chest wall during expiration after a slow maximal inspiration described to the subject as a 'big' breath in. The subjects were asked not to

actively expire. Percussion was applied manually during tidal breathing for at least 30 seconds per application. During both vibration and percussion the position of the subjects was side-lying, with the reported most productive side uppermost. If there was no difference in reported sputum production between the left and right lungs, then the side-lying position was randomised. The forces applied to the subject's chest wall during both percussion and vibration were to the greatest tolerated level of comfort.

The breathing manoeuvre during the use of the devices of PEP, Flutter and Acapella was an inspiration slightly deeper than a normal tidal breath with an end-inspiratory pause of 2-3 seconds. Expiration was slightly active but not forceful, lasting for approximately five seconds. During the use of PEP, the resistor selected was one where a positive expiratory pressure between 15-20 cmH₂O [11] for 5 seconds was achieved by each subject. During the use of Flutter, the device was in almost neutral position (ie. horizontal) to maximise the oscillation amplitude and to target the mid frequency range of oscillation of 10-15 Hz [12]. This mid-frequency range is proposed to have maximal effect on mucus transport [1, 13]. The Acapella was used according to manufacturer's recommendations (DHD Healthcare) with the resistance set by adjusting the numerical dial to the minimal level where each subject could exhale for 3-4 seconds while feeling the vibratory effects.

During huff_{HIGH} and cough the subjects were instructed to inspire maximally followed by either a forced expiration with glottis open for huff_{HIGH} or a cough. For TLC_{relax} subjects were instructed to inspire to total lung capacity and expire passively.

The subjects were instructed to inspire as slowly as possible for all interventions (except percussion) to maximise an expiratory bias to airflow. The subjects were sitting upright for

all of the interventions with the exception of vibration and percussion. Between repetitions of an intervention and between each intervention, the subjects were encouraged to rest and perform relaxed tidal breathing until they had returned to the baseline effort of breathing.

Ethical approval for the study was obtained from Central Sydney Area Health Service Ethics Committee and The University of Sydney's Human Research Ethics Committee and informed consent was obtained from each subject.

Measurements

The inspiratory and expiratory flow rates during all interventions were measured via a mouthpiece with a heated pneumotachograph (Hans Rudolph Model 3813, Hans Rudolph Inc, Kansas City, Missouri). Flow signals were integrated to provide volume measurements [14]. The devices of PEP, Flutter and Acapella were attached to the expiratory port of a two-way non-rebreathing valve (2700 series Hans Rudolph Inc, Kansas) which was connected to the pneumotachograph. This enabled inspiration through the pneumotachograph and expiration through the pneumotachograph and the devices. The pneumotachograph was calibrated and used according to recommended guidelines [15]. Data was collected at a sampling frequency of 100Hz. The subjects were asked to score perceived breathing effort of each intervention using the modified Borg scale for breathlessness [16]. During each intervention the number of spontaneous coughs elicited was recorded with an audiotape for later analysis.

Data Analysis

The oscillation frequency of the interventions which had a proposed oscillation component (ie. vibration, percussion, Flutter and Acapella) was determined by a frequency spectral analysis (Fourier Transform of the auto-correlation function) [17] of the flow data (Matlab

6.5, Student Version, Release 13). An analysis of oscillation frequency of cough and huff_{HIGH} was not performed. After each session the number of spontaneous coughs stimulated during each intervention was recorded from the audiotape.

To calculate sample size, a power analysis was performed on the data of the first 14 subjects, to detect a 25% difference in peak expiratory flow rates between vibration and one of the other interventions of Flutter, Acapella, percussion or PEP. A sample size of 12 subjects was required for this repeated measure study to have 80% power with α of 0.05.

In order to minimise the risk of type one error due to too many statistical tests, only the inspiratory volume (V_I), PEFR, cough frequency and oscillation frequency of vibration were compared with the other physiotherapy interventions of percussion, PEP, Flutter and Acapella. An additional analysis, comparing V_I and PEFR of vibration with cough and huff_{HIGH}, was performed. A repeated measures ANOVA and a post hoc Dunnett's test were used. A Friedman ANOVA for repeated measures was used to compare the subjects reported breathing effort during vibration with the physiotherapy interventions of Acapella, Flutter, PEP and percussion as the data were not normally distributed. A one-way ANOVA was used to compare the effect of disease severity on the PEFR of vibration. Disease severity was based on forced expiratory volume in one second (FEV_1) percent predicted where severe lung disease was $FEV_1 \leq 40\%$ predicted, moderate lung disease was $40\% < FEV_1 \leq 70\%$ predicted, mild lung disease was $70\% < FEV_1 < 90\%$ predicted and normal lung function was $FEV_1 \geq 90\%$ predicted [18]. The results report the mean, standard deviation (SD) and SD of means of the triplicate measures of each intervention of each subject. Significance was set at the 0.05 level.

RESULTS

Subjects

Eighteen subjects (7 female) diagnosed with cystic fibrosis volunteered for the study. The mean (SD) age of these subjects was 28.5 (6.2) years with a body mass index of 20.8 (2.8) kg/m². The mean forced expiratory volume in one second (FEV₁) was 55% of predicted, with a FEV₁/forced vital capacity (FVC) ratio of 58%. Six subjects had severe lung disease, eight had a moderate lung disease, one had mild lung disease and three subjects had normal lung function.

Effects of physiotherapy interventions on respiratory flow rates and volumes

The mean PEFR of vibration was 1.4 times greater than Flutter ($p=0.002$), 1.9 times greater than percussion ($p<0.001$), 2.7 times greater than Acapella ($p<0.001$) and 3.6 times greater than PEP ($p<0.001$) (Table 1, Figure 1). Figure 2 shows the time course of expiratory flow of the interventions of one subject. There was no significant effect of disease severity on the PEFR of vibration ($p = 0.17$).




Figure 1 The peak expiratory flow rate (PEFR) of physiotherapy interventions. Data are means and SD of mean data of each subject. Significantly different from vibration at * $p < 0.001$, † $p = 0.002$. PEP = positive expiratory pressure device.




Figure 2 The time course of the expiratory flow of the interventions in one subject. The eight traces have been separated vertically for clarity.

There was no significant difference in inspired volumes between vibration and those of PEP, Flutter and Acapella (Table 1). The inspired volume of vibration was greater than percussion ($p < 0.001$). There was no difference in the number of coughs stimulated between vibration, percussion, Flutter, Acapella and PEP ($p = 0.7$).

Table 1 Effects of physiotherapy interventions on peak flow rates respiratory volumes and stimulation of coughs.

Intervention	PEFR (L/s)	PIFR (L/s)	PEFR/PIFR	V_I (L)	V_E (L)	coughs stimulated	n
vibration	1.58 (0.73)	1.06 (0.27)	1.51	1.78 (0.87)	2.44 (1.06)	0.7 (1.0)	17‡
percussion	0.83 * (0.14)	0.84 (0.10)	0.99	0.91 * (0.37)	1.03 (0.50)	0.5 (0.9)	18
PEP	0.44 * (0.15)	0.96 (0.20)	0.47	1.64 (0.40)	1.96 (0.57)	0.5 (0.6)	18
Flutter	1.13 † (0.30)	1.05 (0.27)	1.15	1.62 (0.52)	1.81 (0.57)	0.4 (0.7)	17‡
Acapella	0.59 * (0.08)	0.98 (0.27)	0.64	1.55 (0.46)	1.68 (0.50)	0.8 (1.0)	18
TLC _{relax}	0.66 (0.16)	1.01 (0.40)	0.73	1.79 (0.66)	2.24 (0.79)	0	15§

Data are means and SD of mean data of each subject. Legend: significantly different from vibration at * $p < 0.001$, † $p = 0.002$; PIFR = peak inspiratory flow rate, V_I = inspired volume, V_E = expired volume, n = number of subjects used in analysis, PEP = positive expiratory pressure device, TLC = total lung capacity; ‡ data lost due to technical difficulties (data from different interventions lost in different subjects); § data only collected in 15 subjects.

Frequency of oscillation of physiotherapy interventions

The frequency of oscillation of airflow of vibration, percussion, Flutter and Acapella are shown in Table 2.

Table 2 The frequency of oscillation of the physiotherapy interventions as determined by frequency spectral analysis.

Intervention	Frequency (Hz) Mean (SD)	Range (Hz)
vibration	8.4 (0.4)	7.3-10.0
percussion	7.3 (0.3)	6.5-8.0
Flutter	11.3 (1.5)*	7.5-13.7
Acapella	13.5 (1.7)*	10.0-18.3

Data are mean, SD of mean data of each subject and range. Legend: Significantly different from vibration * $p < 0.001$.

Forced expiratory manoeuvres

The mean PEF_R of cough and huff_{HIGH} were 2.9 times ($p < 0.001$) and 3.2 times ($p < 0.001$) greater than vibration respectively. The mean inspired volume of vibration, cough and huff_{HIGH} were similar (Table 3).

Table 3 Effects of forced expiratory manoeuvres compared to vibration on peak flow rates respiratory volumes and stimulation of coughs.

Intervention	PEFR (L/s)	PIFR (L/s)	PEFR/PIFR	V _I (L)	V _E (L)	coughs stimulated	n
vibration	1.58 (0.73)	1.06 (0.27)	1.51	1.78 (0.87)	2.44 (1.06)	0.7 (1.0)	17‡
cough	4.67 * (1.91)	1.68 (0.74)	3.07	1.78 (0.85)	2.02 (0.92)	1.9 (2.3)	17‡
huff _{HIGH}	5.04 * (2.30)	2.08 (1.42)	2.80	1.95 (0.82)	2.25 (0.91)	2.4 (3.7)	17‡

Data are means and SD of mean data of each subject. Legend: significantly different from vibration at * $p < 0.001$; PIFR = peak inspiratory flow rate, V_I = inspired volume, V_E = expired volume, n = number of subjects used in analysis, ‡ data lost due to technical difficulties (data from different interventions lost in different subjects).

There were no reported adverse effects during any of the interventions during this study. There was no significant difference in the perceived effort between the interventions and there was no significant effect of the order of the interventions on perceived effort.

DISCUSSION

This is the first study to report the physiological measures of expiratory flow rates, inspiratory volumes, oscillation frequencies and cough stimulation of vibration and to compare these to other physiotherapy interventions used for secretion clearance in patients with cystic fibrosis. Vibration had greater expiratory flow rates and a higher PEFR/PIFR ratio than the other physiotherapy interventions. However Flutter and Acapella had higher oscillation frequencies than vibration. These data allow inferences to be made about the possible effects of the physiotherapy interventions on secretion clearance based on theoretical rationale.

The PEFR of vibration was greater than all the other physiotherapy interventions and greater than PEFR of TLC_{relax} . This may be due to the fact that the compressive and oscillatory forces of vibration, applied to the chest wall, may increase intrapleural pressure and may therefore increase expiratory flow rates above those of elastic lung recoil as measured by TLC_{relax} . In addition, vibration did not have the added resistance at the mouth to impede expiratory flow as occurred during PEP, Acapella and Flutter. This shows that external forces applied to the chest wall during vibration increase peak expiratory flow rate and thus vibration may be a beneficial intervention for secretion clearance.

The absolute PEFR of vibration was greater than the other physiotherapy interventions.

Whether the absolute PEFR of vibration is adequate to clear secretions is not known, as the

critical absolute PEFR required for secretion clearance is dependant upon the volume of secretion and the viscoelasticity of the secretions [4, 19, 21, 22]. However, vibration had a PEFR/PIFR ratio greater than 1.1 which is the critical level required, for annular flow of secretions towards the oropharynx [4]. Therefore, if increasing expiratory flow contributes to the mechanisms for secretion clearance vibration is likely to be more effective for secretion clearance than the other physiotherapy interventions. The efficacy of vibration for secretion clearance needs to be investigated with radioaerosol clearance studies [7].

It could be argued that performing all the interventions at the same testing session may have resulted in an order effect in which the performance of an intervention may have altered the expiratory flow of subsequent interventions. To reduce this possible order effect, interventions were performed in a random order and bronchodilators were administered prior to testing to those subjects diagnosed with responsive airways. In addition, subjects did not report any increase in perceived breathing effort, suggesting that fatigue did not occur due to the multiple interventions.

The mean number of spontaneous coughs elicited during vibration was less than one and was not significantly different from the other physiotherapy interventions. These results suggest that stimulation of a cough is not a mechanism by which these physiotherapy interventions may aid secretion clearance.

The interventions of PEP and Acapella had slower PEFR's compared to vibration. The PEFR of PEP and Acapella was 3.6 and 2.7 times slower than vibration respectively. This was most likely due to the resistance provided by these two devices impeded expiratory flow rates. If the main physiological mechanisms by which secretion clearance is

enhanced are a high PEFr and an expiratory bias to airflow, then neither PEP nor Acapella would be the intervention of choice. However the results of this study cannot determine whether PEP or Acapella are ineffective in moving secretions to the central airway. The theoretical rationale of the positive expiratory pressure of these devices is to provide back-pressure within the airways during expiration. This increased pressure is proposed to stabilise collapsible airways, thus increasing expiratory flow in these airways [22] and to recruit the collateral ventilation [23] allowing gas behind the secretions, thus aiding the movement of these secretions towards the oropharynx. The results of this current study cannot discount the possibility of this mechanism occurring during PEP or Acapella.

Secretion clearance may be aided by the oscillation of airflow [1]. All the physiotherapy interventions, with the exception of PEP, had an oscillation frequency within the 3-17 Hz range shown to facilitate mucociliary clearance [1]. These oscillation frequencies may assist secretion clearance by altering rheology of the mucus [1] and increasing ciliary beat through stimulation of the ciliated epithelial cells [24, 25]. The oscillation frequency of Flutter (11.3 Hz) and Acapella (13.5 Hz) were close to the cited optimal frequency for secretion clearance of 13 Hz [1, 13]. The natural frequency of the ciliary beat is 11-15 Hz, and if airflow oscillates at a similar frequency, this resonance may increase the amplitude of the cephalad-ciliary beat which may increase mucus transport [1]. Flutter and Acapella may increase mucus transport by this resonance mechanism. There have been no known previous studies that have reported the oscillation frequencies of these interventions in patients with cystic fibrosis. The oscillation frequencies of all these physiotherapy interventions are within the range demonstrated to improve secretion clearance and therefore provide some evidence for their use during physiotherapy.

This is the only known study to report the expiratory flow rates of cough and huff in patients with cystic fibrosis. The PEFR's of cough (4.67 L/s) and huff_{HIGH} (5.04 L/s) in these subjects with cystic fibrosis are lower than the reported PEFR of these manoeuvres in normal subjects but similar to other subjects with airway obstruction [26, 27]. The forced expiratory manoeuvres of cough and huff_{HIGH} have been demonstrated to be effective for secretion clearance in patients with chronic obstructive pulmonary disease and bronchiectasis [8, 9, 10]. It is not surprising that cough and huff_{HIGH} had greater PEFR and higher PEFR/PIFR ratio than vibration. The results of this current study provide some evidence to support that cough and huff would be effective interventions to assist secretion clearance in patients with cystic fibrosis due to the high expiratory flow rates. However these forced expiratory manoeuvres do not have the benefit of the oscillation of airflow that may increase ciliary beat or alter the rheology of the secretions to aid secretion clearance. However, it would be interesting to measure the combined physiological effects of vibration applied during the forced expiratory manoeuvres.

The data from this study has provided some evidence to enable inferences to be made about the possible effects of vibration on secretion clearance based on theoretical rationale. Physiotherapists now have some evidence upon which they may base their decision making for treatment selection, as previous research noted that vibration is widely used in patients with excessive secretions [2]. Recommendations for clinical practice based on these inferences are that cough and huff_{HIGH} may be used to increase PEFR and optimise the PEFR/PIFR ratio, thus aiding secretion clearance by annular flow. If patients with cystic fibrosis are unable to cough or huff effectively, for example when fatigued, then vibration may be used as this resulted in the fastest PEFR and highest PEFR/PIFR ratio of the physiotherapy interventions. If the patients with cystic fibrosis do not have a carer to

apply vibration during airway clearance treatment then Flutter may be the intervention of choice as the PEFR/PIFR ratio was also >1.1 . The subjects in this study were encouraged to inspire as slowly as possible to optimise the chances of increasing the expiratory bias to airflow to increase the likelihood of clearance of secretions by annular flow. Theoretically, if secretions are to be cleared effectively, patients should be encouraged to inspire as slowly as possible.

Limitations

The principle aim of this study was to measure expiratory flow rates and oscillation frequencies of vibration and other physiotherapy interventions, not to measure the effects of these interventions on secretion clearance. Thus, we did not determine if the subjects had excessive secretions, nor did we measure secretion clearance. However, all subjects performed physiotherapy for airway clearance at least daily, implying that the subjects perceived they had excessive secretions. Secondly, the PEFR does not reflect flow in the peripheral airways, from where the physiotherapy interventions are proposed to clear secretions. A measure of interest would have been the maximal expiratory flow rates during the latter half of expiration as an indication of peripheral airflow. However, in order to compare this measure between interventions, the flow rates at similar lung volumes are required. The physiotherapy interventions were performed at different lung volumes reflecting current clinical practice thus precluding such a comparison.

CONCLUSIONS

This study provided physiological evidence to suggest that in subjects with cystic fibrosis with stable lung function, vibration may be effective in secretion clearance by assisting with the movement of secretions according to the theoretical rationale of increasing

absolute expiratory flow rates, PEFR/PIFR ratio > 1.1 and increased oscillation frequency. The effects of vibration on secretion clearance are still unknown. However, the data from this study has provided some evidence to enable inferences to be made about the possible effects of these physiotherapy interventions on secretion clearance based on theoretical rationale.

3749 words

Acknowledgements

We wish to acknowledge Ruth Dentice and Carmel Moriarty for assistance with subject recruitment and the financial support of the NHMRC and the Cumberland Foundation.

REFERENCES

1. King M, Phillips DM, Gross D, Vartian V, Chang HK, Zidulka A. Enhanced tracheal mucus clearance with high frequency chest wall compression. *Am Rev Resp Dis* 1983; 128: 511-515.
2. McCarren B, Alison JA, Lansbury G. The use of vibration in public hospitals in Australia. *Physiother Theory Pract* 2003; 19: 87-98.
3. Webber BA, Pryor JA, Bethune DD, Potter HM, McKenzie D. Physiotherapy techniques. In: Pryor JA, Webber BA, eds. *Physiotherapy for respiratory and cardiac problems*. 2nd ed. Edinburgh: Churchill Livingstone, 1998: 149.
4. Kim CS, Iglesias AJ, Sackner MA. Mucus clearance by two-phase gas-liquid flow mechanism: asymmetric periodic flow model. *J App Physiol* 1987; 62: 959-971.
5. Mellins RB Pulmonary physiotherapy in paediatric age group. *Am Rev Resp Dis* 1974; 110: 137-142.
6. Thomas J, Cook DJ, Brooks D. Chest physical therapy management of patents with cystic fibrosis. A meta-analysis. *Am J Resp Crit Care Med* 1995; 151: 846-850.
7. van der Schans C, Prasad A, Main E. Chest physiotherapy compared to no chest physiotherapy for cystic fibrosis. *The Cochrane Database of Systematic Reviews* 2003; 3: (Accessed, November 2, 2004, at <http://80-gateway1.ovid.com.opac.library.usyd.edu.au/ovidweb.cgi>).
8. Hasani A, Pavia D, Agnew JE, Clarke SW. Regional clearance during cough and forced expiration technique (FET): effects of flow and viscoelasticity. *Thorax* 1994; 49: 557-561.
9. Oldenburg Jr FA, Dolovich MB, Montgomery JM, Newhouse MT. Effects of postural drainage, exercise and cough on mucus clearance in chronic bronchitis. *Am Rev Resp Dis* 1979; 120: 739-745.
10. van der Schans CP, Piers DA, Beekhuis H, Koëter GH, van der Mark TW, Postma DS. Effect of forced expirations on mucus clearance in patients with chronic airflow limitation: effect of lung recoil pressure. *Thorax* 1990; 45: 623-627.
11. Mortensen J, Falk M, Groth S, Jensen C. The effects of postural drainage and positive expiratory pressure on tracheobronchial clearance in cystic fibrosis. *Chest* 1991; 100: 1350-1357.
12. Volsko TA, DiFiore JM, Chatburn RL. Performance comparison of two oscillating expiratory pressure devices: Acapella versus Flutter. *Resp Care* 2003; 48: 124-130.
13. Gross D, Zidulka A, O'Brien C, Wight D, Fraser R, Rosenthal L, King M. Peripheral mucociliary clearance with high-frequency chest wall compression. *J App Physiol* 1985; 58: 1157-1163.
14. Maxwell L, Crosbie J, Ellis ER. A new system for examining pattern of ventilation. *Aust J Physiother* 2001; 147: 271.

15. ATS/ERS Taskforce. Standardisation of spirometry *Eur Resp J* 2005;26: 319-338.
16. Wilson RC, Jones PW. Long term reproducibility of Borg scale estimates of breathlessness during exercise. *Clin Sc* 1991; 80: 309-312.
17. Winter DA, Patla AE. *Signal Processing and Linear Systems for the Movement Sciences*. Ontario: Waterloo Biomechanics. 1997; pp. 51-76.
18. Cystic Fibrosis Australia. *Cystic Fibrosis in Australia and New Zealand 2001. Annual report from the Australasia Cystic Fibrosis Data Registry*. pp 20-22, 43-44. 2003.
19. Kim CS, Greene MA, Sankaran S, Sackner MA. Mucus transport in the airways by two-phase gas-liquid flow mechanism: continuous flow model. *J App Physiol* 1986a; 60: 908-917.
20. Kim CS, Rodriguez CR, Eldridge MA, Sackner MA. Criteria for mucus transport in the airways by two-phase gas-liquid flow mechanism. *J App Physiol* 1986b; 60: 901-907.
21. Sackner MA, Kim CS. Phasic flow mechanisms of mucus clearance. *Eur J Resp Dis* 1987; 71 (Supple 153): 159-164.
22. Oberwaldner B, Evans JC, Zach MS. Forced expirations against a variable resistance: a new chest physiotherapy method in cystic fibrosis. *Ped Pulm* 1986; 2: 358-367.
23. Menkes HA, Traystman RJ. Collateral Ventilation. *Am Rev Resp Dis* 1977; 116: 287-309.
24. Sanderson MJ, Chow I, Dirkson ER. Intercellular communication between ciliated cells in culture. *Am J Physiol* 254 (Cell Physiol) 1988; 23: C63-C74.
25. Sanderson MJ, Charles AC, Dirksen ER. Mechanical stimulation and intercellular communication increases intracellular Ca²⁺ in epithelial cells. *Cell Regul* 1990; 1: 585-596.
26. Langlands J. The dynamics in cough in health and in chronic bronchitis. *Thorax* 1967; 22: 88-96.
27. Loudon RG, Shaw GB. Mechanics of cough in normal subjects and in patients with obstructive respiratory disease. *Am Rev Resp Dis* 1967; 96: 666-677.