

Procalcitonin for Reduced Antibiotic Exposure in Ventilator Associated Pneumonia - A Randomized Study

Daiana Stolz, M.D.,^{1, 2} Nicholas Smyrniotis, M.D.,³ Philippe Eggimann, M.D.,⁴ Hans Pargger, M.D.,⁵ Nehal Thakkar, M.D.,³ Martin Siegemund, M.D.,⁵ Stephan Marsch, M.D.,⁶ Janko Rakic, M.D.,¹ Beat Mueller, M.D.,⁷ and Michael Tamm, M.D.¹

From the Clinic of Pulmonary Medicine,¹ Division of Anesthesiology and Surgical Intensive Care Medicine,⁵ Division of Medical Intensive Care Medicine,⁶ Basel CH-4031, Switzerland; Department of Internal Medicine,⁷ Kantonsspital, CH-5001 Aarau, Switzerland; the Division of Pulmonary, Allergy and Critical Care Medicine,³ UMass Memorial Medical Center, 01655 - Worcester, MA, USA; the Department of Adult Critical Care Medicine,⁴ University Hospital, CH-1011 Lausanne, Switzerland; and the Harvard School of Public Health,² 02115, Boston, MA, USA

Corresponding author: Daiana Stolz, MD MPH, University Hospital, Petersgraben 4, CH-4031 Basel, Switzerland Phone: +41 (61) 265 5184, Fax: +41 (61) 265 4785, E-mail: dstolz@uhbs.ch

Running Head: Procalcitonin in VAP

Word count: 4748 words

Funding: Daiana Stolz was supported by grants from the Swiss National Foundation, the Liechtenstein Foundation and the “Freiwillige Akademische Gesellschaft Basel”. Nehal Thakkar was supported by the Will Rogers Foundation. Additional funding was granted by the Clinic of Pulmonary Medicine, University Hospital Basel, Switzerland. Funding obtained from Brahms (Hennigsdorf, Germany) was used for assay material and logistics. The sponsors of this investigator-initiated project had no involvement in design and conduct of the study, collection, management, analysis, and interpretation of the data, nor in the preparation, review, and approval of the manuscript or decision to submit the manuscript.

ABSTRACT

In patients with ventilator-associated pneumonia, guidelines recommend antibiotic therapy adjustment according to microbiology results after 72 hours. Circulating procalcitonin levels may provide evidence that facilitates reduction of antibiotic therapy.

In a multicenter, randomized, controlled trial, 101 patients with VAP were assigned to an antibiotic discontinuation strategy according to guidelines (control group) or to serum procalcitonin concentrations (procalcitonin group) with an antibiotic regimen selected by the treating physician. The primary endpoint was antibiotic-free days alive assessed 28 days after VAP onset and analyzed on an intent-to-treat basis. Procalcitonin determination significantly increased the number of antibiotic free-days alive 28 days after VAP onset (13 days [2-21] versus 9.5 days [1.5-17]). This translated into a reduction in the overall duration of antibiotic therapy of 27% in the procalcitonin group ($p=0.038$). After adjustment for age, microbiology, and center-effect, the rate of antibiotic discontinuation on day 28 remained higher in the procalcitonin group compared to patients treated according to guidelines (Hazard rate 1.6; 95 percent CI, 1.02-2.71). The number of mechanical ventilation-free days alive, ICU-free days alive, length of hospital stay and mortality rate on day 28 for the two groups were similar.

Serum procalcitonin reduces antibiotic therapy exposure in patients with ventilator associated pneumonia.

Key words: ventilator-associated pneumonia, antibiotic therapy, procalcitonin, biomarker

Word count: 200

INTRODUCTION

Ventilator associated pneumonia (VAP) is the leading cause of death from nosocomial infection in the United States.[1-3]. The mortality attributable to VAP has been reported to range between 24% and 50%.[4, 5] The estimated excess cost for an episode of VAP can be as high as \$40,000 per patient.[5, 6] Based on an estimate of 250,000 cases of VAP annually, the cost of treatment easily approaches \$10 billion in the United States alone.[7]

At the present time, there is no “gold standard” for the diagnosis of VAP and the optimal approach for diagnosing VAP remains to be defined.[8-10] Prompt initiation of broad-spectrum antibiotic therapy is a cornerstone of treatment of VAP because even relatively short delays in administering adequate antibiotic therapy are associated with an increased mortality rate.[11-13] Unfortunately, exposure to antibiotics, especially for ≥ 7 days, has been associated with subsequent emergence of infection with antibiotic-resistant bacteria and worse outcome.[14-18] Clinicians managing patients with suspected VAP should employ antimicrobial treatment strategies that minimize prolonged and potentially unnecessary antibiotic exposure to curtail resistance.[19] In this context, the concept of de-escalation therapy is emerging as an effective strategy to reduce the development of bacterial resistance in patients treated for VAP.[10, 20] Appropriately shortening the treatment duration is an important aspect of decreasing antibiotic-associated costs, minimizing selection pressures for resistant organism in the ICUs and improving outcomes.[8] The most recent treatment guidelines by the American Thoracic Society (ATS) and Infectious Disease Society of America (IDSA) have put forth a strategy to address early stoppage of antibiotics.[21] Accordingly, upon suspicion of VAP, lower respiratory tract sampling for cultures are obtained and empiric broad-spectrum antibiotics started. A systematic re-evaluation at 48 to 72 h, should allow to strongly considering stopping antibiotics if the patient is improving and cultures are negative. Should the culture results be positive and the patient has improved, then antibiotics being administered are tailored to the culture results, including potential for monotherapy.[10] Regrettably, this treatment algorithm often is not adhered to in daily clinical practice.[19] Some authors have suggested that inability

to obtain uncontaminated lower respiratory tract secretions and the pressures to treat an increasingly ill population are some of the factors that limit adherence to this approach.[22]

Procalcitonin, the precursor molecule of calcitonin, is up-regulated in severe bacterial infections and sepsis. [23, 24] Procalcitonin may be considered as valuable biomarkers for the diagnosis of bacterial infections, including VAP, where its levels are correlated with outcome. [25, 26] Its usefulness for guiding antibiotic therapy has been shown in patients admitted for community-acquired respiratory tract infections. [27-29] It has been recently suggested that serial measurements of procalcitonin allow to reduce the duration of antibiotic therapy without resulting in more adverse outcomes in critically ill patients with severe sepsis and septic shock.[30] Despite the growing body of literature in favor of this biomarker [27-31], it is unknown whether procalcitonin enables a refinement in the current ATS-IDSA antibiotic stoppage strategy in VAP. Therefore, we undertook this multicenter, randomized trial to determine if procalcitonin serum determination reduces antibiotic exposure in patients with clinically diagnosed VAP with a similar clinical and laboratory outcome.

METHODS (word count 1437)

Setting and Study Population

This is a multination, randomized, controlled open intervention trial in patients with VAP requiring treatment at an Intensive Care Unit (ICU) performed in seven ICUs (UMass Memorial Medical Center, Worcester, MA, USA; University Hospital Lausanne, Switzerland and University Hospital, Basel, Switzerland). The results are reported following the consolidated standards of reporting trials (CONSORT) statement. **(Figure 1)**[32]. We compared antibiotic therapy duration in patients treated according to current guidelines (control group) with patients in whom antibiotic treatment was guided by serum procalcitonin levels (procalcitonin group). The study was approved by the institutional review boards of all participating institutions and registered in the Current Controlled Trials Database as “ProVAP”-Study [ISRCTN61015974].[33] Written informed consent was obtained from all included patients or their legal representatives. All data were held and analyzed by the authors.

Intensive Care Unit patients intubated for mechanical ventilation for at least 48 hours were eligible for the study if they met all the following criteria: (1) older than 18 years; (2) clinically diagnosed VAP as defined by the American Thoracic Guidelines (new or persistent infiltrate on chest radiography associated with at least 2 of the following: purulent tracheal secretions, temperature $> 38^{\circ}\text{C}$, leukocyte count higher than $11000/\mu\text{L}$ or lower than $3000/\mu\text{L}$)[21]. Patients were excluded if they (1) were pregnant, (2) were enrolled in another trial; (3) had received immunosuppressants or on long-term corticosteroid therapy ($\geq 0.5 \text{ mg/kg}$ per day for > 1 month); (4) were severely immunosuppressed, including acquired immunodeficiency syndrome; (5) had a coexisting extrapulmonary infection diagnosed between day 1 and 3 requiring antibiotic therapy for greater than three days.

Randomization

Patients were randomly assigned to one of the two approaches after agreement of the attending physician. Randomisation was through arbitrary allocation to one of the two treatment assignments based on sealed, opaque envelopes. Block size was 20 envelopes. Treating physicians were not aware of

envelope contents before randomisation. In both groups, antibiotic therapy was started at inclusion (day 0), according to the decision of the attending physician, who was unaware of procalcitonin levels. Educational sessions, posters and instruction cards highlighting the current standard antibiotic de-escalation strategy according to the American Thoracic Society were provided to attending physicians and those responsible for patient's treatment decisions [21]. Thereby, particular emphasis was set on the recommendations regarding the stoppage of antibiotics according to microbiology results after 72 hours and duration of antibiotic therapy as suggested by the American Thoracic Society.[21]

After 72 hours (day 2), daily procalcitonin levels were notified to the attending physician, automatically and/or by personal communication, for patients randomized to the procalcitonin group. Thereafter, the physician in charge was advised to classify the patients into four groups, according to the probability of on going bacterial infection.[27] A procalcitonin level of $<0.25 \mu\text{g/L}$ suggested the absence of VAP and discontinuation of antibiotics was strongly encouraged. A procalcitonin level between $0.25 \mu\text{g/L}$ and $0.5 \mu\text{g/L}$ or a decrease by $\geq 80\%$ as compared to day 0 indicated that bacterial infection was unlikely and reduction or discontinuation of antibiotics was encouraged. A procalcitonin level $\geq 0.5 \mu\text{g/L}$ or decrease by $< 80\%$ as compared to day 0 was considered to indicate unresolved bacterial infection and reduction or discontinuation of antibiotic was discouraged. A procalcitonin level of $>1 \mu\text{g/L}$ strongly suggested unresolved bacterial infection and antibiotic discontinuation was strongly discouraged. After day 2, evaluation of procalcitonin levels was performed by comparing daily procalcitonin levels with the immediately previous value. Reevaluation of the clinical status and measurement of serum procalcitonin levels were performed daily for both groups up to day 10 after inclusion. Attending physicians responsible for patients in the control group remained unaware of procalcitonin throughout the study.

Measurement of Serum Procalcitonin

Measurements were done using a time-resolved amplified cryptate emission (TRACE) technology assay (Kryptor[®] PCT, Brahms AG, Hennigsdorf, Germany) [27] with a functional assay sensitivity of

0.06 µg/L, about four-fold above mean normal levels [34]. Assay time is less than 20 minutes and results were routinely available within one hour.

Antibiotic Treatment

Drug selection was left to the discretion of the treating physicians, including any adaptation considered necessary as a function of clinical, laboratorial, radiological or microbiological results, including bacteriology susceptibility patterns. Upon inclusion in the study and before initiation of antibiotic therapy, sampling of respiratory secretions (endotracheal aspirates, bronchoalveolar lavage, or protected brush specimens) was strongly recommended. The attending and resident physicians in participating ICUs were periodically re-educated on standard antibiotic reduction strategy recommendations. [21] In all situations, treating physicians retained control of antibiotic treatment and were empowered to refrain from antibiotic discontinuation if the duration of antibiotic treatment was considered inadequate or a patient's condition deteriorated.

Baseline Assessment and Follow-up

At the time of enrollment the following information was recorded from each subject: age, sex, preexisting comorbidities, severity of the underlying medical condition(s), primary reason for initiating mechanical ventilation, duration of prior mechanical ventilation, use of any antibiotics within 14 days of VAP onset, presence of infiltrates on chest radiography, fever $\geq 38^{\circ}\text{C}$, leukocytosis or leukopenia, purulent tracheal secretions, body temperature, heart rate, mean arterial pressure, oxygen saturation, ratio of partial pressure of arterial oxygen to the fraction of inspired oxygen ($\text{PaO}_2/\text{FIO}_2$), leukocyte counts and serum procalcitonin values.. **(Table 1).**

The following indices were calculated: CPIS score, simplified acute physiologic score II (SAPS II); organ dysfunction and/or infection score (ODIN); sepsis-related organ failure assessment score (SOFA).

During the 28 day follow-up period the following information was recorded: body temperature, heart rate, mean arterial pressure, oxygen saturation, ratio of partial pressure of arterial oxygen to the fraction of inspired oxygen, CPIS, SOFA, leukocyte counts and serum procalcitonin values for 10 consecutive days; mechanical ventilation status and antibiotic use daily throughout the 28-day study period; vital signs, leukocyte counts and SOFA scores at days 14, 21 and 28; and the patient's status at discharge from the hospital. Any antibiotic use was recorded daily up to day 28. The number of antibiotic-agent days, i.e. number of antimicrobial agents per day x number of days) was also documented. We calculated the number of antibiotic free days alive as the number of days during the 28 days after living patients had been randomized and had not received any antibiotic, as previously described.[16] VAP-related clinical and/or radiological deterioration was defined as an increase in clinical pulmonary infection score (CPIS) of more than two points.

Outcome Measures

The primary endpoint was the number of antibiotic free days alive assessed 28 days after enrollment in the study. Any antibiotic exposure after inclusion, i.e. total antibiotic exposure days and total antibiotic-agent days, regardless of indication, was taken into account for antibiotic exposure analyses. Antibiotic therapy discontinuation related to death was considered a censored event. Secondary endpoints were the number of mechanical ventilation-free days, the number of ICU-free days alive, the evolution of the signs and symptoms potentially linked to pulmonary infection, oxygen saturation, ratio of partial pressure of arterial oxygen to the fraction of inspired oxygen, the evolution of the SOFA, ODIN and CPIS scores, length of hospital stay, the VAP-related clinical deterioration rate, and overall mortality at 28 days. Patients discharged prior to the end of the study (28 days) were followed up at home or in the post-acute institution for the outcomes of interest.

Statistical Analysis

The trial was designed to demonstrate the superiority of the algorithm incorporating procalcitonin in terms of increasing the number of antibiotic-free days in patients alive within 28 days of inclusion in the study. The sample size was calculated based on the antibiotic-use days in a prior study [16]. Considering 13 antibiotic-free days in the control group and 18 antibiotic-free days in the procalcitonin group, a sample size of 84 patients (42 per group) was necessary to detect a significant difference in antibiotic-free days alive between both groups with a power of 90% and an alpha error of 0.05 using a two-tailed test. Assuming 8% lost to follow up, we planned the inclusion of 100 subjects.

Discrete variables are expressed as counts (percentage) and continuous variables as means (standard deviation) or medians (interquartile ranges). Endpoints were predefined and analyzed on the basis of intention-to-treat. Comparability of the control group and the procalcitonin group was analyzed by χ^2 test or Fisher's exact test for categorical variables and non-parametric Mann-Whitney U test or unpaired t test for continuous variables, as appropriate. Cumulative-events curves were estimated with the Kaplan-Meier method. Time to discontinuation of antibiotic treatment was compared between the two study groups by use of the log-rank test. Cox proportional hazards regression analysis was used to evaluate the occurrence of events on day 28, adjusting by age, microbiology of respiratory samples, and center-effect ("a priori decision").

RESULTS

Baseline characteristics of the patients

During the study period, of the 164 patients with VAP screened for eligibility, 101 were eligible and randomized into the procalcitonin group (n=51) or into the control group (n=50) (**Figure 1**). The clinical characteristics of these 101 patients at baseline were similar, except that the ODIN score was slightly higher ($p=0.042$) for the control group (**Table 1**). Three-fourths of the patients had received antibiotics within 14 days prior to inclusion in the study, without a significant difference between the groups ($p=0.119$). At inclusion, serum procalcitonin levels were <0.25 $\mu\text{g/L}$ in 29 patients (15 in the control group and 14 in the procalcitonin group; $p = 0.82$), 0.25 to 0.499 $\mu\text{g/L}$ in 13 patients (4 in the control group and 9 in the procalcitonin group; $p = 0.23$), and ≥ 0.5 $\mu\text{g/L}$ in 56 patients (30 in the control group and 26 in the procalcitonin group; $p = 0.42$). Median [interquartile range] procalcitonin levels in patients pre-treated with antibiotics was 0.73 $\mu\text{g/L}$ [0.22-2.36], which was not significantly different from patients without antibiotic exposure 0.72 $\mu\text{g/L}$ [0.15-2.36].

Microbiology

Microbiological investigation of respiratory tract secretion samples was performed in 97 patients (96 %). In 74 patients (73%) a causative microorganism was identified. The rate of positive bacterial cultures was similar in the control group and the procalcitonin group (76 % vs. 71 %). Microorganisms considered responsible for VAP are listed in **Table 2**. In both groups the most frequently isolated gram negative microorganism was *Pseudomonas aeruginosa* followed by *Klebsiella* species. Most commonly isolated gram positive bacteria were methicillin-sensitive *Staphylococcus aureus* and methicillin-resistant *Staphylococcus aureus*. Microbiologic cultures from others sites were positive in 20 patients (10 control group versus 9 procalcitonin group, $p=0.625$). In the control group, infection sites were catheter tips (2 *staphylococcus non-aureus*, 1 coagulase negative *staphylococcus*), spinal fluid (1 *streptococcus viridans*, 1 *staphylococcus aureus*), urin (1 *enterococcus*), wound swabs (1 coagulase negative *staphylococcus*, 1 methicillin-resistant *staphylococcus aureus*, 1 *enterococcus*), and blood cultures (1 *staphylococcus*

aureus). Correspondingly, in the procalcitonin group, infections sites were catheter tips (1 *candida albicans*), spinal fluid (2 *streptococcus pneumonia*), urine (1 *enterococcus*), wound swabs (1 *enterococcus*, vancomycin-resistant *enterococcus*, 1 *enterobacter cloacae* and *proteus mirabilis*), feces (1 *clostridium difficile*), and blood cultures (1 *candida albicans*).

Primary Endpoint – Reduction of Antibiotic Exposure

The number of antibiotic-free days alive within 28 days of diagnosis of VAP was significantly higher in the procalcitonin group than in the control group (median [IQR], 13 days [2-21] versus 9.5 days [1.5-17], **Figure 2A**). This translated into a reduction in the overall duration of antibiotic therapy of 27% in the procalcitonin group (median [IQR] 15 [10-23] versus 10 [6-16] days, $p=0.038$). The total number of antibiotic-agent days was higher in the control group than in the procalcitonin group (1341 days versus 1077 days). At the end of serial procalcitonin measurements (day 10), the rate of antibiotic discontinuation was significantly higher in the procalcitonin group compared to patients treated according to guidelines (Hazard rate 2.235; 95 percent CI 1.077-4.64, $p=0.031$). Cox regression-based adjustment of the baseline variables did not substantially modify these findings. The adjusted rate for antibiotic discontinuation of patients in the procalcitonin group versus those in the control group on day 28 was 1.66 (95 percent CI, 1.02-2.71, **Figure 2B**) after adjustment for age, respiratory tract culture results, and center-effect.

Overall, the median [IQR] duration of antibiotic therapy for VAP was 11 days [6-17]. Antibiotic therapy was continued beyond 7 days in 82 percent of subjects in the control group and in 65 percent in the procalcitonin group, ($p=0.044$, **Figure 3A**). In the 28 days following the diagnosis of VAP, antimicrobial monotherapy was administered in 22 percent of cases (8 control group, 14 procalcitonin group); combination therapy with two antimicrobials in 24 percent (13 control group, 11 procalcitonin group); three antimicrobials in 21 percent (12 control group, 9 procalcitonin group); four antimicrobials in 17 percent (8 control group, 9 procalcitonin group); and five or more antimicrobials in 16 percent (9 control group, 7 procalcitonin group). The most commonly administered antibiotics in both groups

included piperacillin/tazobactam in 78 patients (77%), vancomycin in 44 (44%), meropenem/imipenem in 31 (31%), levofloxacin in 29 (29%), and amoxicillin/clavulanic acid in 25 (25%). Appropriate initial empiric antibiotic therapy, defined as a regimen combining an aminoglycoside or a fluoroquinolone plus a betalactam or an antipseudomonal carbapenem, was given in 86 percent of the cases without difference between the groups ($p=0.345$). After 72 hours, combination therapy with two or more drugs could be reduced to monotherapy in 54 percent of patients in the procalcitonin group as compared to 28.6 percent of patients in the control group ($p=0.008$, **Figure 3B**).

A microbiologically confirmed VAP influenced maintenance of antibiotic therapy in the control group (Hazard rate 2.30; 95 percent CI, 1.11-4.77) but not in the procalcitonin group (Hazard rate 1.17; 95 percent CI 0.59-2.32). Correspondingly, patients with microbiologically diagnosed VAP ($n = 73$) had significantly more median [IQR] antibiotic-free days alive in the procalcitonin group compared to the patients treated in the control group (median [IQR] 14 [1-21]) versus 7 [0-13.5], $p=0.017$). Conversely, the number of antibiotic-free days alive were similar in those with negative microbiology ($p=0.563$). Neither in the procalcitonin group nor in the control group did CPIS scores influence discontinuation of antibiotic therapy ($p=0.845$ and $p=0.175$).

Reasons for antibiotic discontinuation in the control group were clinical and laboratorial improvement (28%, $n= 14$), extubation (16%, $n = 8$), negative microbiology results (16%, $n=8$) and improvement in CPIS score (14%, $n=7$). In 10% of the cases ($n=5$) antibiotic therapy was maintain beyond 28 days. A total of 8 patients deceased while on antibiotic therapy for VAP. Strategies for discontinuation of antibiotic therapy in VAP included cessation of monotherapy (20%, $n=10$), dual therapy (14%, $n=7$), or triple therapy (2%, $n=1$), de-escalation from triple to dual therapy (16%, $n=8$), de-escalation from triple or dual to monotherapy (22%, $n=11$) or adjustment of antibiotics without reduction of the number of antibiotic agents (26%, $n=13$). Median time to cessation of monotherapy was 9 days [5-12] and to de-escalation from dual to monotherapy was 2 days [1-5]. In the procalcitonin group, a total of 8 patients with improved CPIS scores have received prolonged antibiotic treatment despite low procalcitonin values. In 4 cases (8%), positive blood cultures with gram negative bacilli were observed

and in another 4 cases (8%), treating physicians decided to prolong antibiotic treatment due to documented pulmonary infection with gram negative bacilli (*Acinetobacter baumannii*, *Pseudomonas aeruginosa*, *Stenotrophomonas maltophilia*, and *klebsiella*). Duration of antibiotic therapy for VAP was significantly longer in these patients (15 days [14-19] versus 7 days [4-16], $p=0.023$). The incidence of acquired or persistent microorganisms up to 28 days after VAP onset did not differ between randomized groups (16 versus 13, $p=0.5149$, **Table 3**).

Secondary Endpoints - Clinical and Laboratory Outcome

None of the secondary outcome events (**Table 4**) or physiologic changes seen from day 1 through day 28 differed significantly between patients in the control and procalcitonin groups (**Supplementary Data**). The two groups had a similar median (IQR) number of mechanical ventilation and ICU free days alive. In both groups, median mechanical-ventilation-free days alive and ICU-free days alive were lower for patients with infection due to GNB and MRSA. VAP-related clinical and radiological deterioration rate did not differ in both groups. Hospital discharge rate and post-discharge institution placement pattern was also comparable in both groups.

Deaths in the two groups were due to traumatic brain injury/subarachnoid hemorrhage (4 in the control group / 4 in the procalcitonin group), septic shock (2 / 1), respiratory failure (3 / 1), cardiogenic shock (1 / 1), ARDS (1 / 0), multiorgan failure (0/1), and acute liver failure (1 / 0). Patients who died during their hospitalization had significantly higher levels of procalcitonin on admission than those who did not die (median [interquartile range] 1.29 [0.21-2.00] vs. 0.58 [0.45-5.43], $p=0.02$), higher SAPS II scores (47.5 [43.4-55] vs. 38 [31-47], $p<0.001$), and higher SOFA scores (9 [7-13.8] vs. 6 [6-9], $p=0.004$). CPIS scores ($p=0.935$) and leukocyte counts ($p=0.309$) were similar in subjects who died and those who did not die.

Procalcitonin levels at VAP diagnosis correlated significantly with the increasing severity of the disease, as defined by the SAPS II score ($r^2 = 0.358$, $p<0.001$), the number and the severity of acute organ failures related to ICU mortality ($r^2 = 0.474$, $p<0.001$), and the presence or absence of organ dysfunctions

and/or infection ($r^2 = 0.254$, $p=0.015$). Circulating procalcitonin levels decreased similarly within 28 day of VAP onset in both randomized groups (**Figure 4**).

DISCUSSION

In this multicenter, randomized clinical trial we observed a benefit of incorporating procalcitonin into the antibiotic reduction strategy suggested by current ATS-IDSA guidelines. The average number of antibiotic free-days alive from day 1 to day 28 was 27% higher for patients who had been randomized to the procalcitonin group than for patients assigned to the control group. The procalcitonin-guided approach allowed to reduce the overall number of antibiotic-free days alive at 28 days by the same amount as reported by Chastre et al. in his landmark trial, which randomized patients to receive 8 days or 15 days of antibiotic therapy for VAP.[16] Procalcitonin-guided strategy was able to significantly reduce the median duration of overall antibiotic treatment by one third, i.e. from 15 to 10 days. Despite precise guidelines based on strong evidence for reducing the duration of antibiotic treatment for VAP to 8 days [16], such practice has not been reported to be successfully translated in current clinical practice.[21] Accordingly, the long duration of antibiotic treatment in the control group in our study may be close to the current clinical practice in many centers. This seems to be true particularly for US critical care centers, in which we observed a significantly longer median [IQR] overall duration of antibiotic therapy (15 days [9-23] versus 11 days [6-17], $p=0.02$) and antibiotic therapy for VAP (14.5 days [7-20] versus 10 [5-15], $p=0.027$) as well as higher total number of antibiotic-agent days (26.6 days [18.5-53.5] versus 12 days [7-24], $p<0.001$, respectively) as compared to European centers. In contrast, the median overall antibiotic duration of 10 days in the procalcitonin-guided group, very close to the 8 days targeted by the recommendations for VAP, suggest that the use of this biomarker in addition to the other usual clinical, laboratory and radiological information may lead to effective bedside antibiotic discontinuation. The benefit associated with procalcitonin-guidance was achieved by an increase in the number of antibiotic free days alive after VAP onset. One shall notice that the primary end-point of this study accomplished to evaluate whether a procalcitonin-guided strategy provides a *real* benefit in decreased *overall* antibiotic exposure – by assessing the number of days without *any* antibiotic exposure. In this context, it represents a rather robust, i.e. conservative, outcome.

The absence of differences in outcome parameters including the evolution of clinical and radiological signs and symptoms of pulmonary infection, the number of mechanical-ventilation-free days alive, the number of ICU-free days alive, the length of hospital stay and overall mortality suggest that procalcitonin-guided antibiotic reduction is not associated with a worse outcome in VAP. This also supports the concept that complications associated with long-term broad-spectrum antibiotic use may also have been avoided.[7]

Currently, duration of antibiotic therapy in VAP is based on empirical rules and physician's preference.[10, 16] De-escalation of therapy is advocated for as an appropriate strategy for managing antibiotics for VAP. The de-escalation approach provides clinical balance between one extreme of using broad-spectrum, empiric antimicrobial agents as the sole treatment strategy and the other extreme of delaying the initiation of targeted therapy pending bacteriologic results.[35] The most recent ATS-IDSA guidelines emphasize the need for early and appropriate antibiotic therapy followed by de-escalation whenever possible, based on culture and patient response.[21] Clinical strategies, based on evidence of clinical improvement, as defined by reduction in serial CPIS or improvement of the $\text{PaO}_2/\text{FiO}_2$ ratio at days 3 to 5, have also succeeded in limiting antibiotic use in patients with mild VAP in the context of clinical trials.[15, 17, 18, 36] Furthermore, cultures obtained from respiratory tract sampling consistently result in a change of antibiotic regimen allowing narrowing of antibiotics or monotherapy once microorganisms and their susceptibilities have been identified.[37, 38] Such practices were included in our strategies to reduce antibiotic exposure in both groups and may have contributed to our results.

It has been proposed that biomarkers might contribute to customize the duration of antibiotic treatment at a patient level, resulting in a greater tendency to limit or discontinue antibiotics. [19, 30] Procalcitonin concentration closely parallels the severity and evolution of infection, including VAP, for which it has also a prognostic value.[39-41] This was also observed in our patients. We believe it is fair to assume that physicians taking care of patients with known (low) procalcitonin levels (procalcitonin group) tended to feel more confident in discontinuing antibiotic therapy than physicians taking care of patients, in whom procalcitonin levels were unknown (control group). Procalcitonin levels might have

facilitated the discontinuation of one or all antimicrobial agents at the same time. This might explain why not only the number of antibiotic free days alive was higher in patients randomized to the procalcitonin group, but also why patients randomized to the procalcitonin group received a smaller amount of antibiotic agent-days. In agreement with others, we observed a wide variation in serum procalcitonin concentrations at onset of VAP. [41, 42] This suggests that individual VAP patients may require different lengths of antibiotic therapy according to different virulence of the causative microorganism and to individual host response to infection, factors potentially reflected by the evolution of serum procalcitonin levels.[30]

The utility provided by the procalcitonin-guided strategy was more impressive in those patients with microbiologically diagnosed VAP. In our study, despite a high proportion of previous antibiotic exposure, VAP was microbiologically documented in 74% of the patients. This may again support that a procalcitonin-guided antibiotic discontinuation might provide an additional benefit to the currently applied microbiologically-guided de-escalation strategy.

In contrast to the impressive reduction in antibiotic exposure reported in patients admitted with community acquired respiratory tract infections [27-29], we achieved only modest gains in antibiotic use for VAP. This may be due to additional difficulties in the interpretation of procalcitonin results in such ill patients, potentially related to higher proportion of previous systemic inflammatory response syndromes, multiorgan failure, previous infection and/or surgery, all known to raise the procalcitonin levels. [25, 43] Importantly, in the present study, procalcitonin kinetics assessed up to day 10 was used to support antibiotic discontinuation following at least 72 hours empiric antimicrobial therapy for VAP. Hence, the results of this study do not allow primarily withholding of antibiotics based on procalcitonin levels at the time of VAP onset.

Our study has several limitations. First, we included a relatively small number of patients and our results should not be generalized to other settings such as hospital acquired pneumonia and non-critically ill patients. It may not apply in the context of de-escalation strategies other than those suggested by ATS-IDSA guidelines. Second, these findings have to be validated in a larger cohort of patients, allowing sub-

group analyses for selected populations, in whom infections may be more difficult to treat, e.g. severe COPD, immunocompromised patients. Third, our intervention was performed in a clinical setting, in which the final decision to de-escalate antibiotic treatment was left to the discretion of the attending physician. Thereby and in contrast to the study by Chastre et al., physicians were not obliged to always conform to the study protocol. However, protocol overruling would result in a “conservative bias”, potentially underestimating the benefit of a procalcitonin-guided approach. Given the complexity of monitoring adherence to guidelines, we believe that it is fair to expect results of a study mimicking daily bedside practice to be more likely reproduced in a non-investigation setting. Fourth, we should emphasize that our trial was not designed to test the potential efficacy of a shorter duration of antibiotic treatment. In contrast, we demonstrated that a shorter duration of antibiotic therapy was not associated with a worse outcome. Finally, we did not conduct a formal cost-benefit study. Larger clinical trials are needed to explore the overall clinical and economic impact of the reduction of exposure to antibiotics in VAP patients. Strengths from our study are the multicentric design, the primary end-point evaluating the total number of antibiotic-free days alive up to 28 days irrespective of the indication for antibiotic therapy and the potential reproducibility of the results in daily clinical practice.

In conclusion, serum procalcitonin reduces antibiotic exposure in critically ill patients treated for ventilator-associated pneumonia.

Acknowledgments

We thank Anja Meyer, Stephen Heard, Jaclyn Longtine, and Karin Longtine for continuous support throughout the study; Andrea Azzola for critical discussions on the initial study protocol; and Andy Schötzau for statistical advice.

LEGENDS

Figure 1 Trial Profile

CAP denotes community-acquired pneumonia, ITT denotes intention to treat

Figure 1




Figure 2

A Median Number of Antibiotic-Free Days Alive in Patients in the Control and the Procalcitonin Group at 28 Days after Onset of Ventilator Associated Pneumonia

Results of the procalcitonin group are shown in black, results of the control group in gray, $p = 0.049$

B Cumulative Frequency Distribution Curve for the Time to Discontinuation of Antibiotics in Patients in the Control and the Procalcitonin Group on Day 28

Results of the procalcitonin group are shown in black, results of the control group in gray

Adjusted by age, respiratory tract culture results, and center-effect, $p=0.043$

Figure 2A




Figure 2B




Figure 3

A Duration of Antibiotic Therapy

Percent of patients in both randomized groups on antibiotics beyond 2 days, 7 days, 14 days and 21 days.

Results of the procalcitonin group are shown in black, results of the control group in gray.

B Antibiotic Reduction Profile

Percent of patients in control group and procalcitonin group on monotherapy and combination therapy with two or three antibiotics after 72 hours, $p=0.008$. Results of the procalcitonin group are shown in black, results of the control group in gray.

Figure 3A




Figure 3B







Figure 4


A-F Physiological and Functional Score Changes from Day 0 to Day 28


Results of the procalcitonin group are shown in black, results of the control group in gray.














Figure 5 Serum Procalcitonin Changes from Day 0 to Day 28




Table 1 Baseline Characteristics of the 101 Patients Randomized to Control Group or Procalcitonin group*

Characteristic	Control group (n = 50)	Procalcitonin group (n = 51)
Age - years (min-max)	59 (18-83)	53 (21-88)
Male sex - no. (%)	37 (74)	38 (75)
Admission category - no. (%)		
- Medical	26 (52)	27 (53)
- Emergency surgery	20 (40)	23 (45)
- Elective surgery	3 (6)	1 (2)
Origin - no. (%)		
- Home	26 (52)	35 (69)
- Other institution	24 (48)	16 (31)
Coexisting illnesses – no. (%)		
- Coronary artery disease	4 (8)	9 (18)
- Hypertensive heart disease	8 (16)	8 (16)
- Congestive heart failure	27 (54)	21 (41)
- Renal dysfunction	7 (14)	9 (18)
- Liver disease	3 (6)	4 (8)
- Diabetes mellitus	13 (26)	10 (20)
- Chronic obstructive pulmonary disease	11 (22)	8 (16)
- Neoplastic disease	5 (10)	3 (6)
- Substance abuse	8 (16)	5 (10)

Reason for mechanical ventilation - no. (%)		
- Cardiovascular failure	16 (32)	16 (31)
- Acute respiratory failure	29 (58)	24 (47)
- Trauma	13 (26)	20 (39)
- Neurologic failure	12 (24)	18 (35)
- Sepsis	15 (30)	10 (20)
- Miscellaneous	4 (8)	6 (12)
Duration of MV before VAP onset - median (IQR), d	6 (4-10)	6 (3-7)
Antimicrobial therapy 14 days before VAP – no. (%)	41 (82)	35 (69)
Duration of antimicrobial therapy 14 days before VAP - median (IQR), d	5 (1-9.5)	3 (0-8)
Criteria for VAP diagnosis - no. (%)		
- Infiltrate on chest-x-ray	50 (100)	51 (100)
- Fever $\geq 38^{\circ}\text{C}$	30 (60)	26 (51)
- Leukocytosis or leucopenia	33 (66)	32 (63)
- Purulent tracheal secretions	28 (56)	31 (61)
- CPIS ≥ 6	47 (94)	43 (84)
- $\text{FiO}_2/\text{PO}_2 < 200$	24 (48)	25 (50)
- Shock	12 (24)	11 (22)
- Positive blood cultures within 48 hours	18 (36)	14 (28)
Diagnostic score – mean (SD)‡		
- SAPS II	45 (14)	42 (13)
- ODIN	2.3 (1.0)	1.9 (0.9)
- SOFA	8.2 (3.4)	7.3 (3.4)
Clinical and laboratorial findings – mean (SD)		
- Body temperature - $^{\circ}\text{C}$	38.2 (1.2)	37.9 (0.9)
- Heart rate	91 (23)	88 (18)
- Mean arterial pressure - mmHg	79 (16)	81 (16)
- SaO_2 - %	97 (3)	96 (4)
- $\text{PaO}_2/\text{FiO}_2$ - mmHg	203 (97)	218 (114)
- Leukocyte count - μL	13.3 (5.9)	12.0 (6.6)
- Procalcitonin – $\mu\text{g/L}$ median [IQ range]	0.73 (0.21-2.36)	0.66 (0.22-2.69)

no. = number, min = minute, mm Hg = millimeter Hg, SD = standard deviation, IQ = interquartile, SAPS II = simplified acute physiologic score II, ODIN = organ dysfunction and/or infection, SOFA = sepsis-related organ failure assessment. ‡ODIN $p=0.042$. There were no other significant differences between the groups with respect to any other characteristic. Because of rounding, percentages may not sum to 100. The conversion factor for procalcitonin is as follows: $\mu\text{g/L} \times 0.161 = \text{nmol/L}$.

Table 2. Microbiological Cultures Results From Bronchoscopic and/or Endotracheal Aspirates Specimens in Patients with ventilator-associated pneumonia according to the Treatment Algorithm*

Organism	No. (%) of Pathogens	
	Control group (n = 50)	Procalcitonin group (n = 51)
Bacilli – n (%)		
<i>Pseudomonas species</i>	15 (30)	10 (20)
<i>Acinetobacter species</i>	0 (0)	4 (8)
<i>Stenotrophomonas maltophilia</i>	2 (4)	5 (10)
<i>Escherichia coli</i>	8 (16)	3 (6)
<i>Enterobacter</i>	6 (12)	2 (4)
<i>Proteus</i>	0 (0)	1 (2)
<i>Serratia</i>	4 (8)	2 (4)
<i>Klebsiella</i>	5 (10)	8 (16)
<i>Citrobacter</i>	1 (2)	2 (4)
<i>Morganella morganii</i>	0 (0)	1 (2)
<i>Hemophilus</i>	5 (10)	4 (8)
Others	2 (4)	5 (10)
Cocci – n (%)		
MSSA	10 (20)	10 (20)
MRSA	5 (10)	5 (10)
Coagulase-negative staphylococci	1 (2)	3 (6)
<i>Streptococcus</i>	1 (2)	6 (12)
<i>Neisseria</i>	4 (8)	1 (2)
<i>Enterococcus</i>	2 (4)	0 (0)
Others	0 (0)	2 (4)

MSSA = methicillin-susceptible *Staphylococcus aureus*, MRSA = methicillin-resistant *S. aureus*.

Organisms shown are those isolated at significant concentrations from semi-quantitative endotracheal aspirates or quantitative cultures of protected specimen brush ($\geq 10^3$ colony-forming units/mL) and/or bronchoalveolar lavage fluid ($\geq 10^4$ colony-forming units/mL). Percentages do not sum to 100 because of concomitant isolated microorganisms.

Table 3. Data on Acquired or Persistent Microorganisms in Patients with ventilator-associated pneumonia according to the Treatment Algorithm

Control group (n = 50)			Procalcitonin group (n = 51)		
Day after VAP onset	Microorganism		Day after VAP onset	Microorganism	
5	<i>Pseudomonas</i>	persistent	4	<i>Stenotrophomonas</i>	new
5	<i>Pseudomonas</i>	persistent	4	MRSA	new
5	MRSA	persistent	5	<i>Pseudomonas</i>	persistent
8	<i>Pseudomonas</i>	new	9	MRSA +	new
10	<i>Pseudomonas</i>	persistent		<i>Klebsiella</i>	
12	MRSA + <i>Enterobacter</i>	new	9	<i>Alcaligenes</i> +	new
15	<i>Serratia</i>	persistent		<i>Acinetobacter</i>	
20	<i>Stenotrophomonas</i>	persistent	10	MRSA	new
21	<i>E. coli</i>	new	10	<i>Acinetobacter</i>	new
21	<i>Pseudomonas</i>	persistent	15	<i>Klebsiella</i>	persistent
24	<i>Enterococcus</i>	new	15	MRSA	persistent
28	<i>Klebsiella</i>	persistent	21	<i>Clostridium difficile</i>	new
28	<i>Pseudomonas</i>	new	28	<i>Enterobacter</i>	persistent
28	MRSA	persistent	28	<i>Pseudomonas</i>	persistent
28	<i>Pseudomonas</i>	persistent	30	<i>Pseudomonas</i>	persistent
28	MRSA +	new			
	<i>Pseudomonas</i>	persistent			

MRSA = methicillin-resistant *S. aureus*. Organisms shown are those isolated at significant concentrations from semi-quantitative endotracheal aspirates or quantitative cultures of protected specimen brush ($\geq 10^3$ colony-forming units/mL) and/or bronchoalveolar lavage fluid ($\geq 10^4$ colony-forming units/mL).

Table 4. Secondary Study Outcomes in Patients with Ventilator-Associated Pneumonia according to the Treatment Algorithm*

	Control group (n = 50)	Procalcitonin group (n = 51)	P Value
MV-free days alive, days 1 to 28 – median (IQR)			
All patients	19 (8.5-22.5)	21 (2-24)	0.455
Non-fermenting GNB	15 (7-20)	12 (0.3-23)	0.867
MRSA	15 (0-22)	14 (6-21)	1
Other bacteria	19 (1.8-24.8)	22 (15.5-24.5)	0.284
No bacteria	21.5 (18.8-23.3)	23 (3.5-27)	0.563
ICU-free days alive, days 1 to 28* – median (IQR)			
All patients	8.5 (0-18)	10 (0-18)	0.526
Nonfermenting GNB	0 (0-11)	4 (0-13.5)	0.683
MRSA	15 (2-17)	9 (1.5-15)	0.548
Other bacteria	7.5 (1-17.8)	14 (7.5-20)	0.139
No bacteria	18 (1.5-20)	10 (1.-19.5)	0.554
Length of hospital stay, d – median (IQR)			
All patients	26 (16.8-22.3)	26 (7-21)	0.153
Nonfermenting GNB	34 (26-46)	31 (13-35.5)	0.130
MRSA	28 (17-41)	26 (23.5-37.5)	1.0
Other bacteria	24 (16.5-32.5)	21.5 (14-28)	0.442
No bacteria	28.5 (16-38)	29 (9.5-33)	0.343
VAP-related clinical deterioration, days 1 to 28 – n (%)‡	7 (14)	5 (10)	0.759
Discharge home, days 1 to 28– n (%)	3 (6)	5 (10)	0.479
Discharge to another institution, days 1 to 28 – n (%)	32 (64)	35 (69)	0.509
Death from all causes, days 1 to 28 – n (%)	12 (24)	8 (16)	0.327
In hospital mortality – n (%)	14 (28)	10 (20)	0.322

* GNB = gram-negative bacilli; ICU = intensive care unit; VAP = ventilator-associated pneumonia; IQ

interquartile range, n number. ‡ VAP-related clinical deterioration was defined as an increase in clinical pulmonary infection score (CPIS) > two points

REFERENCES

1. National Nosocomial Infections Surveillance (NNIS) System Report, data summary from January 1992 through June 2004, issued October 2004. *American journal of infection control* 2004; 32(8): 470-485.
2. Vincent JL, Bihari DJ, Suter PM, Bruining HA, White J, Nicolas-Chanoin MH, Wolff M, Spencer RC, Hemmer M. The prevalence of nosocomial infection in intensive care units in Europe. Results of the European Prevalence of Infection in Intensive Care (EPIC) Study. EPIC International Advisory Committee. *Jama* 1995; 274(8): 639-644.
3. Kollef MH. Ventilator-associated pneumonia. A multivariate analysis. *Jama* 1993; 270(16): 1965-1970.
4. Chastre J, Fagon JY. Ventilator-associated pneumonia. *American journal of respiratory and critical care medicine* 2002; 165(7): 867-903.
5. Rello J, Ollendorf DA, Oster G, Vera-Llonch M, Bellm L, Redman R, Kollef MH. Epidemiology and outcomes of ventilator-associated pneumonia in a large US database. *Chest* 2002; 122(6): 2115-2121.
6. Warren DK, Shukla SJ, Olsen MA, Kollef MH, Hollenbeak CS, Cox MJ, Cohen MM, Fraser VJ. Outcome and attributable cost of ventilator-associated pneumonia among intensive care unit patients in a suburban medical center. *Critical care medicine* 2003; 31(5): 1312-1317.
7. Soo Hoo GW, Wen YE, Nguyen TV, Goetz MB. Impact of clinical guidelines in the management of severe hospital-acquired pneumonia. *Chest* 2005; 128(4): 2778-2787.
8. Porzecanski I, Bowton DL. Diagnosis and treatment of ventilator-associated pneumonia. *Chest* 2006; 130(2): 597-604.
9. Michaud S, Suzuki S, Harbarth S. Effect of design-related bias in studies of diagnostic tests for ventilator-associated pneumonia. *American journal of respiratory and critical care medicine* 2002; 166(10): 1320-1325.
10. Koenig SM, Truwit JD. Ventilator-associated pneumonia: diagnosis, treatment, and prevention. *Clinical microbiology reviews* 2006; 19(4): 637-657.
11. Rello J, Gallego M, Mariscal D, Sonora R, Valles J. The value of routine microbial investigation in ventilator-associated pneumonia. *American journal of respiratory and critical care medicine* 1997; 156(1): 196-200.
12. Fagon JY, Chastre J, Wolff M, Gervais C, Parer-Aubas S, Stephan F, Similowski T, Mercat A, Diehl JL, Sollet JP, Tenaillon A. Invasive and noninvasive strategies for management of suspected ventilator-associated pneumonia. A randomized trial. *Annals of internal medicine* 2000; 132(8): 621-630.
13. Kollef MH, Sherman G, Ward S, Fraser VJ. Inadequate antimicrobial treatment of infections: a risk factor for hospital mortality among critically ill patients. *Chest* 1999; 115(2): 462-474.
14. Trouillet JL, Chastre J, Vuagnat A, Joly-Guillou ML, Combaux D, Dombret MC, Gibert C. Ventilator-associated pneumonia caused by potentially drug-resistant bacteria. *American journal of respiratory and critical care medicine* 1998; 157(2): 531-539.
15. Ibrahim EH, Ward S, Sherman G, Schaiff R, Fraser VJ, Kollef MH. Experience with a clinical guideline for the treatment of ventilator-associated pneumonia. *Critical care medicine* 2001; 29(6): 1109-1115.
16. Chastre J, Wolff M, Fagon JY, Chevret S, Thomas F, Wermert D, Clementi E, Gonzalez J, Jusserand D, Asfar P, Perrin D, Fieux F, Aubas S. Comparison of 8 vs 15 days of antibiotic therapy for ventilator-associated pneumonia in adults: a randomized trial. *Jama* 2003; 290(19): 2588-2598.
17. Singh N, Rogers P, Atwood CW, Wagener MM, Yu VL. Short-course empiric antibiotic therapy for patients with pulmonary infiltrates in the intensive care unit. A proposed solution for indiscriminate antibiotic prescription. *American journal of respiratory and critical care medicine* 2000; 162(2 Pt 1): 505-511.
18. Dennesen PJ, van der Ven AJ, Kessels AG, Ramsay G, Bonten MJ. Resolution of infectious parameters after antimicrobial therapy in patients with ventilator-associated pneumonia. *American journal of respiratory and critical care medicine* 2001; 163(6): 1371-1375.

19. Kollef MH. Microbiological diagnosis of ventilator-associated pneumonia - Using the data to optimize clinical outcomes. *American journal of respiratory and critical care medicine* 2008; 173: 1182-1183.
20. Kollef MH. Providing appropriate antimicrobial therapy in the intensive care unit: surveillance vs. de-escalation. *Critical care medicine* 2006; 34(3): 903-905.
21. Guidelines for the management of adults with hospital-acquired, ventilator-associated, and healthcare-associated pneumonia. *American journal of respiratory and critical care medicine* 2005; 171(4): 388-416.
22. Morrow LE, Shorr AF. The seven deadly sins of ventilator-associated pneumonia. *Chest* 2008; 134(2): 225-226.
23. Muller B, Becker KL, Schachinger H, Rickenbacher PR, Huber PR, Zimmerli W, Ritz R. Calcitonin precursors are reliable markers of sepsis in a medical intensive care unit. *Crit Care Med* 2000; 28(4): 977-983.
24. Becker KL, Nylen ES, White JC, Muller B, Snider RH, Jr. Clinical review 167: Procalcitonin and the calcitonin gene family of peptides in inflammation, infection, and sepsis: a journey from calcitonin back to its precursors. *J Clin Endocrinol Metab* 2004; 89(4): 1512-1525.
25. Luyt CE, Combes A, Reynaud C, Hekimian G, Nieszkowska A, Tonnellier M, Aubry A, Trouillet JL, Bernard M, Chastre J. Usefulness of procalcitonin for the diagnosis of ventilator-associated pneumonia. *Intensive care medicine* 2008; 34(8): 1434-1440.
26. Pelosi P, Barassi A, Severgnini P, Gomiero B, Finazzi S, Merlini G, d'Eril GM, Chiaranda M, Niederman MS. Prognostic role of clinical and laboratory criteria to identify early ventilator-associated pneumonia in brain injury. *Chest* 2008; 134(1): 101-108.
27. Christ-Crain M, Jaccard-Stolz D, Bingisser R, Gencay MM, Huber PR, Tamm M, Muller B. Effect of procalcitonin-guided treatment on antibiotic use and outcome in lower respiratory tract infections: cluster-randomised, single-blinded intervention trial. *Lancet* 2004; 363(9409): 600-607.
28. Christ-Crain M, Stolz D, Bingisser R, Muller C, Miedinger D, Huber PR, Zimmerli W, Harbarth S, Tamm M, Muller B. Procalcitonin guidance of antibiotic therapy in community-acquired pneumonia: a randomized trial. *American journal of respiratory and critical care medicine* 2006; 174(1): 84-93.
29. Stolz D, Christ-Crain M, Bingisser R, Leuppi J, Miedinger D, Muller C, Huber P, Muller B, Tamm M. Antibiotic treatment of exacerbations of COPD: a randomized, controlled trial comparing procalcitonin-guidance with standard therapy. *Chest* 2007; 131(1): 9-19.
30. Nobre V, Harbarth S, Graf JD, Rohner P, Pugin J. Use of procalcitonin to shorten antibiotic treatment duration in septic patients: a randomized trial. *American journal of respiratory and critical care medicine* 2008; 177(5): 498-505.
31. Stolz D, Stulz A, Muller B, Gratwohl A, Tamm M. BAL neutrophils, serum procalcitonin, and C-reactive protein to predict bacterial infection in the immunocompromised host. *Chest* 2007; 132(2): 504-514.
32. Altman DG, Schulz KF, Moher D, Egger M, Davidoff F, Elbourne D, Gotzsche PC, Lang T. The revised CONSORT statement for reporting randomized trials: explanation and elaboration. *Ann Intern Med* 2001; 134(8): 663-694.
33. <http://www.controlled-trials.com/ISRCTN61015974>. [cited; Available from:
34. Nylen ES, Muller B, Becker KL, Snyder RH. The future diagnostic role of procalcitonin levels: the need for improved sensitivity. *Clin Infect Dis* 2003; 36: 823-824.
35. Kollef MH, Morrow LE, Niederman MS, Leeper KV, Anzueto A, Benz-Scott L, Rodino FJ. Clinical characteristics and treatment patterns among patients with ventilator-associated pneumonia. *Chest* 2006; 129(5): 1210-1218.
36. Luna CM, Blanzaco D, Niederman MS, Matarucco W, Baredes NC, Desmery P, Palizas F, Menga G, Rios F, Apezteguia C. Resolution of ventilator-associated pneumonia: prospective evaluation of the clinical pulmonary infection score as an early clinical predictor of outcome. *Critical care medicine* 2003; 31(3): 676-682.

37. Shorr AF, Sherner JH, Jackson WL, Kollef MH. Invasive approaches to the diagnosis of ventilator-associated pneumonia: a meta-analysis. *Critical care medicine* 2005; 33(1): 46-53.
38. Rello J, Vidaur L, Sandiumenge A, Rodriguez A, Gualis B, Boque C, Diaz E. De-escalation therapy in ventilator-associated pneumonia. *Critical care medicine* 2004; 32(11): 2183-2190.
39. Wanner GA, Keel M, Steckholzer U, Beier W, Stocker R, Ertel W. Relationship between procalcitonin plasma levels and severity of injury, sepsis, organ failure, and mortality in injured patients. *Critical care medicine* 2000; 28(4): 950-957.
40. Harbarth S, Holeckova K, Froidevaux C, Pittet D, Ricou B, Grau GE, Vadas L, Pugin J. Diagnostic value of procalcitonin, interleukin-6, and interleukin-8 in critically ill patients admitted with suspected sepsis. *American journal of respiratory and critical care medicine* 2001; 164(3): 396-402.
41. Luyt CE, Guerin V, Combes A, Trouillet JL, Ayed SB, Bernard M, Gibert C, Chastre J. Procalcitonin kinetics as a prognostic marker of ventilator-associated pneumonia. *American journal of respiratory and critical care medicine* 2005; 171(1): 48-53.
42. Seligman R, Meisner M, Lisboa TC, Hertz FT, Filippin TB, Fachel JM, Teixeira PJ. Decreases in procalcitonin and C-reactive protein are strong predictors of survival in ventilator-associated pneumonia. *Critical care (London, England)* 2006; 10(5): R125.
43. Suprin E, Camus C, Gacouin A, Le Tulzo Y, Lavoue S, Feuillu A, Thomas R. Procalcitonin: a valuable indicator of infection in a medical ICU? *Intensive care medicine* 2000; 26(9): 1232-1238.