

# European Respiratory Society Annual Congress 2012

**Abstract Number:** 735

**Publication Number:** P1455

**Abstract Group:** 7.3. Cystic Fibrosis

**Keyword 1:** Exercise **Keyword 2:** Physical activity **Keyword 3:** Physiology

**Title:** Habitual physical activity in cystic fibrosis

Dr. Daniela 5828 Savi danielasavi1@virgilio.it MD <sup>1</sup>, Dr. Mattia 5829 Internullo m.internullo@virgilio.it MD <sup>2</sup>, Dr. Giuseppe 5830 Cimino ciminolo@tiscali.it MD <sup>1</sup>, Dr. Elena 5831 Angelici elena.angelici@uniroma1.it MD <sup>2</sup>, Dr. Serena 5832 Quattrucci serena.quattrucci@uniroma1.it MD <sup>1</sup>, Dr. Patrizia 5842 Troiani patrizia.troiani@fastwebnet.it MD <sup>1</sup>, Prof. Salvatore 5845 Cucchiara salvatore.cucchiara@uniroma1.it MD <sup>1</sup> and Prof. Paolo 5853 Palange paolo.palange@uniroma1.it MD <sup>2</sup>. <sup>1</sup> Regional Cystic Fibrosis Centre - Pediatrics Department, Sapienza - University of Rome, Italy, 00161 and <sup>2</sup> Department of Public Health and Infectious Diseases, Sapienza - University of Rome, Italy, 00161 .

**Body:** Introduction. Habitual physical activity has important clinical implications in Cystic Fibrosis (CF) and has therapeutic effects on sputum clearance, respiratory muscle strength and quality of life. While recent studies have shown that aerobic fitness is related to survival, that physically active CF patients have higher peak oxygen uptake (VO<sub>2</sub>peak) and lower rate of decline of FEV<sub>1</sub>, less is known about the relationship between physical activity levels and patients's functional status, including exercise tolerance. Methods. Eleven stable CF patients (mean (SD) age 32 (9) yrs; FEV<sub>1</sub> 2.7 (0.8) l; IC 3.8 (0.9) l), were studied at rest, during symptom-limited incremental exercise test (CPET) and during 6MWT. We assessed daily physical activity using both SW (SenseWear, SW) that subjects wore for 4 consecutive days from waking until going to bed including weekends, and the Habitual Activity Estimation Scale Questionnaire (HAESQ). Results. There was no difference between the weekdays and weekends recordings in any activity variable. We found no agreement in physical activity measured by SW vs HAESQ. By contrast, we found a close relationship between SW step count and FEV<sub>1</sub> (r<sup>2</sup>=0.9; p<0.02) and SW energy expenditure vs VO<sub>2</sub>peak (r<sup>2</sup>=0.8; p<0.05) at CPET. Distance at 6MWT was not correlated to any physical activity variables, either at SW or HAESQ. Conclusions. Adults CF patients have similar habitual physical activity levels at weekdays and weekends. Activity levels measured by the subjective methods and by the objective monitoring, i.e., SW vs HAESQ, seems to provide different information on the level of daily physical activity. Physical activity is related to degree of airflow obstruction and to the maximum exercise capacity.