

**TITLE PAGE**

**Intrabronchial activated protein C enhances lipopolysaccharide-induced pulmonary responses**

Liesbeth M. Kager,<sup>1,2</sup> J. Daan de Boer,<sup>1</sup> Paul Bresser,<sup>3,7</sup> Jaring S. van der Zee,<sup>3,7</sup> Sacha Zeerleder,<sup>4,8</sup> Joost C. M. Meijers,<sup>5</sup> Cornelis van 't Veer,<sup>1,2</sup> and Tom van der Poll,<sup>1,2,6</sup>

Academic Medical Center/University of Amsterdam, The Netherlands, <sup>1</sup>Center of Experimental and Molecular Medicine, <sup>2</sup>Center for Infection and Immunity Amsterdam, <sup>3</sup>Department of Pulmonology, <sup>4</sup>Department of Hematology, <sup>5</sup>Department of Experimental Vascular Medicine, <sup>6</sup>Division of Infectious Diseases.

Onze Lieve Vrouwe Gasthuis, Amsterdam, The Netherlands, <sup>7</sup>Department of Pulmonology, Sanquin Research at CLB and Landsteiner Laboratory of the AMC, Amsterdam, The Netherlands, <sup>8</sup>Department of Immunopathology.

**Corresponding author & requests for reprints:**

Liesbeth M. Kager, MD  
Center for Experimental and Molecular Medicine  
Academic Medical Center, University of Amsterdam  
Meibergdreef 9, Room G2-130  
1105 AZ Amsterdam, The Netherlands  
Fax +31 20 697 7192,  
Telephone +31 20 566 5910  
E-mail: l.m.kager@amc.uva.nl

**Running title:** Effects of intrabronchial APC after LPS challenge

**Keywords:** lung, pneumonia, endotoxin, lipopolysaccharide, activated protein C, inflammation, cytokines, chemokines, coagulation, fibrinolysis

**Word count abstract:** 197

**Word count (body of manuscript):** 3033

## **ABSTRACT**

Intravenous administration of activated protein C (APC) inhibits coagulation and inflammation in the lungs of humans and animals. Investigations in rodents demonstrated that direct intrapulmonary delivery of APC also exerts anticoagulant and anti-inflammatory effects. The effect of intrabronchial administration of recombinant human (rh)APC on lipopolysaccharide (LPS)-induced hemostatic and inflammatory alterations in the bronchoalveolar space of humans was studied.

Eight subjects received rhAPC via intrabronchial instillation by bronchoscope, while in a contralateral subsegment subjects received saline; all subjects were challenged bilaterally with LPS in the same lung subsegments. Four additional subjects received rhAPC (75 µg), with saline as control in the contralateral subsegment, while they were bilaterally 'challenged' with saline. After 6 hours a bronchoalveolar lavage was performed and coagulation and inflammatory parameters were measured.

rhAPC enhanced LPS-induced coagulation activation in the bronchoalveolar space, when compared with control side. In addition, rhAPC amplified LPS-induced proinflammatory responses, as indicated by higher concentrations of cytokines and chemokines. RhAPC alone did not have procoagulant or proinflammatory effects.

Locally administered rhAPC has unexpected procoagulant and proinflammatory effects in LPS-challenged lung subsegments. These data argue against a role for intrapulmonary delivery of rhAPC as a treatment strategy for lung inflammatory disorders in humans. (ClinicalTrials.gov number: NCT00943267; Nederlands Trial register number: NTR1544).

### **Keywords:**

coagulation, fibrinolysis, inflammation, lung, recombinant activated protein C

## INTRODUCTION

The airways represent a body site in which procoagulant and anticoagulant mechanisms can be initiated and regulated locally[1, 2]. Activated protein C (APC), an important anticoagulant, has been shown to have a reduced activity in the bronchoalveolar space in various lung inflammatory diseases [3-11]. Diminished APC levels may contribute to the disturbed hemostatic equilibrium in the lungs of patients suffering from these conditions, shifting the physiological anticoagulant environment into a net procoagulant state. Besides its anticoagulant effects, APC has been implicated in a variety of anti-inflammatory and/or cytoprotective effects, characterized by down-regulation of pro-inflammatory pathways, upregulation of anti-inflammatory pathways and inhibition of neutrophil activity[12, 13].

We recently demonstrated that intrabronchial instillation of lipopolysaccharide (LPS), a constituent of the cell wall of gram-negative bacteria, can reproduce hemostatic alterations found in inflammatory lung disorders, including a reduction in APC levels in the lungs of healthy volunteers[14, 15]. This human model of intrabronchial LPS delivery also has been used to study the *in vivo* effects of intravenous recombinant human (rh)APC treatment on pulmonary inflammation and coagulation[15, 16]. Intravenous rhAPC reduced LPS-induced neutrophil accumulation in the bronchoalveolar space[16] and inhibited local activation of coagulation[15]. Concurrently, LPS-induced suppression of fibrinolysis was partially prevented by intravenous rhAPC[15].

When administered intravenously, the strong anticoagulant effects of rhAPC may contribute to serious systemic side effects such as intracranial hemorrhages[17-20], which may be circumvented by local administration of rhAPC. Although preclinical animal studies have shown that administration of APC via the airways exerts local anti-inflammatory and anticoagulant effects[21-24], no studies exist that evaluated the effects of direct rhAPC administration into the lungs of humans. The present study therefore sought to determine the

feasibility of local, intrabronchial administration of rhAPC to inhibit LPS-induced lung inflammation and coagulation in humans.

## METHODS

### *Study subjects & materials*

Twelve nonsmoking males (age,  $23.2 \pm 1.1$  year) were enrolled. Screening, consisting of a questionnaire, physical examination, routine blood and urine investigation, electrocardiogram and spirometry did not reveal any abnormality. The protocol was approved by the institutional Medical Ethics Review Committee. All subjects provided written informed consent before enrollment. LPS was derived from *Escherichia coli* O:113 (Reference Endotoxin, CC-RE-Lot 3, National Institutes of Health, Bethesda, MD). RhAPC (drotrecogin alpha (activated), Eli Lilly, Indianapolis, IN) was purchased from a commercial supplier.

### *Study design*

The study was set-up as a single blind intervention study with a dose-escalation and a follow-up phase. A bilateral challenge of two contralateral lung subsegments was performed by intrabronchial instillation of LPS (4 ng/kg) by bronchoscopy, diluted in 10 mL sterile saline. In one lung subsegment the instillation of LPS was followed immediately by administration of rhAPC (in 10 mL) in the same lung subsegment, whereas in the contralateral lung subsegment LPS administration was followed by instillation of 10 mL saline alone. The subjects were randomized to left or right lungs for rhAPC instillation. In the dose-escalating phase of the study, 5-fold increases in the rhAPC dose were intended to be given in cohorts of four subjects per dose, starting at 15  $\mu$ g. The predefined primary end-point was to reach a 30% decrease in thrombin-antithrombin complex (TATc) concentrations in the rhAPC-treated lung 6 hours after intrabronchial LPS challenge relative to the contralateral LPS challenged side; this end-point was based on the capacity of intravenous rhAPC to inhibit the rise in LPS-induced TATc concentrations in bronchoalveolar lavage fluid (BALF) by approximately 30% in our previous study[15]. Per study design, the sample size of the rhAPC dose group

that would reach this primary end-point would be increased to a total of 12 subjects in the follow-up phase of the study. After completion of the first two cohorts in the dose-escalation phase (15 and 75 µg; N=4 for each dose), we decided to stop the study because it was considered highly unlikely that the primary study end-point could be reached. Instead, 4 additional subjects were included who received 75 µg rhAPC in one lung subsegment without previous intrabronchial LPS challenge; in these volunteers saline was given as a control in the contralateral lung subsegment.

### *Methods*

A bilateral bronchoalveolar lavage (BAL) was performed 6 hours post-challenge in a standardized fashion[14, 25]. Citrated and heparinised blood was taken by venipuncture before the first (t=0) and the second (t=6 hours) bronchoscopy. The assays used are described in the on-line supplement.

### *Statistical analysis*

Values are expressed as means ( $\pm$  SEM). Paired *t* tests were used to establish significance between data sets. *P*-values < 0.05 were considered statistically significant.

## RESULTS

### *Clinical signs and systemic inflammatory response*

Instillation of LPS in two contralateral lung subsegments (combined with rhAPC in one lung segment) was well tolerated. A modest rise in body temperature was recorded 6 hours after instillation of LPS (from  $36.0 \pm 0.2^\circ\text{C}$  at T=0h to  $36.6 \pm 0.1^\circ\text{C}$  at T=6h,  $P<0.05$ ). Intrabronchial LPS caused a rise in neutrophil counts in blood (from  $2.3 \pm 0.2$  to  $6.6 \pm 0.8 \times 10^9/\text{L}$ ,  $P<0.001$ ), while in the saline challenged group no significant rises in neutrophils counts were seen (from  $2.0 \pm 0.2$  to  $4.1 \pm 0.8 \times 10^9/\text{L}$ ). Finally, intrabronchial instillation of either LPS or saline, with addition of rhAPC in one lung subsegment, was not associated with changes in plasma concentrations of TATc, D-dimer, tumor necrosis factor (TNF)- $\alpha$  or interleukin (IL)-6 (data not shown).

### *Bronchoalveolar APC concentrations*

Six hours after intrabronchial instillation of rhAPC, APC was measurable in all BAL fluid (BALF) samples obtained from the APC-treated side: APC levels were  $0.33 \pm 0.14$  and  $1.60 \pm 0.57$  ng/mL in LPS-challenged subjects treated with 15 and 75  $\mu\text{g}$  rhAPC, respectively ( $P=0.07$  and  $<0.05$  for lung subsegments treated with 15 and 75  $\mu\text{g}$  rhAPC versus control lung subsegments; Figure 1A-B). In the saline-challenged group treated with 75  $\mu\text{g}$  rhAPC, BALF APC levels were  $3.2 \pm 1.0$  ng/mL ( $P<0.05$ ; Figure 1C). APC levels in the 75  $\mu\text{g}$  rhAPC-challenged subsegments were not different between LPS-challenged and saline-challenged subjects. APC could not be detected in any of the control lung subsegments (i.e. not administered with rhAPC).

### *Intrabronchial rhAPC enhances bronchoalveolar coagulation activation after LPS challenge*

The intrabronchial concentrations of TATc have been used to determine the extent of bronchoalveolar coagulation activation in patients with acute lung injury and/or pneumonia [7, 8, 10]. Intrabronchial instillation of LPS in healthy humans is also associated with a rise in

TATc levels in BALF[14, 15]. An earlier study showed that intravenously administered rhAPC could reduce LPS-induced elevations in TATc-concentrations in BALF by approximately 30%[15]. The main objective of the current study was to determine whether intrabronchially instilled rhAPC was able to inhibit LPS-induced TATc elevation in BALF and if so whether this anticoagulant effect was accompanied by anti-inflammatory effects. Considering the results of our previous investigation using intravenous rhAPC[15], we here predefined the primary endpoint of the current study as establishing the intrabronchial rhAPC dose that induced a 30% reduction in LPS-induced TATc concentrations in BALF. We chose to administer LPS in two contralateral lung subsegments, thereby eliminating inter-individual variation in LPS responsiveness and allowing every subject to serve as his own control with regard to rhAPC effects. Much to our surprise, intrabronchially instilled rhAPC, given at either 15 µg or 75 µg, induced an increase rather than a reduction in BALF TATc concentrations after LPS-challenge when compared with the contralateral side not treated with rhAPC (rhAPC 15 µg:  $13.5 \pm 5.2$  versus  $9.8 \pm 4.8$  µg/L at the contralateral side,  $P < 0.01$ , Figure 2A; rhAPC 75 µg:  $19.4 \pm 4.2$  versus  $2.9 \pm 0.8$  µg/L at the contralateral side,  $P < 0.05$ , Figure 2B). As these results made it highly unlikely that we could reach our predefined endpoint, we decided not to proceed with another 5-fold increase in rhAPC dose, but instead, added an extra study group without LPS challenge, seeking to determine whether intrabronchial rhAPC per se elicited a procoagulant response. Thus, subjects received a bilateral challenge with saline, combined with rhAPC (75 µg) on one side. These studies revealed no procoagulant effect of rhAPC: BALF TATc levels were similar in lung subsegments instilled with rhAPC ( $1.6 \pm 1.4$  µg/L) when compared with the contralateral side ( $2.4 \pm 1.2$  µg/L, Figure 2C). Protein C activity, total and free protein S, soluble thrombomodulin, protein C inhibitor and von Willebrand factor antigen concentrations were all below detection limits in BALF.

#### *Effect of intrabronchial rhAPC on fibrinolysis after LPS challenge*

Evidence derived from *in vitro* investigations indicates that APC may stimulate fibrinolysis by inhibiting PAI-1[26]. We measured levels of PAI-1, tissue-type plasminogen activator (tPA)

and plasmin- $\alpha$ 2-antiplasmin complexes. While intrabronchial rhAPC given at a dose of 15  $\mu$ g did not influence LPS-induced PAI-1 release in BALF ( $P=0.57$ , Figure 3A), rhAPC administered at 75  $\mu$ g increased LPS-induced PAI-1 BALF levels in 3 of 4 subjects ( $192.5 \pm 42.1$  versus  $55.0 \pm 17.1$  pg/mL at the contralateral side,  $P=0.07$ , Figure 3B). Notably, intrabronchial rhAPC also tended to increase PAI-1 levels in BALF from subjects who did not receive LPS ( $70.0 \pm 45.3$  versus  $7.5 \pm 7.5$  pg/mL at the contralateral side,  $P=0.26$ , Figure 3C). Neither LPS nor rhAPC influenced BALF concentrations of plasmin- $\alpha$ 2-antiplasmin complexes, whereas tPA levels were below detection limit in all BALF samples (data not shown).

*Intrabronchial rhAPC is associated with enhanced concentrations of tissue-factor pathway inhibitor (TFPI) after LPS challenge*

Recent data suggest that APC can have a procoagulant effect by virtue of its proteolytic activity. APC was reported to shed the Kunitz-1 domain from TFPI, resulting in increased tissue factor activity and consequently, coagulation activation[27]. We measured levels of TFPI in BALF. While intrabronchial rhAPC given at a dose of 15  $\mu$ g did not influence LPS-induced TFPI release in BALF (Figure 4A), rhAPC administered at 75  $\mu$ g increased LPS-induced TFPI BALF levels in all 4 subjects ( $48.1 \pm 9.9$  versus  $87.9 \pm 14.8$  pg/mL at the contralateral side,  $P<0.05$ , Figure 4B). Intrabronchial rhAPC did not change levels of TFPI in BALF from subjects who did not receive LPS (Figure 4C).

*Intrabronchial rhAPC increases leukocyte recruitment after LPS administration*

Intravenous rhAPC has been reported to reduce neutrophil recruitment into the bronchoalveolar space upon intrabronchial instillation of LPS in healthy humans[16]. In contrast, intrabronchial rhAPC treatment increased the total number of white blood cells in BALF 6 hours after bilateral LPS-challenge, although the differences with the contralateral control side did not reach statistical significance (APC 15  $\mu$ g:  $33.6 \pm 14.6 \times 10^4$  versus  $18.6 \pm$

4.1x10<sup>4</sup> cells/mL at the contralateral side,  $P=0.25$ , Figure 5A; APC 75 µg: 52.0 ± 22.4x10<sup>4</sup> versus 11.1 ± 1.5x10<sup>4</sup> cells/mL at the contralateral side,  $P=0.13$ , Figure 5B). The rhAPC induced increase in total cell number in lung subsegments challenged with LPS was due to a rise in the number of neutrophils (APC 15 µg: 16.8 ± 9.0 x 10<sup>4</sup> versus 7.5±2.7 x 10<sup>4</sup> neutrophils/mL,  $P=0.25$ , Figure 5D; APC 75 µg: 34.1 ± 15.8x10<sup>4</sup> versus 1.5 ± 0.6x10<sup>4</sup> neutrophils/mL,  $P=0.13$ , Figure 5E). Intrabronchial rhAPC did not impact cell counts in lung subsegments not challenged with LPS (Figure 5C,F).

*Intrabronchial rhAPC enhances the local release of several cytokines and chemokines after LPS instillation*

To investigate the effect of local rhAPC administration on LPS-induced inflammatory mediator release we measured BALF concentrations of 48 cytokines, chemokines and growth factors. The results of these analyses are shown in table E1 in the on-line data supplement. In the lower rhAPC dose cohort (15 µg), rhAPC significantly enhanced LPS-induced release of IL-6, CCL8 and CXCL6 while not influencing the levels of other mediators (Table E1). In the rhAPC 75 µg cohort, rhAPC significantly enhanced LPS-induced TNF-α, CCL4, CCL8 and TNF-related apoptosis-inducing ligand (TRAIL) concentrations (Table E1, Figure 5). In addition, IL-1β, CCL14a, CCL20, CXCL7, CXCL9 and CXCL11 showed a strong trend toward higher levels in this higher rhAPC dose cohort ( $P = 0.052, 0.081, 0.078, 0.075, 0.092$  and  $0.074$  respectively; Table E1, Figure 6). RhAPC given at 75 µg without LPS did not induce the release of inflammatory mediators in the lung with the exception of a modest rise in CCL15 concentrations (Table E1). Taken together these results show that intrabronchial instillation of rhAPC in LPS-challenged lung subsegments enhanced the release of multiple proinflammatory mediators.

## DISCUSSION

Several preclinical studies have shown that APC may protect the lung from injury caused by inflammation[28]. Moreover, intravenous administration of rhAPC was reported to exert anticoagulant and anti-inflammatory effects in the bronchoalveolar space of healthy humans challenged with LPS in a lung subsegment by bronchoscope[15, 16]. Animal studies demonstrated that intrabronchially instilled APC exerts local anticoagulant and anti-inflammatory effects[21-24, 29-32]. Considering the bleeding risk associated with intravenous APC, we here for the first time investigated the potential of intrabronchial rhAPC administration to inhibit coagulation and inflammation in the human lung, arguing that this route of delivery would circumvent serious side effects [17-20]. For this we used the established human model of lung inflammation induced by intrabronchial instillation of LPS via a bronchoscope[14-16, 25]. The main finding of our study was remarkable: intrabronchially administered rhAPC induced a procoagulant and proinflammatory response in lung subsegments challenged with LPS rather than the expected inhibitory effects on coagulation and inflammation. Our results from subjects not challenged with LPS via the airways further suggest that rhAPC augments LPS effects in the human lung, while not eliciting detectable inflammation or coagulation per se.

Our study had a small sample-size. As a consequence, although rhAPC clearly enhanced the LPS-induced release of biomarkers of coagulation and inflammation, the differences with the placebo control side often did not reach statistical significance. Of note, however, the current study was designed to demonstrate an inhibitory effect of rhAPC on bronchoalveolar coagulation and inflammation. After completion of the rhAPC 75 µg cohort, we considered it unlikely that the predefined endpoint (i.e. a 30% reduction in BALF TATc levels) could be reached. Therefore, we decided to prematurely terminate the study. In addition, we considered it not ethical to enlarge the sample size of our cohorts in order to strengthen the statistical power of our study, arguing that procoagulant and proinflammatory effects of

locally administered rhAPC would not be of clinical value for patient groups and not wishing to expose additional healthy subjects to invasive procedures.

In the earlier study, in which intravenous rhAPC was found to inhibit coagulation and neutrophil influx in the same human LPS-induced lung inflammation model[15, 16], APC concentrations in BALF were approximately 15 ng/mL as measured 2 hours after discontinuation of the rhAPC infusion (24 µg/kg/hour)[15]. Although BALF APC levels detected after intravenous rhAPC infusion cannot be readily translated to a feasible rhAPC dose to be administered in the bronchoalveolar space, we conservatively started the dose-escalation study with 15 µg of intrabronchial rhAPC, taking into account the dilution caused by the BAL procedure and an estimated recovery of 50-60%. BALF APC levels measured in the current study, 6 hours after instillation, were considerably lower than in the previous study using intravenous APC administration[15] (mean levels of 0.33 ng/mL and 1.60 ng/mL in the rhAPC 15 µg and 75 µg cohorts respectively). Arguably, BALF APC concentrations were higher directly after rhAPC administration. Clearance of APC from the bronchoalveolar space is difficult to investigate in humans in light of the invasive procedure needed; nonetheless, evidence suggests that APC is slowly cleared from the alveolar compartment. Indeed, while the majority of patients with sepsis who are treated with intravenous rhAPC, do not have detectable APC in plasma 2 hours after discontinuation of the infusion[33], APC could still be recovered from BALF of healthy humans in whom a similar rhAPC infusion was stopped 2 hours earlier[15]. Hence, although BALF APC concentrations measured in the present (intrabronchial rhAPC) and previous (intravenous rhAPC)[15] investigation cannot be compared easily, these combined data suggest that the APC levels achieved in BALF here were likely within the same or lower range of those attained after intravenous rhAPC infusion.

How can the apparently differential effects of intravenous rhAPC[15] and intrabronchial rhAPC in LPS-induced human lung inflammation be explained? APC can exert anti-inflammatory, anti-apoptotic and barrier protective signals in endothelial cells via protease

activated receptor (PAR)1 by a mechanism that requires binding of APC to the endothelial cell protein C receptor (EPCR)[34]. Cytoprotective anti-inflammatory APC-PAR1 signaling has also been demonstrated in other cell types, in particular dendritic cells (EPCR dependent[35]) and macrophages (CD11/CD18 dependent[35, 36]). Furthermore, APC directly binds to activated  $\alpha_3\beta_1$ ,  $\alpha_5\beta_1$ , and  $\alpha_v\beta_3$  integrins and this interaction was essential for APC-induced inhibition of neutrophil extravasation into the bronchoalveolar space of mice[37]. These “cytoprotective” and/or anti-inflammatory APC effects *in vivo* likely depend on which cell types are affected and thereby on the route of rhAPC administration. Furthermore, the distribution of receptors involved in these APC effects may vary in different body compartments. In addition, recent data suggest that APC can have a procoagulant effect by virtue of its proteolytic activity: APC was reported to shed the Kunitz-1 domain from TFPI, resulting in increased tissue factor activity and consequently, coagulation activation[27]. In this study we measured TFPI levels in BALF and showed that administration of rhAPC in the 75  $\mu$ g cohort was associated with significantly enhanced TFPI levels. We hypothesize that increased levels of TFPI may have been a result of the pro-inflammatory state due to hAPC administration. Of note, the TFPI assay used detects both full-length and cleaved TFPI. Hence, it is possible that the elevated TFPI levels were the result of cleavage of the intact protein by rhAPC, thereby contributing to a procoagulant effect. We do not have a clear explanation for the discrepancy between our current findings and earlier reports on the effect of intrapulmonary delivery of APC in mice[21-24], although differences in the exact way of administration, the dose and species studied may be involved.

Our study does not provide a clear mechanism underlying the unexpected proinflammatory and procoagulant effect of rhAPC in the human lung. We tried to reproduce the effects of APC using a mixed cell culture system using respiratory epithelial cells (A549) and either

primary human alveolar macrophages or macrophage-like THP-1 cells but were unsuccessful (data not shown).

In conclusion, this is the first study on the effects of intrabronchially administered rhAPC in a controlled lung inflammation model in humans. Our data show that locally administered rhAPC has procoagulant and proinflammatory effects in lung segments exposed to LPS. These data argue against a role for intrapulmonary delivery of rhAPC as a treatment strategy for lung inflammatory disorders in humans.

### **Acknowledgements**

The authors thank A. F. de Vos, D. Kruijswijk and D. C. Blok for their assistance during the study days, G. van Mierlo, W. F. Kopatz, M. A. Weijne, L. M. Leverink and J. A. Marquart for assistance in performing the coagulation measurements and E. M. Kemper, in-hospital pharmacist, for assistance in preparation of the study medication.

### **Competing interests**

None of the authors has competing interests to declare.

### **Funding**

This study was supported by research grants of ZonMW (to L. M. K., grant number: 92003504), the Netherlands Asthma Foundation (J.D.d.B; project 3.2.08.009) and the Stichting BeGeTu (to L. M. K.).

## Reference List

- [1] Levi M, Schultz MJ, Rijneveld AW, van der Poll T. Bronchoalveolar coagulation and fibrinolysis in endotoxemia and pneumonia. *Crit Care Med* 2003; **31**:S238-S242.
- [2] Wygrecka M, Jablonska E, Guenther A, Preissner KT, Markart P. Current view on alveolar coagulation and fibrinolysis in acute inflammatory and chronic interstitial lung diseases. *Thromb Haemost* 2008; **99**:494-501.
- [3] Choi G, Schultz MJ, Levi M, van der Poll T, Millo JL, Garrard CS. Protein C in pneumonia. *Thorax* 2005; **60**:705-706.
- [4] Hataji O, Taguchi O, Gabazza EC, Yuda H, Fujimoto H, Suzuki K *et al.* Activation of protein C pathway in the airways. *Lung* 2002; **180**:47-59.
- [5] Imokawa S, Sato A, Hayakawa H, Kotani M, Urano T, Takada A. Tissue factor expression and fibrin deposition in the lungs of patients with idiopathic pulmonary fibrosis and systemic sclerosis. *Am J Respir Crit Care Med* 1997; **156**:631-636.
- [6] Schouten M, VAN DE Pol MA, Levi M, van der Poll T, van der Zee JS. Early activation of coagulation after allergen challenge in patients with allergic asthma. *J Thromb Haemost* 2009; **7**:1592-1594.
- [7] Schultz MJ, Millo J, Levi M, Hack CE, Weverling GJ, Garrard CS *et al.* Local activation of coagulation and inhibition of fibrinolysis in the lung during ventilator associated pneumonia. *Thorax* 2004; **59**:130-135.
- [8] Choi G, Schultz MJ, van Till JW, Bresser P, van der Zee JS, Boermeester MA *et al.* Disturbed alveolar fibrin turnover during pneumonia is restricted to the site of infection. *Eur Respir J* 2004; **24**:786-789.
- [9] Ware LB, Fang X, Matthay MA. Protein C and thrombomodulin in human acute lung injury. *Am J Physiol Lung Cell Mol Physiol* 2003; **285**:L514-L521.
- [10] Gunther A, Mosavi P, Heinemann S, Ruppert C, Muth H, Markart P *et al.* Alveolar fibrin formation caused by enhanced procoagulant and depressed fibrinolytic capacities in severe pneumonia. Comparison with the acute respiratory distress syndrome. *Am J Respir Crit Care Med* 2000; **161**:454-462.
- [11] Arndt PG, Young SK, Worthen GS. Regulation of lipopolysaccharide-induced lung inflammation by plasminogen activator Inhibitor-1 through a JNK-mediated pathway. *J Immunol* 2005; **175**:4049-4059.
- [12] Mosnier LO, Zlokovic BV, Griffin JH. The cytoprotective protein C pathway. *Blood* 2007; **109**:3161-3172.
- [13] Weiler H. Regulation of inflammation by the protein C system. *Crit Care Med* 2010; **38**:S18-S25.
- [14] Hoogerwerf JJ, de Vos AF, Levi M, Bresser P, van der Zee JS, Draing C *et al.* Activation of coagulation and inhibition of fibrinolysis in the human lung on bronchial instillation of lipoteichoic acid and lipopolysaccharide. *Crit Care Med* 2009; **37**:619-625.
- [15] van der Poll T, Levi M, Nick JA, Abraham E. Activated protein C inhibits local coagulation after intrapulmonary delivery of endotoxin in humans. *Am J Respir Crit Care Med* 2005; **171**:1125-1128.

- [16] Nick JA, Coldren CD, Geraci MW, Poch KR, Fouty BW, O'Brien J *et al.* Recombinant human activated protein C reduces human endotoxin-induced pulmonary inflammation via inhibition of neutrophil chemotaxis. *Blood* 2004; **104**:3878-3885.
- [17] Bernard GR, Vincent JL, Laterre PF, LaRosa SP, Dhainaut JF, Lopez-Rodriguez A *et al.* Efficacy and safety of recombinant human activated protein C for severe sepsis. *N Engl J Med* 2001; **344**:699-709.
- [18] Vincent JL, Bernard GR, Beale R, Doig C, Putensen C, Dhainaut JF *et al.* Drotrecogin alfa (activated) treatment in severe sepsis from the global open-label trial ENHANCE: further evidence for survival and safety and implications for early treatment. *Crit Care Med* 2005; **33**:2266-2277.
- [19] Abraham E, Laterre PF, Garg R, Levy H, Talwar D, Trzaskoma BL *et al.* Drotrecogin alfa (activated) for adults with severe sepsis and a low risk of death. *N Engl J Med* 2005; **353**:1332-1341.
- [20] Marti-Carvajal AJ, Sola I, Lathyris D, Cardona AF. Human recombinant activated protein C for severe sepsis. *Cochrane Database Syst Rev* 2011;CD004388.
- [21] Hofstra JJ, Vlaar AP, Cornet AD, Dixon B, Roelofs JJ, Choi G *et al.* Nebulized anticoagulants limit pulmonary coagulopathy, but not inflammation, in a model of experimental lung injury. *J Aerosol Med Pulm Drug Deliv* 2010; **23**:105-111.
- [22] Cornet AD, Hofstra JJ, Vlaar AP, Van Den Boogaard FE, Roelofs JJ, van der Poll T *et al.* Nebulized anticoagulants limit coagulopathy but not inflammation in pseudomonas aeruginosa-induced pneumonia in rats. *Shock* 2011; **36**:417-423.
- [23] Slofstra SH, Groot AP, Maris NA, Reitsma PH, Cate HT, Spek CA. Inhalation of activated protein C inhibits endotoxin-induced pulmonary inflammation in mice independent of neutrophil recruitment. *Br J Pharmacol* 2006; **149**:740-746.
- [24] Maniatis NA, Letsiou E, Orfanos SE, Kardara M, Dimopoulou I, Nakos G *et al.* Inhaled activated protein C protects mice from ventilator-induced lung injury. *Crit Care* 2010; **14**:R70.
- [25] Hoogerwerf JJ, de Vos AF, Bresser P, van der Zee JS, Pater JM, de BA *et al.* Lung inflammation induced by lipoteichoic acid or lipopolysaccharide in humans. *Am J Respir Crit Care Med* 2008; **178**:34-41.
- [26] Bajzar L, Nesheim ME, Tracy PB. The profibrinolytic effect of activated protein C in clots formed from plasma is TAFI-dependent. *Blood* 1996; **88**:2093-2100.
- [27] Schuepbach RA, Velez K, Riewald M. Activated protein C up-regulates procoagulant tissue factor activity on endothelial cells by shedding the TFPI Kunitz 1 domain. *Blood* 2011; **117**:6338-6346.
- [28] Suzuki K, Gabazza EC, Hayashi T, Kamada H, Adachi Y, Taguchi O. Protective role of activated protein C in lung and airway remodeling. *Crit Care Med* 2004; **32**:S262-S265.
- [29] Yuda H, Adachi Y, Taguchi O, Gabazza EC, Hataji O, Fujimoto H *et al.* Activated protein C inhibits bronchial hyperresponsiveness and Th2 cytokine expression in mice. *Blood* 2004; **103**:2196-2204.


- [30] Yasui H, Gabazza EC, Tamaki S, Kobayashi T, Hataji O, Yuda H *et al.* Intratracheal administration of activated protein C inhibits bleomycin-induced lung fibrosis in the mouse. *Am J Respir Crit Care Med* 2001; **163**:1660-1668.
- [31] Shimizu S, Gabazza EC, Taguchi O, Yasui H, Taguchi Y, Hayashi T *et al.* Activated protein C inhibits the expression of platelet-derived growth factor in the lung. *Am J Respir Crit Care Med* 2003; **167**:1416-1426.
- [32] Waerhaug K, Kuzkov VV, Kuklin VN, Mortensen R, Nordhus KC, Kirov MY *et al.* Inhaled aerosolised recombinant human activated protein C ameliorates endotoxin-induced lung injury in anaesthetised sheep. *Crit Care* 2009; **13**:R51.
- [33] Macias WL, Dhainaut JF, Yan SC, Helterbrand JD, Seger M, Johnson G3 *et al.* Pharmacokinetic-pharmacodynamic analysis of drotrecogin alfa (activated) in patients with severe sepsis. *Clin Pharmacol Ther* 2002; **72**:391-402.
- [34] Danese S, Vetrano S, Zhang L, Poplis VA, Castellino FJ. The protein C pathway in tissue inflammation and injury: pathogenic role and therapeutic implications. *Blood* 2010; **115**:1121-1130.
- [35] Kerschen E, Hernandez I, Zogg M, Jia S, Hessner MJ, Fernandez JA *et al.* Activated protein C targets CD8+ dendritic cells to reduce the mortality of endotoxemia in mice. *J Clin Invest* 2010; **120**:3167-3178.
- [36] Cao C, Gao Y, Li Y, Antalis TM, Castellino FJ, Zhang L. The efficacy of activated protein C in murine endotoxemia is dependent on integrin CD11b. *J Clin Invest* 2010; **120**:1971-1980.
- [37] Elphick GF, Sarangi PP, Hyun YM, Hollenbaugh JA, Ayala A, Biffl WL *et al.* Recombinant human activated protein C inhibits integrin-mediated neutrophil migration. *Blood* 2009; **113**:4078-4085.
- [38] Stephenson DA, Tolti LJ, Beaudin S, Liaw PC. Modulation of monocyte function by activated protein C, a natural anticoagulant. *J Immunol* 2006; **177**:2115-2122.
- [39] White B, Schmidt M, Murphy C, Livingstone W, O'Toole D, Lawler M *et al.* Activated protein C inhibits lipopolysaccharide-induced nuclear translocation of nuclear factor kappaB (NF-kappaB) and tumour necrosis factor alpha (TNF-alpha) production in the THP-1 monocytic cell line. *Br J Haematol* 2000; **110**:130-134.
- [40] Brueckmann M, Hoffmann U, De RL, Weiler HM, Liebe V, Lang S *et al.* Activated protein C inhibits the release of macrophage inflammatory protein-1-alpha from THP-1 cells and from human monocytes. *Cytokine* 2004; **26**:106-113.
- [41] Yuksel M, Okajima K, Uchiba M, Horiuchi S, Okabe H. Activated protein C inhibits lipopolysaccharide-induced tumor necrosis factor-alpha production by inhibiting activation of both nuclear factor-kappa B and activator protein-1 in human monocytes. *Thromb Haemost* 2002; **88**:267-273.

## FIGURE LEGENDS

**Figure 1.**

**Activated protein C (APC) concentrations in bronchoalveolar lavage samples.** Healthy subjects (N = 4 per group) were challenged bilaterally in a lung subsegment via a bronchoscope with lipopolysaccharide (4 ng/kg body weight, A-B) or saline (C). Directly thereafter subjects received recombinant human (rh) APC 15  $\mu$ g (A) or 75  $\mu$ g (B-C) dissolved in 10 mL of saline in one subsegment and 10 mL of saline in the other lung subsegment. Bilateral bronchoalveolar lavage was done 6 hours after challenge. APC levels were undetectable in any of the control lung subsegments not administered with rhAPC. \* $P < 0.05$ .

Figure 1.



**Figure 2.**

**Intrabronchial instillation of recombinant human activated protein C (rhAPC) augments lipopolysaccharide-induced coagulation in the bronchoalveolar space.**

Healthy subjects (N = 4 per group) were challenged bilaterally in a lung subsegment via a bronchoscope with lipopolysaccharide (4 ng/kg body weight, A-B) or saline (C). Directly thereafter subjects received recombinant human (rh) APC 15  $\mu$ g (A) or 75  $\mu$ g (B-C) dissolved in 10 mL of saline in one subsegment and 10 mL of saline in the other lung subsegment. Thrombin-antithrombin complex (TATc) levels were measured in bronchoalveolar lavage samples obtained 6 hours after challenge. \* $P < 0.05$ , \*\* $P < 0.01$ .

Figure 2.




Figure 3.

**Impact of intrabronchial recombinant human activated protein C (rhAPC) on lipopolysaccharide-induced release of plasminogen activator inhibitor type I in the bronchoalveolar space.** Healthy subjects (N = 4 per group) were challenged bilaterally in a lung subsegment via a bronchoscope with lipopolysaccharide (4 ng/kg body weight, A-B) or saline (C). Directly thereafter subjects received recombinant human (rh) APC 15 µg (A) or 75 µg (B-C) dissolved in 10 mL of saline in one subsegment and 10 mL of saline in the other lung subsegment. Plasminogen activator inhibitor type I (PAI-1) levels were measured in bronchoalveolar lavage samples obtained 6 hours after challenge.

Figure 3.




Figure 4.

**Impact of intrabronchial recombinant human activated protein C (rhAPC) on lipopolysaccharide-induced release of tissue-factor pathway inhibitor in the bronchoalveolar space.** Healthy subjects (N = 4 per group) were challenged bilaterally in a lung subsegment via a bronchoscope with lipopolysaccharide (4 ng/kg body weight, A-B) or

saline (C). Directly thereafter subjects received recombinant human (rh)APC 15  $\mu\text{g}$  (A) or 75  $\mu\text{g}$  (B-C) dissolved in 10 mL of saline in one subsegment and 10 mL of saline in the other lung subsegment. Tissue-factor pathway inhibitor (TFPI) levels were measured in bronchoalveolar lavage samples obtained 6 hours after challenge. \* $P < 0.05$ .

Figure 4.




Figure 5.

**Impact of intrabronchial recombinant human activated protein C (rhAPC) on the lipopolysaccharide-induced increase in total leukocyte and neutrophil counts in the bronchoalveolar space.** Healthy subjects (N = 4 per group) were challenged bilaterally in a lung subsegment via a bronchoscope with lipopolysaccharide (4 ng/kg body weight, A-B) or saline (C). Directly thereafter subjects received recombinant human (rh) APC 15  $\mu\text{g}$  (A) or 75  $\mu\text{g}$  (B-C) dissolved in 10 mL of saline in one subsegment and 10 mL of saline in the other lung subsegment. White blood cell counts (WBCs) and neutrophil counts were determined in bronchoalveolar lavage samples obtained 6 hours after challenge.

Figure 5.




Figure 6.

**Intrabronchial instillation of recombinant human activated protein C (rhAPC) augments lipopolysaccharide-induced release of inflammatory mediators in the bronchoalveolar space.** Healthy subjects (N = 4 per group) were challenged bilaterally in a lung subsegment via a bronchoscope with lipopolysaccharide (4 ng/kg body weight). Directly thereafter subjects received recombinant human (rh) APC 75 µg dissolved in 10 mL of saline in one subsegment and 10 mL of saline in the other lung subsegment. Mediator levels were measured in bronchoalveolar lavage samples obtained 6 hours after challenge. \* $P < 0.05$ , \*\* $P < 0.01$ .

Figure 6.

