



# Aclidinium inhibits cholinergic and tobacco smoke-induced MUC5AC in human airways

J. Cortijo<sup>\*,#</sup>, M. Mata<sup>#,†</sup>, J. Milara<sup>#</sup>, E. Donet<sup>+</sup>, A. Gavalda<sup>†</sup>,  
M. Miralpeix<sup>#,†</sup> and E.J. Morcillo<sup>\*,§</sup>

**ABSTRACT:** Mucus hypersecretion and mucin MUC5AC overexpression are pathological features of chronic obstructive pulmonary disease (COPD). This study examines the inhibitory effect of acclidinium, a new long-acting muscarinic antagonist, on MUC5AC expression in human airway epithelial cells.

MUC5AC mRNA (RT-PCR) and protein expression (ELISA and immunohistochemistry) were studied in human bronchial tissue and differentiated human airway epithelial cells activated with carbachol (100 µM) or cigarette smoke extract in the absence or presence of acclidinium.

Carbachol increased MUC5AC mRNA and protein expression in human bronchus and cultured epithelial cells. Aclidinium inhibited the carbachol-induced MUC5AC mRNA and protein expression with potency (half maximal inhibitory concentration) ~1 nM in human bronchus and cultured airway epithelial cells. AG1478, a selective inhibitor of epidermal growth factor receptor (EGFR) tyrosine kinase, inhibited carbachol-induced MUC5AC responses, indicating EGFR transactivation. Aclidinium inhibited carbachol-induced phospho-EGFR and phospho-p44/42 MAPK expression. In cultured airway epithelial cells transfected with small interfering (si)RNA against muscarinic receptor subtypes, siRNA-M3 but not siRNA-M2 blocked carbachol-induced MUC5AC expression. Cigarette smoke-induced MUC5AC upregulation in cultured airway epithelial cells was suppressed by acclidinium.

In conclusion, acclidinium decreases carbachol and tobacco smoke-induced MUC5AC overexpression in human airway epithelial cells. This effect may contribute to the clinical efficacy of acclidinium in mucus hypersecretory diseases including COPD.

**KEYWORDS:** Aclidinium, human airway epithelial cells, human isolated bronchus, mucin MUC5AC, muscarinic receptor subtypes, small interfering RNA

Mucus hypersecretion is an important feature of chronic inflammatory airway diseases such as chronic obstructive pulmonary disease (COPD) and asthma, and contributes to their morbidity and mortality [1]. MUC5AC is the predominant mucin gene expressed in healthy human airway epithelial cells, and its expression is augmented in smokers, COPD patients and asthmatics [2].

COPD and asthma are associated with increased pulmonary vagal activity [3]. Muscarinic antagonists are effective drugs for the treatment of COPD and certain forms of asthma, because they exert an anticholinergic effect that results in relaxation of airway smooth muscle [4]. Furthermore, there is recent awareness of the existence of a non-neuronal cholinergic system in humans. Airway epithelial cells are endowed with this

system, which represents a previously unappreciated regulatory pathway in pulmonary inflammation and remodelling [5]. Dysfunction of the non-neuronal cholinergic system appears to be involved in the pathophysiology of asthma and COPD [6]. Therefore, these potential anti-inflammatory and anti-remodelling effects of the muscarinic antagonists shown in animal models [7] may be of added value to their established bronchodilation in the management of chronic respiratory diseases.

Aclidinium is a novel, long-acting, muscarinic antagonist that has reached phase III clinical development for COPD treatment [8]. In pre-clinical studies, acclidinium demonstrated potent muscarinic antagonist activity, comparable to ipratropium and tiotropium, and long duration of action [9]. The aim of the present study was to

## AFFILIATIONS

<sup>\*</sup>Dept of Pharmacology, Faculty of Medicine, University of Valencia,  
<sup>#</sup>Research Unit, University General Hospital Consortium,  
<sup>†</sup>Centro de Investigación Biomedica en Red de Enfermedades Respiratorias (CIBERES),  
<sup>§</sup>Clinical Pharmacology Unit, Research Foundation and University Clinic Hospital, Valencia, and  
<sup>†</sup>R&D Centre, Almirall, Barcelona, Spain.

## CORRESPONDENCE

E.J. Morcillo  
Clinical Pharmacology  
Faculty of Medicine of the University of Valencia  
Research Foundation of University Clinic Hospital  
Av. Blasco Ibanez 17  
E-46010 Valencia  
Spain  
E-mail: Esteban.Morcillo@uv.es

Received:

Nov 16 2009

Accepted after revision:

May 12 2010

First published online:

June 04 2010

This article has supplementary material accessible from [www.erj.ersjournals.com](http://www.erj.ersjournals.com)

characterise the effect of acridinium on regulating carbachol-induced increase of MUC5AC expression in human isolated bronchus and well-differentiated human primary airway epithelial cells cultured in an air–liquid interface (ALI) system. In addition, since tobacco smoke exposure is associated with the pathogenesis of COPD and steroid resistance in COPD and severe asthma [10], we examined the effects of acridinium on cigarette smoke extract (CSE)-induced increase of MUC5AC expression in human cultured airway epithelial cells. Acridinium was found to effectively decrease carbachol and cigarette smoke-induced MUC5AC overexpression in human airway epithelial cells. Preliminary data from this study have been presented at the American Thoracic Society congress [11].

## METHODS

### *Human bronchial tissue experimental protocol*

Human lung tissue was obtained from patients who had undergone surgery for lung carcinoma, as previously outlined [12]. Experiments were approved by the local ethics committee and informed consent was obtained. Tissue from a total of 12 patients was included in this study. All patients were smokers until the moment of lung carcinoma diagnosis (~1 month before lung surgery). None of the patients included in this study had COPD and none of the patients were chronically treated with theophylline,  $\beta$ -adrenoceptor agonists, corticosteroids or anticholinergic drugs. Clinical details of the different patients are provided in the online supplementary material.

Human bronchial tissues were pretreated with antagonists or their vehicles for 15 min prior to stimulation with carbachol and remained until termination of experiments. Carbachol was selected as a cholinergic agonist as it is widely used in the literature and is resistant to degradation by cholinesterases present in human bronchial epithelium [13]. In this study we used carbachol at 100  $\mu$ M concentration, since we observed that it was near maximal responses, which is in keeping with other studies of goblet cell secretion and cell culture models using cholinomimetics [14]. Therefore, this carbachol concentration was selected for all subsequent experiments.

### *Differentiated human bronchial epithelial cells cultured in ALI*

Human bronchial epithelial cells were cultured and differentiated in Transwell inserts (Corning Costar, High Wycombe, UK) under ALI conditions, as previously described [15]. In brief, a multilayered bronchial epithelium was obtained by seeding cells ( $8.25 \times 10^4$  cells per insert) onto polyester inserts. Cells were submerged in differentiation media (50% DMEM in basal epithelial growth media (BEGM; Clonetics, Wokingham, UK) for the first 7 days. Cells were then cultured for an additional 21 days with the apical surface exposed to air. Based on results from pilot experiments in airway epithelial ALI cultures that studied the time course of the expression of MUC5AC (1, 2, 6, 12 and 24 h), the carbachol response was obtained at 12 h, a time point similar to that studied in conventional culture models [12].

### *Mucin MUC5AC protein expression*

MUC5AC protein was measured by ELISA, as outlined previously [12]. In brief, for MUC5AC ELISA, 100  $\mu$ g total protein extracted from human bronchial tissues was incubated at 40°C until dry. Plates were blocked with 2% bovine serum

albumin for 1 h at room temperature. After three washes, plates were incubated with 50  $\mu$ L of mouse monoclonal antibody (mAb) to MUC5AC (clone 45M1, 1:100; NeoMarkers, Fremont, CA, USA). After incubation with secondary horseradish peroxidase-goat anti-mouse (1:10,000) colour reaction was read at 450 nm. To confirm ELISA results, western blot analysis of MUC5AC was carried out in human bronchial homogenates, as previously reported [12].

For MUC5AC immunohistochemistry of human bronchus, specimens were fixed, cut into sections, stained with haematoxylin–eosin, and with Alcian blue and periodic acid–Schiff (PAS) to visualise goblet cells, and incubated with mouse monoclonal antibody to MUC5AC (clone 45M1, 1:100; NeoMarkers), as previously reported [12].

### *Expression of phospho-EGFR and phospho-p44/42 MAPK*

Protein expression of phospho-EGFR (Tyr845) and phospho-p44/42 MAPK (Thr202/Tyr204) was determined in cultured airway epithelial cells by PathScan® sandwich ELISA kit following the instructions of the manufacturer (Cell Signaling Technology, Beverly, MA, USA). A 15 min incubation time with carbachol was chosen because preliminary experiments indicated that there was little difference in phosphorylation between 5 and 15 min (data not shown). In some experiments, acridinium was added 30 min before the addition of carbachol.

In additional experiments, the expression of phospho-p44/42 MAPK was determined by immunohistochemistry in human bronchial tissue using phospho-p44/42 MAPK rabbit mAb (Thr202/Tyr204, 1:100; catalogue number E7028; Sigma, St Louis, MO, USA) with appropriate rabbit and mouse immunoglobulin G negative controls (Sigma).

### *Real time RT-PCR*

Total RNA was isolated from human bronchial tissue or human epithelial cells differentiated in ALI, as previously outlined [15]. cDNA was amplified with specific primers to MUC5AC, muscarinic receptors (M)1–5 and GAPDH (as endogenous control) using 7900 HT Fast Real-Time PCR System (Applied Biosystems, Foster City, CA, USA). The primers used to amplify cDNA were obtained from TaqMan Gene Expression Assays (Applied Biosystem) to MUC5AC (catalogue number Hs01365616\_m1), M1 (catalogue number Hs00912795\_m1), M2 (catalogue number Hs00265208\_m1), M3 (catalogue number Hs00325478\_m1), M4 (catalogue number Hs00265219\_s1), M5 (catalogue number Hs00255278\_s1) and GAPDH (catalogue number 4352339E). Relative quantification of the different transcripts was determined with the  $2^{-\Delta\Delta Ct}$  method using GAPDH as endogenous control. Results were normalised to respective time control relative expression, as previously outlined [12].

### *Transfection of muscarinic receptor siRNA*

Cultured human bronchial epithelial ALI cells were transfected with a commercial 50 nM small interfering (si)RNA against M2 muscarinic receptor gene (PN 4392421; Ambion, Austin, TX, USA) or with 50 nM siRNA M3 muscarinic receptor gene (PN 4390815; Ambion) or with 50 nM siRNA control (Ambion, Huntingdon, UK), as outlined [16]. Briefly, ALI cultured bronchial epithelial cells were exposed to three successive solutions (4°C) containing: 1) EGTA 10 mM, KCl 120 mM, ATP

5 mM, MgCl<sub>2</sub> 2 mM and TES 20 mM (pH 6.8; 20 min); 2) KCl 120 mM, ATP 5 mM, MgCl<sub>2</sub> 2 mM, TES 20 mM and siRNA 50 nM (pH 6.8; 3 h); and 3) KCl 120 mM, ATP 5 mM, MgCl<sub>2</sub> 10 mM, TES 20 mM and siRNA 50 nM (pH 6.8; 30 min). Subsequently, ALI epithelial cells were bathed in a fourth solution containing NaCl 140 mM, KCl 5 mM, MgCl<sub>2</sub> 10 mM, glucose 5 mM and MOPS 2 mM (pH 7.1, 22°C), in which Ca<sup>2+</sup> concentration was gradually increased from 0.01 to 0.1 to 1.8 mM every 15 min. Then, cells were cultured in differentiation media. The mRNA expression for M2 and M3 transcripts was determined by real time RT-PCR after 72 h post-silencing and compared with siRNA control at the respective time to determine silencing efficiency. Furthermore, the M2 and M3 muscarinic receptor protein expression was measured by western blot after 72 h of silencing using commercial antibodies against muscarinic receptors M2 (catalogue number M9558; Sigma) and M3 (catalogue number M0194; Sigma), as previously described [15]. Protein expression was referred to  $\beta$ -actin (1:1,000; Sigma) expression as internal control. All experiments were performed in triplicate.

#### **Cigarette smoke exposure of differentiated human airway epithelial cells**

Differentiated bronchial epithelial cells were used to examine the MUC5AC responses after exposure to CSE prepared as previously outlined [17]. Aqueous CSE was applied to the apical side of epithelial cells to mimic the *in vivo* pattern of cigarette smoke exposure. Briefly, smoke of research grade cigarettes 2R4F (Tobacco Health Research, University of Kentucky, Lexington, KY, USA) was bubbled into a flask containing 25 mL of pre-warmed (37°C) DMEM medium obtaining a 100% CSE solution that was used within 30 min of CSE preparation. The CSE solutions were assessed by measuring absorbance at 320 nm. Stock solutions with absorbance values of  $3.0 \pm 0.1$  were used. To test for cytotoxicity from CSE exposure, cultured cells were treated with CSE concentrations of up to 30% for 24 h. No significant increase of supernatant lactate dehydrogenase content above control levels were observed (colorimetric assay; Sigma; data not shown).

#### **Statistics**

Data are expressed as mean  $\pm$  SEM of *n* experiments. In concentration–response experiments, the -log inhibitory concentration 50% (IC<sub>50</sub>) was calculated by nonlinear regression to express compound potency (GraphPad Software Inc., San Diego, CA, USA). Statistical analysis was carried out by analysis of variance followed by appropriate *post hoc* tests, including Bonferroni correction. Significance was accepted when  $p < 0.05$ .

#### **Materials**

Drugs used in this study were from Sigma (St Louis, MO, USA), except acridinium bromide, which was synthesised at Amirall (Barcelona, Spain), and tyrphostin-AG1478 (Calbiochem, Nottingham, UK). All products were dissolved in distilled water milli-Q or in dimethyl sulphoxide (DMSO) at the concentration of 10<sup>-3</sup> M, except acridinium, which was dissolved in HCl 1N 1% (v/v) at the concentration of 10<sup>-3</sup> M. Appropriate dilutions were obtained in freshly prepared medium. The DMSO final concentration in the assay solutions was  $\leq 0.1\%$  (v/v); this concentration was without significant effects on MUC5AC expression.

## **RESULTS**

### **Carbachol-induced MUC5AC overexpression in human bronchus is blocked by acridinium**

Carbachol (100  $\mu$ M) augmented the MUC5AC mRNA and protein expression with peak values reached at 1 h and 3 h after carbachol exposure, respectively (fig. 1a). Furthermore, carbachol increased MUC5AC mRNA and protein expression in a dose-dependent manner, reaching a peak value at 100  $\mu$ M (fig. 1b). Thus, we selected this carbachol concentration for the rest of experiments.

The inhibitory potency values (-log IC<sub>50</sub>) of acridinium on carbachol-induced MUC5AC mRNA expression were  $\sim 1$  nM (fig. 1c and d; table 1). Acridinium (100 nM) as well as atropine (1  $\mu$ M) and the selective inhibitor of epidermal growth factor receptor (EGFR) tyrosine kinase, tyrphostin-AG1478 (10  $\mu$ M), abolished the carbachol-induced increase in MUC5AC mRNA and protein (fig. 1e). In contrast, the antagonist of nicotinic receptors hexamethonium (100  $\mu$ M) did not alter the MUC5AC response to carbachol (fig. 1e).

Immunohistochemistry experiments showed that MUC5AC immunoreactivity was localised in Alcian blue and PAS-stained goblet cells. The MUC5AC positive staining in airway epithelium was increased in carbachol-exposed preparations, and this augmentation was reduced in acridinium-treated tissues (fig. 2). Additional immunohistochemical experiments addressed to assert the activation of downstream elements of the EGFR pathway (see below) showed that carbachol augmented the expression of phospho-p44/42 in human bronchial tissue and this effect was also blocked by acridinium (fig. 2).

### **Carbachol-induced MUC5AC overexpression in cultured human ALI airway epithelial cells is blocked by acridinium**


In human bronchial epithelial ALI culture model, carbachol (100  $\mu$ M) increased significantly MUC5AC mRNA and protein after 12 h of incubation (fig. 3a). The carbachol-induced MUC5AC expression was suppressed by acridinium and atropine in a concentration-dependent manner (fig. 3b and c; table 1). Similar findings were obtained with tyrphostin-AG1478 (10  $\mu$ M), but not with hexamethonium (100  $\mu$ M) or with direct stimulation with nicotine (fig. 3d).

Furthermore, we observed that relative protein expression levels of phospho-EGFR and phospho-p44/42 were augmented at 15 min ( $2.02 \pm 0.21$  and  $2.29 \pm 0.17$ -fold increase, respectively;  $n=3$ ;  $p < 0.05$  versus control at 15 min) and declined at 30 and 60 min following carbachol addition (fig. 3e). In this sense, acridinium (100 nM), atropine (1  $\mu$ M) and tyrphostin-AG1478 (10  $\mu$ M) suppressed the carbachol (15 min incubation)-induced EGFR and p42/44 phosphorylation ( $p < 0.05$ ) (fig. 3f).

### **Cigarette smoke-induced MUC5AC expression in cultured human ALI airway epithelial cells is blocked by acridinium**

CSE increased MUC5AC mRNA and protein expression in a time and concentration-dependent manner (fig. 4a and b). Based on these results, CSE (10%, 24 h) was selected for further experiments as producing consistent increases of MUC5AC mRNA and protein similar to those obtained with carbachol.

CSE (10%, 24 h)-induced increase of MUC5AC mRNA and protein expression was suppressed by acridinium and atropine in a dose-dependent fashion (fig. 4c and d; table 1). Furthermore,



**FIGURE 1.** Carbachol (CCh) induces MUC5AC mRNA and protein expression in human isolated bronchus. a) Time course of the relative expression of MUC5AC mRNA (○) and protein (●) in human isolated bronchus. The peak expression for MUC5AC mRNA was observed 1 h after stimulation with CCh (100 μM), thus preceding the peak expression of MUC5AC protein at 3 h. b) CCh dose-dependently increased MUC5AC mRNA (1 h of stimulation; ●) and protein (3 h of stimulation; ○) expression. Concentration-dependent inhibition of the CCh 100 μM-induced overexpression of MUC5AC mRNA (c) and protein (d) obtained in the presence of different concentrations of acridinium (●) and atropine (○) under the experimental conditions described in the methods section. e) Relative quantification of MUC5AC mRNA and protein levels in human bronchus unstimulated (control) or stimulated with CCh (100 μM) in the absence or presence of acridinium (ACL; 100 nM), atropine (ATR; 1 μM), hexamethonium (HEX; 100 μM) or tyrphostin AG1478 (10 μM). Exposure time was 1 h for MUC5AC mRNA determination and 3 h for MUC5AC protein measurements. MUC5AC was determined using real time RT-PCR by the 2<sup>-ΔΔCt</sup> method; points show the increase in expression of MUC5AC relative to GAPDH normalised to respective controls in time as mean ± SEM of the 2<sup>-ΔΔCt</sup> values. MUC5AC protein was determined in bronchial tissue by ELISA; points are mean ± SEM. Data were obtained from three to five different patients. \*: p<0.05 versus basal values; #: p<0.05 versus CCh.

CSE-induced MUC5AC expression was inhibited by acridinium (100 nM) as well as atropine (1 μM) and tyrphostin-AG1478 (10 μM), but remained unaltered in the presence of hexamethonium (100 μM) (fig. 4e).

In other experiments performed in differentiated bronchial epithelial cells, CSE induced EGFR and p42/44 phosphorylation, reaching a peak value at 30 min (fig. 5a). Moreover, the CSE-induced EGFR and p42/44 phosphorylation was partially suppressed by acridinium (100 nM) as well as atropine (1 μM) and tyrphostin-AG1478 (10 μM) (p<0.05) (fig. 5b).

**M3 muscarinic receptor subtype appears involved in MUC5AC response to carbachol and CSE in human cultured airway epithelial cells**

Setting the expression level of M1 mRNA at 1, the amount of M2 mRNA expression in ALI airway epithelial cells was ~5.2 times higher and that of M3 was ~2.4-fold higher than that of M1 (fig. 6a). Additional experiments demonstrated also the

presence of M4 and M5 expression (data not shown). These data were corroborated at protein level, where M2 receptor was the most expressed followed by M3 (fig. 6b).

Furthermore, following CSE exposure, a significant increase in the mRNA and protein expression of M3 was observed without significant changes of the relative levels of M1 and M2 (fig. 6a and b).

Due to the low expression of M1, we studied the role of muscarinic M2 and M3 receptor subtypes in the MUC5AC secretion. To this respect, transfection of human ALI airway epithelial cells with siRNA-M2 or siRNA-M3 decreased the mRNA expression of M2 and M3 by ~62% and ~73%, respectively, with the corresponding protein downregulation (fig. 7a and b). The siRNA-M2 did not affect the carbachol- or CSE-induced MUC5AC mRNA expression, while siRNA-M3 blocked the MUC5AC mRNA response to carbachol and CSE, thus indicating a preferential role of M3 receptors in this response (fig. 7c and d). To further study the implication of

**TABLE 1** Potency values of acclidinium as antagonist of carbachol- and cigarette smoke-induced augmentation of the expression of mucin MUC5AC mRNA and protein expression in human isolated bronchial tissue and human airway epithelial cells cultured in air-liquid interface system


	Potency values as inhibitor of	
	MUC5AC mRNA expression	MUC5AC protein expression
<b>Carbachol in human isolated bronchial tissues</b>		
Acclidinium	9.17 ± 0.14	8.81 ± 0.16
Atropine	8.61 ± 0.08	8.78 ± 0.09
<b>Carbachol in human cultured airway epithelial cells</b>		
Acclidinium	9.13 ± 0.12	8.95 ± 0.09
Atropine	8.69 ± 0.05	8.60 ± 0.06
<b>Cigarette smoke in human cultured airway epithelial cells</b>		
Acclidinium	8.80 ± 0.12	8.67 ± 0.11
Atropine	8.29 ± 0.14	8.24 ± 0.13

Data are presented as mean ± SEM of 3–5 independent experiments. The corresponding values for atropine are also given for comparison. Potency values are given as the  $-\log IC_{50}$  values calculated from the concentration-dependent inhibition of the carbachol 100  $\mu$ M and cigarette smoke 10%-induced overexpression of MUC5AC mRNA and protein obtained in the presence of acclidinium (0.1, 1, 10 and 100 nM) and atropine (1, 10, 100 and 1000 nM) under the experimental conditions described in the methods section.


M1, M2 and M3 receptors in MUC5AC upregulation, we pre-incubated differentiated human ALI epithelial cells with pirenzepine 1  $\mu$ M (M1 antagonist), methoctramine 1  $\mu$ M (M2 antagonist) and p-fluoro-hexahydro siladifenidol (pFHHSid) 1  $\mu$ M (M3 antagonist). We detected that both pirenzepine and methoctramine did not modify significantly the effect of carbachol and CSE on MUC5AC expression. In contrast, pFHHSid effectively attenuated carbachol- and CSE-induced MUC5AC overexpression, confirming data from siRNA experiments (fig. 7e and f).

## DISCUSSION


In this study, we show that acclidinium potently inhibited the carbachol-induced MUC5AC overexpression in human bronchial tissue *in vitro* as well as the carbachol- and cigarette smoke-induced augmentation of MUC5AC expression in differentiated human airway epithelial cells cultured in ALI. This is the first report showing a direct effect of a cholinergic agonist in upregulating MUC5AC expression in human airway epithelial cells by activation of muscarinic receptors and transactivation of EGFR. Also, we showed that cigarette



**FIGURE 2.** Photomicrographs of representative histological sections from human bronchial tissue unstimulated (control) or stimulated with carbachol (CCh; 100  $\mu$ M) for 3 h or 30 min (in case of phospho-p44/42 MAPK immunostaining) in the absence or presence of acclidinium (ACL; 100 nM). Sections show staining with haematoxylin–eosin (H&E), Alcian blue, periodic acid–Schiff (PAS) or immunohistochemical staining of MUC5AC or phospho-p44/42 MAPK. Mucin stores in goblet cells appear as blue staining for alcian blue-positive cells and purple staining for PAS-positive cells. MUC5AC immunoreactivity was observed as brown staining in goblet cells. Ciliated cells showed no staining for MUC5AC. The sections demonstrate increased Alcian blue, PAS, MUC5AC and phospho-p44/42 staining in the tissues exposed to CCh which were prevented by the incubation with ACL for 15 min. Negative control is expressed as nonspecific immunoglobulin (Ig)G. Scale bars=25  $\mu$ m.



**FIGURE 3.** Carbachol (CCh) induces MUC5AC mRNA and protein expression in differentiated human bronchial epithelial cells. a) Time course of the relative expression of MUC5AC mRNA (○) and protein (●) in differentiated human bronchial epithelial cells in air-liquid interface system. The peak expression for MUC5AC mRNA and protein was observed 12 h after stimulation with CCh (100 μM). Concentration-dependent inhibition of the CCh 100 μM-induced overexpression of MUC5AC mRNA (b) and protein (c) obtained in the presence of different concentrations of acclidinium (ACL; ●) and atropine (ATR; ○) under the experimental conditions described in the methods section. d) Relative quantification of MUC5AC mRNA (□) and protein (■) in human bronchial epithelial cells cultured in air-liquid interface system, unstimulated (control) or stimulated with CCh (100 μM) in the absence and presence of ACL (100 nM), ATR (1 μM), hexamethonium (HEX; 100 μM) or AG1478 (10 μM). Nicotine (NIC; 10 μM) was without effect. Exposure time was 12 h for MUC5AC mRNA and protein determination. Columns show the relative increase from control levels at 12 h as mean ± SEM of four independent experiments. \*: p<0.05 from control; #: p<0.05 versus CCh. e) Relative time course of the epidermal growth factor receptor (EGFR) and p44/42 phosphorylation in differentiated human bronchial epithelial cells in air-liquid interface system following CCh 100 μM stimulation. Levels of EGFR (○) and p44/42 (●) phosphorylation are normalised to basal conditions. f) Relative expression of the EGFR and p44/42 phosphorylation in human bronchial epithelial cells cultured in air-liquid interface system, unstimulated (control) or stimulated with CCh 100 μM in the absence and presence of ACL 100 nM, ATR 1 μM, HEX 100 μM or AG1478 10 μM. CCh exposure time was 15 min. Columns show the relative increase from control levels at 15 min as mean ± SEM of four independent experiments. \*: p<0.05 from control; #: p<0.05 versus CCh.



**FIGURE 4.** Cigarette smoke extract (CSE) induces MUC5AC mRNA and protein expression in differentiated human bronchial epithelial cells. a) Time course of the relative expression of MUC5AC mRNA (●) and protein (○) in differentiated human bronchial epithelial cells in air–liquid interface system. The peak expression for MUC5AC mRNA and protein expression was observed 24 h after stimulation with CSE 10%. b) CSE dose-dependently increased MUC5AC mRNA (●) and protein (○) expression. Concentration-dependent inhibition of the CSE 10%-induced overexpression of MUC5AC mRNA (c) and protein (d) obtained in the presence of different concentrations of acclidinium (ACL; ●) and atropine (ATR; ○) under the experimental conditions described in the methods section. e) Relative quantification of MUC5AC mRNA (□) and protein (■) in human bronchial epithelial cells cultured in air–liquid interface system, unstimulated (control) or stimulated with CSE 10% in the absence and presence of ACL (100 nM), ATR (1 μM), hexamethonium (HEX; 100 μM) or AG1478 (10 μM). Exposure time was 24 h for MUC5AC mRNA and protein determination. Columns show the relative increase from control levels at 24 h as mean ± SEM of four independent experiments. \*: p < 0.05 from control; #: p < 0.05 versus CSE.


smoke-induced increase of MUC5AC expression *in vitro* can be effectively suppressed by the use of muscarinic antagonists. Since mucus hypersecretion is considered pathologically relevant in COPD and asthma, this inhibitory effect of acclidinium is of potentially added therapeutic value.

It is well established that goblet cell hypertrophy and hyperplasia occur in the large airways of habitual cigarette smokers with or without airway obstruction and result in epithelial mucin stores that are significantly higher than normal [18]. Goblet cells are located on the bronchial surface epithelium and MUC5AC is the best characterised mucin in this cell type, and its expression is directly correlated with airway obstruction [18–20]. Therefore, MUC5AC was selected for this study. While MUC5AC is increased directly by cigarette smoke in goblet cells [20], the role of other mucins such as MUC5B seems contradictory since MUC5B is not altered or, indeed, diminished in goblet cells from smokers and COPD patients [18, 19]. However, the major place of mucin production is located in submucosal glands in the central airways, where both MUC5AC and MUC5B are abundant in smokers and COPD patients [19, 21]. In this line, MUC5B is

mainly produced in submucosal gland cells and is the most abundant in COPD sputum, while MUC5AC is mainly produced in goblets cells and is the most abundant in smoker sputum without airway obstruction [21]. However, it seems that only MUC5AC may be induced by different stimuli, while MUC5B appears more constitutively expressed and less sensitive to the inducible effect of irritants such as cigarette smoke [19]. This work is focused on MUC5AC production in goblet cells from the surface epithelium due to the inducible characteristic of MUC5AC and its correlation with human airway obstruction found in COPD. However, since the major mucins are released predominantly from submucosal gland cells, this may be considered a limitation of the present study.

The human bronchial tissue *in vitro* is a preparation that has previously been shown to have a basal secretion of mucin MUC5AC produced principally by goblet cells [12].

The rapid time course of MUC5AC secretion found in this work (fig. 1) was in accord with previous reports for epidermal growth factor [12] and for carbachol in goblet cells from rat conjunctiva [14].




**FIGURE 5.** Cigarette smoke extract (CSE)-induced epidermal growth factor receptor (EGFR) and p44/42 phosphorylation is partially suppressed by antimuscarinics. a) Relative time course of the EGFR and p44/42 phosphorylation in differentiated human bronchial epithelial cells in air-liquid interface system following CSE 10% stimulation. Levels of EGFR (○) and p44/42 (●) phosphorylation are normalised to basal conditions. b) Relative expression of the EGFR (■) and p44/42 (□) phosphorylation in human bronchial epithelial cells cultured in air-liquid interface system, unstimulated (control) or stimulated with CSE 10% in the absence and presence of acclidinium (ACL; 100 nM), atropine (ATR; 1 μM) or AG1478 (10 μM). CSE exposure time was 30 min. Columns show the relative increase from control levels at 30 min as mean ± SEM of four independent experiments. \*: p<0.05 from control; #: p<0.05 versus CSE.

In this *in vitro* model, acclidinium inhibited the augmented expression of MUC5AC induced by carbachol in a concentration-dependent fashion, with potency values of ~1 nM, which is in the range of the potency of this muscarinic antagonist against cholinergic contraction in isolated trachea [22].

Immunohistochemical analysis of human bronchial tissues confirmed that carbachol exposure resulted in an augmented expression of MUC5AC-positive stained cells in the airway epithelium, which was paralleled by an increased presence of Alcian blue-positive and PAS-stained goblet cells. Treatment with acclidinium effectively prevented this carbachol-induced overexpression of MUC5AC. In this point, it is interesting to note that the histochemical analysis of goblet cells was not always correlated with the immunohistochemical staining of MUC5AC, as previously reported [19, 20], which may be taken in account in the evaluation of this study.


EGFR and its tyrosine receptor kinase-based pathway mediate mucin production by airway epithelial cells in response to a variety of pathogenic and environmental insults [23]. The transactivation of EGFR following the stimulation of muscarinic receptors has been described in different cell types, including conjunctival goblet cells in relation to mucin secretion [24]. We confirmed and extended this finding by showing that carbachol-induced augmentation of MUC5AC mRNA and protein expression was mediated by the transactivation of EGFR and the subsequent phosphorylation of p44/42 MAPK. Thus, blockade of muscarinic receptors by acclidinium avoided transactivation of EGFR.

Because the human bronchial tissue contains a variety of different cell types, we also aimed to study the direct impact of carbachol on well-differentiated human primary airway epithelial cells with relation to MUC5AC expression. We found that carbachol-induced MUC5AC overexpression was abolished



**FIGURE 6.** Relative expression of muscarinic receptor subtypes M1, M2 and M3 in human bronchial epithelial cells cultured as air-liquid interface in the absence (control; □) and presence of exposure to cigarette smoke extract (CSE; 10%, 24 h; ■). M receptor levels were measured by quantitative real-time PCR (a) and western blot (b). The level of the mRNA expression of M2 and M3 receptors in the controls is referred to the expression level of M1 receptors taken as unity. Columns are mean ± SEM of five independent experiments. \*: p<0.05 from M1; #: p<0.05 from the corresponding control at 24 h.





**FIGURE 7.** Effects of M2- and M3-targeted small interfering (si)RNA on the expression of their corresponding receptors (a) and on the carbachol (CCh)- (b) and cigarette smoke extract (CSE)-induced (c) MUC5AC expression in differentiated human bronchial epithelial cells in air-liquid interface system. siRNA-M2 and siRNA-M3 resulted in a significant reduction in their respective muscarinic M2 and M3 mRNA and protein expression (a). Muscarinic receptor mRNA transcripts were measured by quantitative RT-PCR and protein expression by western blotting. The MUC5AC mRNA overexpression elicited by CCh (100  $\mu$ M) or CSE 10% was unaltered in the presence of siRNA-M2 but blocked by siRNA-M3 (b, c). Columns are mean  $\pm$  SEM of three independent experiments per condition. \*:  $p < 0.05$  from corresponding controls; #:  $p < 0.05$  from the response to CCh or CSE without siRNA-M3. In other experiments, differentiated human bronchial epithelial cells were incubated in presence or absence of pirenzepine (M1 antagonist at 1  $\mu$ M), methoctramine (M2 antagonist at 1  $\mu$ M) or pFHHSid (M3 antagonist at 1  $\mu$ M) for 30 min before the stimulation with d) CCh (100  $\mu$ M, 12 h) or e) CSE (10%, 24 h). MUC5AC mRNA was then quantified by real-time PCR. Columns are mean  $\pm$  SEM of three independent experiments. \*:  $p < 0.05$  from basal conditions; #:  $p < 0.05$  from CCh or CSE.

by acridinium and atropine, in a manner similar to that found in a tissue bronchium *in vitro* model, while hexamethonium did not reduce carbachol responses. Furthermore, nicotine (10  $\mu$ M) did not evoke a MUC5AC response, which rules out nicotinic receptor activation in this mechanism. Consistent with the results observed in bronchial tissue, MUC5AC overexpression was a consequence of the transactivation of the EGFR and the subsequent triggering of the phosphorylated p44/42 MAPK signalling cascade, as previously suggested for goblet cells from rat conjunctiva [24].

Next we investigated the role of muscarinic receptor subtypes on carbachol-induced MUC5AC overexpression. The presence of the five subtypes of muscarinic receptors was detected but

we focused our study on M1, M2 and M3 receptors, since only these subtypes exert well known physiological effects in the lungs [25]. We found a predominance of M2 and M3 muscarinic subtypes, with low amounts of M1 receptors. A similar pattern of expression was described for cultured respiratory epithelial cells [5, 26]. In this sense, we found that siRNA specific for siRNA-M3 and the M3 antagonist pFHHSid attenuated the MUC5AC response to carbachol and CSE, while siRNA-M2 and the M2 antagonist methoctramine were without significant effects. These results suggest that the activation of muscarinic M3 receptor appears involved in the mucin MUC5AC response after cholinergic activation in differentiated human airway epithelium. The low expression of M1 receptors impeded the use of specific siRNA in the present

study. However the use of the M1 antagonist pirenzepine was without effect, which discounts the M1 receptor in this process.

Cigarette smoke is widely used in *in vitro* studies due to its relevance in the pathogenesis of COPD [10]. The extent of the increases of MUC5AC expression obtained in the present study after cigarette smoke exposure, and the time point selected for this expression, are in keeping with findings from other studies [27]. In this work we observed that acclidinium with potency close to 1 nM, as well as atropine, inhibited the CSE-induced MUC5AC. In contrast, nicotinic receptors did not affect the MUC5AC upregulation. In this mechanism, the transactivation of EGFR after cigarette smoke exposure was also demonstrated by direct phosphorylation after CSE exposure and by the inhibition of MUC5AC expression obtained in the presence of tyrphostin-AG1478, a selective inhibitor of the EGFR tyrosine kinase. This result is consistent with the overproduction of mucin *via* EGFR found in response to cigarette smoke in the airway epithelial cell line NCI-H292 [28].

The activation of muscarinic receptors in response to cigarette smoke requires explanation. The epithelium of the airways possesses a non-neuronal cholinergic system and local release of acetylcholine may serve a variety of autocrine and paracrine functions [5]. Bronchial epithelial cells in culture express cholinesterases that inactivate acetylcholine [28]. Cigarette smoke is known to impose an important oxidative burden on exposed cells [10] that may degrade acetylcholinesterase and butyrylcholinesterase molecules, while cholineacetyltransferase is not affected, thus resulting in tissue accumulation of acetylcholine [29]. Conversely, we found an upregulation of muscarinic M3 receptors in airway epithelial cells exposed to CSE. This finding would be consistent with a recent study showing that human lung fibroblasts exposed to CSE dramatically increased cholineacetyltransferase and M3 muscarinic receptor expression [30], which may explain the effect of anti-muscarinics on tobacco smoke responses.

The results of this study indicate that acclidinium effectively decreased the MUC5AC overexpression elicited by cholinergic activation and cigarette smoke exposure in human bronchial epithelial cells. This direct inhibitory effect may have an additive effect in the clinical efficacy of acclidinium in mucus hypersecretory diseases, such as COPD.

#### SUPPORT STATEMENT

This work was supported by Almirall S.A., Barcelona, Spain and in part by grants SAF2008-03113 (J. Cortijo), SAF2009-08913 (E.J. Morcillo), CIBERES (CB06/06/0027) and CAIBER (CAI08/01/0039) from the Ministry of Science and Innovation and Health Institute "Carlos III" of the Spanish Government, and research grants from Regional Government (Prometeo/2008/045; "Generalitat Valenciana"). Support from CENIT programme (Spanish Government) and Almirall (Barcelona, Spain) was obtained. J. Milara has a research contract from "Fondo de Investigaciones Sanitarias" of Health Institute "Carlos III" of the Ministry of Health (Spain).

#### STATEMENT OF INTEREST

Statements of interest for J. Cortijo, M. Mata, A. Gavaldà, M. Miralpeix and E.J. Morcillo, and for the study itself, can be found at [www.erj.ersjournals.com/site/misc/statements.xhtml](http://www.erj.ersjournals.com/site/misc/statements.xhtml)

#### ACKNOWLEDGEMENTS

We are grateful for the valuable help of M. Cerdá of the Dept of Pathology, Valencia University Clinic Hospital, Valencia, Spain with the histology and immunohistochemistry experiments. The technical assistance of P. Santamaria of the Dept of Pharmacology, Faculty of Medicine, University of Valencia, Valencia, Spain is also gratefully acknowledged.

#### REFERENCES

- 1 Voynow JA, Rubin BK. Mucins, mucus, and sputum. *Chest* 2009; 135: 505–512.
- 2 Morcillo EJ, Cortijo J. Mucus and MUC in asthma. *Curr Opin Pulm Med* 2006; 12: 1–6.
- 3 Udem BJ, Kollarik M. The role of vagal afferent nerves in chronic obstructive pulmonary disease. *Proc Am Thorac Soc* 2005; 2: 355–360.
- 4 Gross NJ. Anticholinergic agents in asthma and COPD. *Eur J Pharmacol* 2006; 533: 36–39.
- 5 Gwilt CR, Donnelly LE, Rogers DF. The non-neuronal cholinergic system in the airways: an unappreciated regulatory role in pulmonary inflammation? *Pharmacol Ther* 2007; 115: 208–222.
- 6 Gosens R, Zaagsma J, Meurs H, *et al.* Muscarinic receptor signaling in the pathophysiology of asthma and COPD. *Respir Res* 2006; 7: 73.
- 7 Bos IS, Gosens R, Zuidhof AB, *et al.* Inhibition of allergen-induced airway remodelling by tiotropium and budesonide: a comparison. *Eur Respir J* 2007; 30: 653–661.
- 8 Cazzola M, Matera MG. Novel long-acting bronchodilators for COPD and asthma. *Br J Pharmacol* 2008; 155: 291–299.
- 9 Gavaldà A, Miralpeix M, Ramos I, *et al.* Characterization of acclidinium bromide, a novel inhaled muscarinic antagonist, with long duration of action and a favorable pharmacological profile. *J Pharmacol Exp Ther* 2009; 331: 740–751.
- 10 Adenuga D, Yao H, March TH, *et al.* Histone deacetylase 2 is phosphorylated, ubiquitinated, and degraded by cigarette smoke. *Am J Respir Cell Mol Biol* 2009; 40: 464–473.
- 11 Mata M, Milara J, Bedrina B, *et al.* Acclidinium, a new long-acting antimuscarinic, inhibits cholinergic and cigarette smoke-induced up-regulation of mucin MUC5AC expression in human airway epithelial cells. *Am J Respir Crit Care Med* 2009; 179: A6179.
- 12 Mata M, Sarria B, Buenestado A, *et al.* Phosphodiesterase 4 inhibition decreases MUC5AC expression induced by epidermal growth factor in human airway epithelial cells. *Thorax* 2005; 60: 144–152.
- 13 Norel X, Angrisani M, Labat C, *et al.* Degradation of acetylcholine in human airways: role of butyrylcholinesterase. *Br J Pharmacol* 1993; 108: 914–919.
- 14 Rios JD, Zoukhri D, Rawe IM, *et al.* Immunolocalization of muscarinic and VIP receptor subtypes and their role in stimulating goblet cell secretion. *Invest Ophthalmol Vis Sci* 1999; 40: 1102–1111.
- 15 Cortijo J, Milara J, Mata M, *et al.* Nickel induces intracellular calcium mobilization and pathophysiological responses in human cultured airway epithelial cells. *Chem Biol Interact* 2010; 183: 25–33.
- 16 Corteling RL, Brett SE, Yin H, *et al.* The functional consequence of RhoA knockdown by RNA interference in rat cerebral arteries. *Am J Physiol Heart Circ Physiol* 2007; 293: H440–H447.
- 17 Wyatt TA, Schmidt SC, Rennard SI, *et al.* Acetaldehyde-stimulated PKC activity in airway epithelial cells treated with smoke extract from normal and smokeless cigarettes. *Proc Soc Exp Biol Med* 2000; 225: 91–97.
- 18 Innes AL, Woodruff PG, Ferrando RE, *et al.* Epithelial mucin stores are increased in the large airways of smokers with airflow obstruction. *Chest* 2006; 130: 1102–1108.

- 19 Caramori G, Casolari P, Di Gregorio C, *et al.* MUC5AC expression is increased in bronchial submucosal glands of stable COPD patients. *Histopathology* 2009; 55: 321–331.
- 20 O'Donnell RA, Richter A, Ward J, *et al.* Expression of ErbB receptors and mucins in the airways of long term current smokers. *Thorax* 2004; 59: 1032–1040.
- 21 Kirkham S, Kolsum U, Rousseau K, *et al.* MUC5B is the major mucin in the gel phase of sputum in chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2008; 178: 1033–1039.
- 22 Miralpeix M, Gavalda A, Morcillo E, *et al.* Assessment of the potency and duration of action of aclidinium bromide in guinea pig isolated trachea *in vitro*. *Eur Respir J* 2007; 30: 359s.
- 23 Abdullah LH, Davis CW. Regulation of airway goblet cell mucin secretion by tyrosine phosphorylation signaling pathways. *Am J Physiol Lung Cell Mol Physiol* 2007; 293: L591–L599.
- 24 Kanno H, Horikawa Y, Hodges RR, *et al.* Cholinergic agonists transactivate EGFR and stimulate MAPK to induce goblet cell secretion. *Am J Physiol Cell Physiol* 2003; 284: C988–998.
- 25 Racke K, Matthiesen S. The airway cholinergic system: physiology and pharmacology. *Pulm Pharmacol Ther* 2004; 17: 181–198.
- 26 Buhling F, Lieder N, Kuhlmann UC, *et al.* Tiotropium suppresses acetylcholine-induced release of chemotactic mediators *in vitro*. *Respir Med* 2007; 101: 2386–2394.
- 27 Glader P, Moller S, Lilja J, *et al.* Cigarette smoke extract modulates respiratory defence mechanisms through effects on T-cells and airway epithelial cells. *Respir Med* 2006; 100: 818–827.
- 28 Shao MX, Nakanaga T, Nadel JA. Cigarette smoke induces MUC5AC mucin overproduction *via* tumor necrosis factor- $\alpha$ -converting enzyme in human airway epithelial (NCI-H292) cells. *Am J Physiol Lung Cell Mol Physiol* 2004; 287: L420–L427.
- 29 Schallreuter KU, Elwary S. Hydrogen peroxide regulates the cholinergic signal in a concentration dependent manner. *Life Sci* 2007; 80: 2221–2226.
- 30 Profita M, Bonanno A, Siena L, *et al.* Smoke, choline acetyltransferase, muscarinic receptors, and fibroblast proliferation in chronic obstructive pulmonary disease. *J Pharmacol Exp Ther* 2009; 329: 753–763.