- 8 Fleetham J, West P, Mezon B, *et al.* Sleep, arousals, and oxygen desaturation in chronic obstructive pulmonary disease. The effect of oxygen therapy. *Am Rev Respir Dis* 1982; 126: 429–433.
- 9 Murciano D, Armengaud MH, Cramer PH, *et al.* Acute effects of zolpidem, triazolam and flunitrazepam on arterial blood gases and control of breathing in severe COPD. *Eur Respir J* 1993; 6: 625–629.

Eur Respir J 2015; 45: 565–566 | DOI: 10.1183/09031936.00160514 | Copyright ©ERS 2015

From the authors:

We thank R. Katyal for his comments on our study as he raises some interesting points. Although there are some other physiology studies showing no association between benzodiazepine receipt and negative respiratory effects among individuals with chronic obstructive pulmonary disease (COPD) [1, 2], physiology studies are limited in their ability to provide information on the true clinical impact of benzodiazepines in the COPD population. This is because they generally include small numbers of participants, administration of only a single benzodiazepine dose, short follow-up and do not examine for clinically meaningful respiratory outcomes. In contrast, our study [3] included a large population-based sample, real-world benzodiazepine doses and durations, a longer follow-up period, and important and relevant clinical respiratory outcomes.

R. Katyal writes that it would have been helpful to see an analysis of benzodiazepine-related adverse respiratory effects by COPD severity “according to Global Initiative for Chronic Obstructive Lung Disease (GOLD) guidelines”, probably referring to the forced expiratory volume in 1 s (FEV₁) staging system. The health administrative data on which our analysis was based did not contain patient-level clinical information, such as lung function measurements. Newer GOLD guidelines have moved away from classifying COPD severity by FEV₁ level alone, and instead use a combined assessment that incorporates COPD exacerbation history [4]. We explain in our supplementary material that COPD exacerbation history is an important marker of disease severity [3], being associated both with severity of underlying airflow obstruction [5] and mortality [6], and that it is also the single best predictor of future COPD exacerbation, regardless of lung function [7]. We performed a sensitivity analysis reporting outcomes stratifying by COPD exacerbation history in the year prior to index and consistently showed an increased risk of several negative respiratory outcomes among benzodiazepine users across subgroups of differing COPD exacerbation history. This analysis supports the claim that benzodiazepine use is associated with adverse respiratory outcomes regardless of COPD severity.

R. Katyal states that receipt of lower doses of benzodiazepines among individuals with COPD are not associated with increased hospital admission risk and cite a study by KLINK *et al.* [8] to support this statement. However, the study by KLINK *et al.* [8] does not report data on benzodiazepine drug receipt or hospital admissions. A paper by EKSTRÖM *et al.* [9] shows no increased risk of all-cause hospitalisation among prevalent benzodiazepine users with oxygen-dependent COPD, regardless of drug dose. However, in the study by EKSTRÖM *et al.* [9], respiratory-specific hospitalisations were not examined and prevalent (and not incident) benzodiazepine receipt was considered. In addition, the authors’ approach to quantify drug dose is questionable, as individuals are often given flexibility by their doctors regarding the frequency with which they use benzodiazepines, making accurate benzodiazepine dose quantification challenging.

As mentioned by R. Katyal, sleep symptoms are present with increased frequency among individuals with COPD with more respiratory symptoms [8], thereby implying that our finding of increased adverse respiratory outcomes among older adult benzodiazepine users with COPD may be confounded by the presence of sleep symptomatology. However, our propensity-score matched analysis included doctor-diagnosed sleep disorders and new and non-benzodiazepine users were well-matched on this health characteristic, making confounding as a result of this factor unlikely (table 1 in [3]).

We agree that benzodiazepine half-life may impact risk of negative respiratory outcomes and R. Katyal references a physiology study showing worse adverse respiratory effects following administration of a longer-acting benzodiazepine (flunitrazepam) *versus* a short-acting agent (triazolam) [2]. In our supplementary material [3], we analysed adverse respiratory outcomes stratifying by the half-life of the benzodiazepine received. We reported that both long-acting and intermediate/short-acting benzodiazepines were significantly associated with important adverse respiratory outcomes, with slightly higher relative risks among users in the intermediate/short-acting subgroup compared to the long-acting subgroup. Others have similarly reported that benzodiazepine-related psychomotor effects are not related to medication half-life [10].

@ERSpublications

Benzodiazepines, regardless of half-life, are associated with adverse respiratory outcomes across COPD severity <http://ow.ly/DgoFp>

Nicholas T. Vozoris^{1,2} and Anne L. Stephenson^{1,2,3}

¹Division of Respiriology, Dept of Medicine, St Michael's Hospital, Toronto, ON, Canada. ²Dept of Medicine, University of Toronto, Toronto, ON, Canada. ³Keenan Research Centre in the Li Ka Shing Knowledge Institute, St Michael's Hospital, Toronto, ON, Canada.

Correspondence: Nicholas T. Vozoris, Division of Respiriology, Dept of Medicine, St Michael's Hospital, 30 Bond Street, Toronto, Ontario M5B 1W8, Canada. E-mail: nick.vozoris@utoronto.ca

Received: Sept 21 2014 | Accepted: Sept 22 2014

Conflict of interest: None declared.

References

- 1 Midgren B, Hansson L, Skeidsvoll H, *et al*. The effects of nitrazepam and flunitrazepam on oxygen desaturation during sleep in patients with stable hypoxemic nonhypercapnic COPD. *Chest* 1989; 95: 765–768.
- 2 Murciano D, Armengaud MH, Cramer PH, *et al*. Acute effects of zolpidem, triazolam and flunitrazepam on arterial blood gases and control of breathing in severe COPD. *Eur Respir J* 1993; 6: 625–629.
- 3 Vozoris NT, Fischer HD, Wang X, *et al*. Benzodiazepine drug use and adverse respiratory outcomes among older adults with COPD. *Eur Respir J* 2014; 44: 332–340.
- 4 Global Initiative for Chronic Obstructive Lung Disease. Global Strategy for the Diagnosis, Management, and Prevention of Chronic Obstructive Pulmonary Disease. www.goldcopd.org/uploads/users/files/GOLD_Report_2014_Jun11.pdf Date last updated: 2014. Date last accessed: September 2014.
- 5 Donaldson GC, Seemangal TA, Bhowmik A, *et al*. Relationship between exacerbation frequency and lung function decline in chronic obstructive pulmonary disease. *Thorax* 2002; 57: 847–852.
- 6 Connors AF Jr, Dawson NV, Thomas C, *et al*. Outcomes following acute exacerbation of severe chronic obstructive lung disease. The SUPPORT investigators (Study to Understand Prognoses and Preferences for Outcomes and Risks of Treatments). *Am J Respir Crit Care Med* 1996; 154: 959–967.
- 7 Hurst JR, Vestbo J, Anzueto A, *et al*. Susceptibility to exacerbation in chronic obstructive pulmonary disease. *N Engl J Med* 2010; 363: 1128–1138.
- 8 Klink ME, Dodge R, Quan SF. The relation of sleep complaints to respiratory symptoms in a general population. *Chest* 1994; 105: 151–154.
- 9 Ekström MP, Bornefalk-Hermansson A, Abernethy AP, *et al*. Safety of benzodiazepines and opioids in very severe respiratory disease: national prospective study. *BMJ* 2014; 348: g445.
- 10 Herings RM, Stricker BH, de Boer A, *et al*. Benzodiazepines and the risk of falling leading to femur fractures. Dosage more important than elimination half-life. *Arch Intern Med* 1995; 155: 1801–1807.

Eur Respir J 2015; 45: 566–567 | DOI: 10.1183/09031936.00172714 | Copyright ©ERS 2015

HIV and multidrug-resistant tuberculosis: overlapping risk factors

To the Editor:

Recent meta-analyses have indicated that, on average, new HIV-positive tuberculosis (TB) patients are at increased risk of multidrug-resistant (MDR)-TB compared with HIV-negative patients, while this risk is less clear for previously treated TB patients [1, 2]. In a recent issue of the *European Respiratory Journal*, DEAN *et al*. [3] reported an association between HIV infection and MDR-TB disease based on aggregated data reported annually to the World Health Organization. Out of 24 countries where $\geq 75\%$ of TB patients had a HIV test result and at least one HIV-positive MDR-TB case was reported, 11 showed a significant positive association between HIV and multidrug resistance among TB patients, mostly in countries with a high prevalence of multidrug resistance. One of these countries was Kazakhstan, which added data for 40 975 out of the 104 781 patients included in their report.

The authors pointed out that there are likely to be risk factors common to HIV-positive and MDR-TB patients that could not be explored in their analysis. They refer to a study from Moldova, where detailed analysis of 2007–2010 surveillance data showed that the positive association observed among new patients remained after adjustment for potential confounders [4]. We would like to refer to a similar analysis