- 10 Milgrom H, Fowler-Taylor A, Vidaurre CF, *et al.* Safety and tolerability of omalizumab in children with allergic (IgE-mediated) asthma. *Curr Med Res Opin* 2011; 27: 163–169.
- 11 Chung KF, Wenzel SE, Brozek JL, *et al.* International ERS/ATS guidelines on definition, evaluation and treatment of severe asthma. *Eur Respir J* 2014; 43: 343–373.
- 12 Global Initiative for Asthma. Global Strategy for Asthma Management and Prevention. 2014. www.ginasthma.org
- 13 Lowe PJ, Renard D. Omalizumab decreases IgE production in patients with allergic (IgE-mediated) asthma; PKPD analysis of a biomarker, total IgE. *Br J Clin Pharmacol* 2011; 72: 306–320.
- 14 Molimard M, Mala L, Bourdeix I, *et al.* Observational study in severe asthmatic patients after discontinuation of omalizumab for good asthma control. *Respir Med* 2014; 108: 571–576.
- 15 Busse WW, Trzaskoma B, Omachi TA, *et al.* Evaluating Xolair persistency of response after long-term therapy (XPORT). *Am J Respir Crit Care Med* 2014; 189: A6576.

Eur Respir J 2015; 46: 856–859 | DOI: 10.1183/09031936.00008115 | Copyright ©ERS 2015

Bronchial thermoplasty in asthma: 2-year follow-up using optical coherence tomography

To the Editor:

Bronchial thermoplasty (BT) is a novel, nonpharmacological procedure for treatment of severe asthma. Recently, the Asthma Intervention Research 2 clinical trial demonstrated asthmatics had fewer hospitalisations following BT, which persisted 5 years after therapy [1]. However, it is well recognised that asthma is a heterogeneous disease with distinct asthma phenotypes and, not surprisingly, not all asthmatics in that trial benefited from BT [2].

Although bronchoscopic biopsies, pulmonary function tests, exhaled nitric oxide, sputum eosinophil counts and other biological measures have been proposed as biomarkers for evaluating treatment effects [3], these biomarkers cannot provide regional information to characterise airway remodelling in the targeted airways prior to and longitudinally following treatment. Although imaging approaches, such as computed tomography of the lung [4] and magnetic resonance imaging using inhaled contrast agents [5], do provide regional information, these tests are limited to indirect assessment of the small airways. Optical coherence tomography (OCT) is a minimally invasive imaging technique for visualising airway wall structures with near-histological resolution [6–8]. OCT has been used for the evaluation of airway remodelling [9, 10] and early lung neoplastic changes [8, 11].

Identifying asthma phenotypes with the greatest response to BT is likely to bring the potential for better patient selection and ultimately better patient outcomes, and further research into methods capable of careful patient selection for BT has been strongly recommended [12]. Here, our objective was to provide a pilot study in two asthma patients who underwent BT in order to investigate the role of OCT imaging for evaluating airway remodelling prior to and longitudinally following BT treatment.

Two patients with chronic persistent asthma provided written informed consent. Flexible bronchoscopy was performed under local anaesthesia and conscious sedation [7]; BT was performed according to established protocols (Boston Scientific Corp., Marlborough, MA, USA) [2, 13]. OCT images of the subsegmental branch of the right-lower lobe (RB8a&b and RB9a&b) were acquired prior to and immediately after BT as well as at 3 weeks, 6 weeks, 6 months and 2 years post-BT using a custom-built swept-source OCT system [14] and a C7 Dragonfly Imaging Catheter (St Jude Medical Inc., St Paul, MN, USA). OCT airway segments matched by visual inspection at each time-point were selected for analysis. The lumen area (A_i) and outer wall area (A_o) for three consecutive OCT slices were manually segmented (ImageJ; National Institutes of Health, Bethesda, MD, USA) to generate airway wall (WA) percentage:

$$WA = \frac{A_i}{A_o} \times 100$$

Patient A was a 51-year-old male with no smoking history and an asthma duration of 6 years. Patient B was a 56-year-old female and an ex-smoker at the time of the study (10 pack-years) with an asthma

duration of 27 years. Both patients reported similar forced expiratory volume in 1 s (FEV₁) (patient A: 1.63 L (53% pred); patient B: 1.52 L (54% pred)) as well as similar doses of inhaled steroids (4000 µg beclomethasone equivalent per day), salmeterol (100 µg per day) and tiotropium (18 µg per day). However, patient A reported higher doses of prednisone (patient A: 25 mg per day; patient B: 5–10 mg per day) and salbutamol (patient A: 3500–3700 µg per day; patient B: 300–500 µg per day) than patient B prior to BT.

Figure 1a shows OCT images pre-BT, immediately post-BT, and 3 weeks, 6 weeks, 6 months and 2 years post-BT for patients A and B. Patient A showed a remarkably thickened and inflamed epithelium, irregular basement membrane, and prominent smooth muscle pre-BT. Immediately post-BT, copious secretions and sloughing of the epithelium were observed. At 3 and 6 weeks post-BT, the epithelium appeared thinner and the lamina propria appeared thicker from deposition of collagen, and smooth muscle appeared slightly less prominent. At 6 months and 2 years post-BT, OCT showed recurrence of the inflammatory changes in the epithelium. In contrast to patient A, the epithelial layer of patient B was not inflamed pre-BT. Airway wall oedema was visually apparent 3 weeks post-BT but was reduced at week 6. The smooth muscle layer was also less prominent 6 weeks post-BT, and remained the same at 6 months and 2 years. At 6 months and 2 years post-BT, patient B also showed a normal bronchial epithelium with collagen deposition in the submucosa.

Figure 1b shows OCT airway measurements pre-BT, immediately post-BT, and 3 weeks, 6 weeks, 6 months and 2 years post-BT for patients A and B. For Patient A, airway secretions prevented OCT airway measurements immediately post-BT. At week 3, OCT airway measurements showed an increase in WA. Although WA improved slightly at week 6, WA was increased 6 months and 2 years post-BT. In patient A, there was a transient improvement in FEV₁ at 3 and 6 weeks post-BT (pre-BT: 1.63 L; 3 weeks post-BT: 2.14 L; 6 weeks post-BT: 2.27 L). However, symptoms recurred 4 months after treatment and patient A remained on 20–50 mg per day of prednisone due to recurrent exacerbations. FEV₁ was also worse 6 months and 2 years post-BT than pre-BT (6 months post-BT: 1.08 L; 2 years post-BT: 1.11 L).

As shown in figure 1b for patient B, OCT WA was reduced at 3 and 6 weeks compared to baseline. This reduction in WA persisted 6 months and 2 years post-BT. FEV₁ also progressively improved post-BT (pre-BT: 1.52 L; 3 weeks post-BT: 1.68 L; 6 weeks post-BT: 1.78 L; 6 months post-BT: 2.03 L) but dropped slightly 2 years post-BT (2 years post-BT: 1.82 L). Symptoms decreased along with discontinuation of prednisone and less frequent use of rescue medication.

Figure 1c shows histology of RB8 bronchial biopsies 6 months post-BT. For patient A, histology showed a partially denuded epithelium, thickened basement membrane, moderate-to-severe inflammation and remodelling of the submucosa by collagen deposition. In contrast, for patient B, histology of the same airway showed a normal bronchial epithelium with collagen deposition in the submucosa.

This pilot study illustrates, for the first time, that OCT can be used to evaluate *in vivo* airway remodelling longitudinally following BT in patients with severe asthma. Although the two asthmatics evaluated were similar in age and presented with similar clinical features and spirometry, the asthma patients demonstrated very different responses to BT as well as very distinct OCT airway wall features at baseline.

Following BT in patient A, despite evidence of FEV₁ improvement at weeks 3 and 6, OCT imaging showed no reduction in airway wall thickness and the patient's symptoms returned 4 months post-treatment. In contrast, patient B showed progressive improvements in FEV₁ for at least 6 months and reductions in OCT airway wall thickness, and demonstrated improvements in respiratory symptoms and decreased medication use for 2 years post-BT. Although patient A was a clear BT nonresponder, patient B demonstrated improvements following BT therapy that could not be predicted based on baseline spirometry or clinical features.

Although this study was limited to only two patients, it is certainly hypothesis generating and a number of questions arise from this pilot study. Most important is the question of whether OCT can identify characteristics within the airway wall that predict BT responders. Inflammation is visible on OCT [7, 15] and, notably, in this study, the BT responder showed thickened airway wall without evidence of inflammation in the epithelial layer at baseline compared to the nonresponder.

In summary, we evaluated two severe asthmatics immediately prior to and longitudinally following BT, and demonstrated a reduction in airway wall thickness that persisted 2 years following treatment in the BT responder, as well as differences in airway wall features between the responder and nonresponder prior to treatment. These observations generate hypotheses for a larger study to determine if airway changes defined by OCT imaging can identify asthma patients who will benefit from BT and to determine the long-term effects of the treatment.

FIGURE 1 a) Optical coherence tomography images (scale bars=1 mm) and b) mean±SD airway measurements prior to and immediately after bronchial thermoplasty (BT), 3 weeks post-BT, 6 weeks post-BT, 6 months post-BT and 2 years post-BT, with c) the corresponding bronchial biopsy at 6 months post-BT. Epi: epithelium; BM: basement membrane; SM: smooth muscle; WA: airway wall.

@ERSpublications

Optical coherence tomography small airway imaging may be used for better patient phenotyping and selection for BT <http://ow.ly/O0Cm2>

Miranda Kirby¹, Keishi Ohtani^{2,3}, Rosa Maria Lopez Lisbona^{3,4}, Anthony M.D. Lee³, Wei Zhang^{3,5}, Pierre Lane³, Nina Varfolomeva⁶, Linda Hui⁶, Diana Ionescu⁷, Harvey O. Coxson¹, Calum MacAulay³, J. Mark FitzGerald⁶ and Stephen Lam^{3,6}

¹Centre for Heart Lung Innovation, University of British Columbia and St. Paul's Hospital, Vancouver, BC, Canada.

²Department of Surgery, Tokyo Medical University, Tokyo, Japan. ³Imaging Unit, Integrative Oncology Department, British Columbia Cancer Research Centre, Vancouver, BC, Canada. ⁴Department of Respiratory, Bellvitge University Hospital, Hospitalet de Llobregat, Barcelona, Spain. ⁵Department of Respiratory and Critical Care Medicine, Peking University First Hospital, Beijing, China. ⁶Institute for Heart and Lung Health, University of British Columbia and Vancouver General Hospital, Vancouver, BC, Canada. ⁷Department of Pathology, British Columbia Cancer Agency and the University of British Columbia, Vancouver, BC, Canada.

Correspondence: Stephen Lam, British Columbia Cancer Research Centre, 675 West 10th Avenue, Vancouver, BC, V5Z 1L3, Canada. E-mail: slam2@bccancer.bc.ca

Received: Jan 29 2015 | Accepted after revision: April 10 2015 | First published online: May 28 2014

Support statement: M. Kirby gratefully acknowledges postdoctoral support from the Canadian Institutes of Health Research (CIHR) Bisby award, the CIHR Integrated and Mentored Pulmonary and Cardiovascular Training program (IMPACT), and the Michael Smith Foundation for Health Research (MSFHR). We also gratefully acknowledge the

CIHR–Thoracic Imaging Network of Canada and the United States Public Health Service contracts N01-CN 35000 from the National Cancer Institute for the OCT technology development. The funding sources for this study played no role in the design and conduct of the study; in the collection, management, analysis, and interpretation of the data; or in the preparation of the manuscript. The funding sources did not review the manuscript. S. Lam had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis. Funding information for this article has been deposited with FundRef.

Conflict of interest: Disclosures can be found alongside the online version of this article at erj.ersjournals.com

Acknowledgements: We thank Myles McKinnon, Jennifer Campbell and Chulho Hyun (British Columbia Cancer Agency, Vancouver, BC, Canada) for their assistance with the bronchoscopy procedures.

References

- 1 Wechsler ME, Laviolette M, Rubin AS, *et al.* Bronchial thermoplasty: Long-term safety and effectiveness in patients with severe persistent asthma. *J Allergy Clin Immunol* 2013; 132: 1295–1302.
- 2 Castro M, Rubin AS, Laviolette M, *et al.* Effectiveness and safety of bronchial thermoplasty in the treatment of severe asthma: a multicenter, randomized, double-blind, sham-controlled clinical trial. *Am J Respir Crit Care Med* 2010; 181: 116–124.
- 3 Hanania NA, Wenzel S, Rosen K, *et al.* Exploring the effects of omalizumab in allergic asthma: an analysis of biomarkers in the EXTRA study. *Am J Respir Crit Care Med* 2013; 187: 804–811.
- 4 Nakano Y, Wong JC, de Jong PA, *et al.* The prediction of small airway dimensions using computed tomography. *Am J Respir Crit Care Med* 2005; 171: 142–146.
- 5 Kauczor HU, Ebert M, Kreitner KF, *et al.* Imaging of the lungs using ³He MRI: preliminary clinical experience in 18 patients with and without lung disease. *J Magn Reson Imaging* 1997; 7: 538–543.
- 6 Coxson HO, Quiney B, Sin DD, *et al.* Airway wall thickness assessed using computed tomography and optical coherence tomography. *Am J Respir Crit Care Med* 2008; 177: 1201–1206.
- 7 Lam S, Standish B, Baldwin C, *et al.* *In vivo* optical coherence tomography imaging of preinvasive bronchial lesions. *Clin Cancer Res* 2008; 14: 2006–2011.
- 8 Hariri LP, Applegate MB, Mino-Kenudson M, *et al.* Volumetric optical frequency domain imaging of pulmonary pathology with precise correlation to histopathology. *Chest* 2013; 143: 64–74.
- 9 Williamson JP, McLaughlin RA, Noffsinger WJ, *et al.* Elastic properties of the central airways in obstructive lung diseases measured using anatomical optical coherence tomography. *Am J Respir Crit Care Med* 2011; 183: 612–619.
- 10 Coxson HO, Lam S. Quantitative assessment of the airway wall using computed tomography and optical coherence tomography. *Proc Am Thorac Soc* 2009; 6: 439–443.
- 11 Tsuboi M, Hayashi A, Ikeda N, *et al.* Optical coherence tomography in the diagnosis of bronchial lesions. *Lung Cancer* 2005; 49: 387–394.
- 12 British Thoracic Society, Scottish Intercollegiate Guidelines Network. British guideline on the management of asthma. *Thorax* 2014; 69: 1–192.
- 13 Cox G, Miller JD, McWilliams A, *et al.* Bronchial thermoplasty for asthma. *Am J Respir Crit Care Med* 2006; 173: 965–969.
- 14 Lee AM, Ohtani K, Macaulay C, *et al.* *In vivo* lung microvasculature visualized in three dimensions using fiber-optic color Doppler optical coherence tomography. *J Biomed Opt* 2013; 18: 50501.
- 15 Whiteman SC, Yang Y, Gey van Pittius D, *et al.* Optical coherence tomography: real-time imaging of bronchial airways microstructure and detection of inflammatory/neoplastic morphologic changes. *Clin Cancer Res* 2006; 12: 813–818.

Eur Respir J 2015; 46: 859–862 | DOI: 10.1183/09031936.00016815 | Copyright ©ERS 2015

Effective strategies for managing new *Pseudomonas* cultures in adults with cystic fibrosis

To the Editor:

Pseudomonas aeruginosa is a common chronic pulmonary infection in cystic fibrosis (CF). It is associated with poor outcomes and new detection should prompt an early eradication attempt [1–3].

Strict segregation and eradication policies have led to a declining prevalence of chronic *P. aeruginosa* infection, with only 30% of CF adolescents having chronic infection when they transition to adult services [4]. Adult CF physicians increasingly manage new *P. aeruginosa* cultures in adults without robust evidence to guide decision-making. We evaluated the success of our eradication and suppression strategies for adults