Elisa López-Varela^{1,2}, Orvalho Joaquim Augusto¹, Luis Guerra¹, Durval Respeito¹, Charfudin Saco¹, Jahit Sacarlal¹, Giovanni Battista Migliori³, Giovanni Sotgiu⁴, Pedro L. Alonso^{1,2} and Alberto L. García-Basteiro^{1,2,5}
¹Centro de Investigação em Saúde de Manhiça (CISM), Maputo, Mozambique. ²Instituto de Salud Global de Barcelona (ISGLOBAL), Barcelona, Spain. ³WHO Collaborating Centre for TB and Lung Diseases, Fondazione S. Maugeri, Care and Research Institute, Tradate, Italy. ⁴Clinical Epidemiology and Medical Statistics Unit, Dept of Biomedical Sciences, University of Sassari – Research, Medical Education and Professional Development Unit, AOU Sassari, Sassari, Italy. ⁵Amsterdam Institute for Global Health and Development, Academic Medical Centre, Amsterdam, The Netherlands.

Correspondence: Elisa López-Varela, Centro de Investigação em Saúde de Manhiça (CISM), Maputo, Mozambique.
 E-mail: elisa.lopez@manhica.net

Received: Sept 01 2015 | Accepted after revision: Nov 09 2015 | First published online: Dec 23 2015

Conflict of interest: None declared.

Acknowledgements: The authors thank the participants and their families. The authors are grateful to the National TB Program (Maputo, Mozambique), Denise Naniche (ISGLOBAL, Barcelona, Spain), Alberto Bila Junior (CISM, Manhiça, Mozambique), health staff at the Manhiça District Hospital (Manhiça, Mozambique) and peripheral health centres for their contribution.

References

- 1 World Health Organization. Global Tuberculosis Report 2014. WHO/HTM/TB/2014.08. Geneva, World Health Organization, 2014. Available from: http://www.who.int/tb/publications/global_report/gtbr14_main_text.pdf
- 2 Marais BJ, Graham SM, Maeurer M, *et al.* Progress and challenges in childhood tuberculosis. *Lancet Infect Dis* 2013; 13: 287–289.
- 3 Murray CJ, Ortblad KF, Guinovart C, *et al.* Global, regional, and national incidence and mortality for HIV, tuberculosis, and malaria during 1990–2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet* 2014; 384: 1005–1070.
- 4 Jenkins HE, Tolman AW, Yuen CM, *et al.* Incidence of multidrug-resistant tuberculosis disease in children: systematic review and global estimates. *Lancet* 2014; 383: 1572–1579.
- 5 Dodd PJ, Gardiner E, Coghlan R, *et al.* Burden of childhood tuberculosis in 22 high-burden countries: a mathematical modelling study. *Lancet Glob Health* 2014; 2: e453–e459.
- 6 Lönnroth K, Migliori GB, Abubakar I, *et al.* Towards tuberculosis elimination: an action framework for low-incidence countries. *Eur Respir J* 2015; 45: 928–952.
- 7 García-Basteiro AL, López-Varela E, Respeito D, *et al.* High tuberculosis burden among people living with HIV in southern Mozambique. *Eur Respir J* 2015; 45: 547–549.
- 8 Saco C, Nhacolo A, Nhalungo D, *et al.* Profile: Manhiça Health Research Centre (Manhiça HDSS). *Int J Epidemiol* 2013; 42: 1309–1318.
- 9 López-Varela E, Augusto OJ, Gondo K, *et al.* Incidence of tuberculosis among young children in rural Mozambique. *Pediatr Infect Dis J* 2015; 34: 686–692.
- 10 García-Basteiro AL, López-Varela E, Augusto OJ, *et al.* Radiological findings in young children investigated for tuberculosis in Mozambique. *PLoS One* 2015; 10: e0127323.
- 11 Rudolf F, Wejse C. Tuberculosis case detection revisited: better testing might not improve outcomes. *Lancet Glob Health* 2015; 3: e424–e425.
- 12 Gibbons CL, Mangan MJ, Plass D, *et al.* Measuring underreporting and under-ascertainment in infectious disease datasets: a comparison of methods. *BMC Public Health* 2014; 14: 147.
- 13 Cruz AT, Starke JR. What's in a number? Accurate estimates of childhood tuberculosis. *Lancet Glob Health* 2014; 2: e432–e433.
- 14 Perez-Velez CM, Marais BJ. Tuberculosis in children. *N Engl J Med* 2012; 367: 348–361.

Eur Respir J 2016; 47: 1003–1005 | DOI: 10.1183/13993003.01454-2015 | Copyright ©ERS 2016

Polymers of Z α_1 -antitrypsin are secreted in cell models of disease

To the Editor:

The α_1 -antitrypsin (α_1 -AT) is a 52 kDa glycoprotein that is predominantly synthesised in the liver and secreted into the circulation, where it protects the lungs from the enzyme neutrophil elastase. α_1 -AT deficiency (α_1 -ATD) is caused by mutations in the α_1 -AT gene, with most cases resulting from homozygous inheritance of the Z allele (Glu342Lys). This leads to low levels of circulating α_1 -AT, uncontrolled elastase activity and emphysema [1]. The Z mutation destabilises the native α_1 -AT and causes the formation of aberrant polymers that accumulate within the endoplasmic reticulum (ER) of

hepatocytes, giving rise to inclusion bodies that are the main histological feature of α_1 -ATD [2]. Extracellular polymers have been found in lung lavage, the skin of an individual with panniculitis and the kidney of an individual with vasculitis [1], and are also present in the circulation of all individuals homo- or heterozygous for the Z allele [3]. Circulating polymers originate in the liver, since they became undetectable in the plasma of an individual 4 days after liver transplantation [3], but it is unknown whether polymers can be secreted from hepatocytes or can form in the extracellular environment from secreted monomeric Z α_1 -AT. Extracellular polymers are chemotactic and stimulatory for human neutrophils [4] and may contribute to inflammatory neutrophil infiltration in the lungs, kidney and skin. It is important to understand where these polymers form in order to design effective therapies for emphysema and other pathological manifestations of α_1 -ATD. Here we investigated the origin of extracellular polymers by exploiting our cellular models of α_1 -ATD and conformer-specific and functional monoclonal antibodies (mAb) against Z α_1 -AT [5–8].

We first confirmed the presence of α_1 -AT polymers in the plasma of three PiZZ individuals by non-denaturing PAGE and the polymer-specific mAb-2C1 [5] (figure 1a). Similarly, when expressed in human embryonic kidney (HEK) 293 T-cells Z α_1 -AT was detected in the culture medium as a mix of monomer and polymers (figure 1b). To exclude the possibility that polymers were released into the culture medium from damaged cells, we analysed the glycosylation state of extracellular α_1 -AT. Secretory glycoproteins are synthesised in the ER, modified in the Golgi apparatus where N-linked glycans increase in size and complexity, and arrive to the trans-Golgi network (TGN) for sorting and secretion. We ascertained the state of N-linked glycosylation by treating both M and Z α_1 -AT with endoglycosidase H (endo H) that removes pre-Golgi glycans or peptide-N-glycosidase F (PNGase F) that removes all N-linked glycans. Digestion with endo H showed that all extracellular α_1 -AT contained only mature, endo H-resistant N-glycans, which were removed by PNGase F giving the expected size for non-glycosylated α_1 -AT (figure 1c), demonstrating that extracellular polymers bare mature N-glycans, resulting from their transport through the Golgi or from extracellular polymerisation of monomeric Z α_1 -AT, but not from direct release of immature ER proteins.

We next used our polymerisation-blocking mAb-4B12 [8] to prevent Z α_1 -AT polymerisation in cell culture medium. This antibody blocks polymer formation at a molar ratio of 1:1 for Z α_1 -AT and mAb-4B12 *in vitro*, so adding it in excess to the culture medium should inhibit the polymerisation of secreted monomeric Z α_1 -AT. Chinese hamster ovary (CHO) cells expressing Z α_1 -AT under a Tet-inducible promoter [7] were cultured for 24 h with or without mAb-4B12 at a molar ratio of 1:1 or 1:2 for Z α_1 -AT and mAb-4B12. There was no difference in polymeric Z α_1 -AT in the culture media by ELISA (figure 1d), indicating that polymers were secreted from the cells. The presence of a complex between secreted Z α_1 -AT and mAb-4B12 was confirmed by ELISA, demonstrating binding of mAb-4B12 to secreted Z α_1 -AT (not shown). The secretory origin of extracellular polymers was further supported by the observation that culture medium from CHO cells expressing Z-A1AT induced for 48 h and further incubated in the absence of cells for 72 h showed no increase in polymer levels (not shown).

M α_1 -AT is readily secreted and can be detected by immunofluorescence within the Golgi compartment [5, 7, 9–11]. Although polymers of Z and other mutant variants of α_1 -AT are found in the culture medium of expressing cells, they have never been shown to co-localise with Golgi-resident proteins [5, 7, 10, 11]. This may be due to very low levels of these proteins transiting the Golgi at steady state, and so we used a temperature block by culturing cells at 20°C, which reduces the exit of secretory proteins from the Golgi apparatus without affecting ER to Golgi transport [12]. Secretion of M and Z α_1 -AT from transiently transfected COS-7 cells was reduced 5- and 2-fold upon incubation at 20°C for 4 h (data not shown). In these conditions, we assessed the co-localisation of 2C1-positive polymers with resident proteins of the secretory pathway: BiP (ER), giantin (cis-medial-Golgi) and TGN-46 (TGN). The majority of 2C1-positive Z α_1 -AT co-localised with BiP, confirming polymer accumulation within the ER [5, 7], but we also observed partial co-localisation with giantin and TGN-46 (figure 1e), supporting their transiting through the Golgi apparatus and exiting the cells *via* the canonical secretory pathway.

In order to confirm our results we used myc- and haemagglutinin epitope (HA)-tagged versions of M and Z α_1 -AT (M_{myc}, Z_{myc}, M_{HA} and Z_{HA}) to transiently transfect from HEK293 T-cells. We expressed each version of Z α_1 -AT or different mixes of the M and Z α_1 -AT constructs, as described in figure 1f. We also created a new experimental condition by collecting, mixing and replating cells transfected separately with Z_{HA} and Z_{myc} (figures 1f and g, lane 4). All the transfections were successful in producing similar amounts of α_1 -AT (figure 1f top panel), monomeric for the M α_1 -AT and polymeric for the Z α_1 -AT constructs (figure 1f, bottom panel). We then performed immunoprecipitation of the culture media with anti-HA antibodies, followed by Western blot with an anti-myc-tag antibody and an anti-total α_1 -AT antibody in the same membrane. All samples containing Z_{HA} gave a positive signal with the anti- α_1 -AT antibody, demonstrating an efficient immunoprecipitation (figure 1g, bottom panel), but only cells

FIGURE 1 Extracellular polymers of Z α_1 -antitrypsin (α_1 -AT) do not form in the extracellular medium, but are secreted from cells via the Golgi complex. **a)** Polymers of Z α_1 -AT were found in the plasma of three PiZZ α_1 -AT deficiency individuals but not in plasma from a control donor by non-denaturing PAGE (stacking gel pH 6.8, separating gel pH 8.8) and Western blot analysis using the polymer-specific monoclonal antibodies (mAb) 2C1. **b)** The culture medium of human embryonic kidney (HEK) 293 T-cells expressing M α_1 -AT contained only monomeric protein, while Z α_1 -AT was found as monomeric and polymeric forms after non-denaturing PAGE and Western blot analysis for total α_1 -AT. The differences in band patterns and background levels between the polymer ladders in **a)** and **b)** are probably due to high protein concentrations in plasma samples compared to serum-free culture medium supernatants. **c)** Cell lysates (Lys) and culture media (Med) of cells in **b)** treated or not treated (-) with endoglycosidase H (Endo H) or peptide-N-glycosidase F (PNGase F) and analysed by SDS PAGE and Western blot for total α_1 -AT. All extracellular α_1 -AT carried mature N-glycosylation (55 kDa), while intracellular α_1 -AT showed immature N-glycosylation (H) and deglycosylated α_1 -AT (I). **d)** Expression of Z α_1 -AT was induced in Chinese hamster ovary stable cells with 0.5 $\mu\text{g}\cdot\text{mL}^{-1}$ doxycycline (Dox) and simultaneously incubated with purified mAb-4B12 at a molar ratio of 1:1 or 1:2 for Z α_1 -AT and mAb-4B12 for 24 h; the amount of extracellular polymers as quantified by sandwich ELISA (2C1-Ag-9C5-HRP) did not change in the presence of the polymerisation-blocking antibody. **e)** Immunodetection of Z α_1 -AT polymers with mAb-2C1 (red) in co-localisation with resident proteins (green) of the endoreticulum (BiP), cis/medial Golgi (Giantin) and trans-Golgi network (TGN-46); the cell nucleus was counter-stained with diaminophenylindole. Scale bars=10 μm and 5 μm in the main panel and close-up panel, respectively. All panels were obtained in a Zeiss LSM510META confocal microscope (Zeiss, Jena, Germany) with a $\times 63$ oil objective. **f)** HEK293 T-cells were transiently transfected with vectors encoding Z_{HA} (lane 1) or Z_{myc} (lane 2), or co-transfected with Z_{HA}/Z_{myc} (lane 3), M_{HA}/M_{myc} (lane 5) or Z_{HA}/M_{myc} (lane 6); an additional condition was created by detaching, mixing and replating equal numbers of Z_{HA} and Z_{myc} cells (lane 4) at the end of the fifth transfection. After an overnight incubation in normal culture medium, cells were further incubated for 24 h in serum-free medium, and culture media were analysed by SDS- or non-denaturing PAGE. **g)** Culture media as in **f)** were immunoprecipitated with an anti-haemagglutinin (HA) antibody and analysed by SDS PAGE and immunoblot with an anti-myc HRP mAb (top panel), then stripped and re-probed with the 2G7 anti-total α_1 -AT mAb (bottom panel). **h)** Plasma from four different ZZ patient donors was incubated for the times indicated at 37°C and analysed by sandwich ELISA with the mAb-2C1. The graph shows the amount of Z α_1 -AT polymers normalised to time 0.

co-transfected with Z_{HA} and Z_{myc} showed a band positive to the anti-myc antibody, demonstrating the presence of Z_{HA}/Z_{myc} heteropolymers in their culture medium (fig. 1g top panel). In contrast, cells transfected separately with Z_{HA} and Z_{myc} that were afterwards cultured together showed no signal with the anti-myc antibody. These results support the intracellular origin of secreted polymers.

Taken together, our results indicate that polymers formed within the ER can traffic through this organelle and be secreted *via* the Golgi compartment, where N-glycans acquire their mature configuration. Our biochemical and immunofluorescence results do not exclude polymer formation in post-ER compartments. The existing literature and our present data are in keeping with polymerisation occurring mostly within the ER, where insoluble long-chain polymers are trapped within inclusion bodies, while at least part of the soluble, short-chain polymers can enter transport vesicles for secretion. Our present observations suggest that polymers secreted from hepatocytes can contribute to circulating polymers. Although the concentration of monomeric Z α_1 -AT in plasma is probably high enough to allow extracellular polymerisation, we did not observe an increase in polymer levels when plasma samples from PiZZ patients were incubated at 37°C for 72 h (figure 1h). This can be due to limitations in our experimental conditions or to the presence of polymerisation-inhibitory factors in plasma. Our results are also compatible with local, extracellular polymer formation in the lung during inflammation and particularly by exposure to cigarette smoke [13]. In conclusion, our present results show for the first time that polymers of Z α_1 -AT are secreted from cells, supporting a contribution of intracellular origin to circulating polymers in α_1 -AT D.

@ERSpublications

Pathological polymers of Z alpha1-AT can be secreted from cells through the canonical secretory pathway <http://ow.ly/WHxZh>

Annamaria Fra¹, Francesca Cosmi^{2,8}, Adriana Ordoñez^{3,8}, Romina Berardelli¹, Juan Perez⁴, Noemi A. Guadagno², Luciano Corda⁵, Stefan J. Marciniak³, David A. Lomas⁶ and Elena Miranda^{2,7}

¹Dept of Molecular and Translational Medicine, University of Brescia, Brescia, Italy. ²Dept of Biology and Biotechnologies 'Charles Darwin', Sapienza University of Rome, Rome, Italy. ³Dept of Medicine, Cambridge Institute for Medical Research, University of Cambridge, Cambridge, UK. ⁴Dept of Cell Biology, Genetics and Physiology, University of Malaga, Malaga, Spain. ⁵Division of Internal Medicine, Spedali Civili, Brescia, Italy. ⁶Wolfson Institute for Biomedical Research, University College London, London, UK. ⁷Pasteur Institute – Cenci Bolognetti Foundation, Sapienza University of Rome, Rome, Italy. ⁸Both authors contributed equally to this work.

Correspondence: Annamaria Fra, Department of Molecular and Translational Medicine, University of Brescia, Viale Europa 11, Brescia, 25123, Italy. E-mail: annamaria.fra@unibs.it

Received: June 16 2015 | Accepted after revision: Dec 15 2015 | First published online: Feb 04 2016

Support statement: A. Fra is supported by Fondazione Cariplo (Italy), grant number 2013-0967. F. Cosmi was supported by a mobility grant from the Italian ABCD association. D.A. Lomas has funding provided by the Medical Research Council (UK) and the UK NIHR UCLH Biomedical Research Centre. E. Miranda was supported by the Pasteur Institute – Cenci Bolognetti Foundation (Italy) and by the Telethon Foundation (Italy) grant number GGP11057.

Conflict of interest: Disclosures can be found alongside the online version of this article at erjersjournals.com

Acknowledgements: We thank Giulia Baldini (University of Arkansas for Medical Sciences, Little Rock, AR, USA) for providing tagged α_1 -antitrypsin vectors, Riccardo Ronzoni (University of Brescia, Brescia, Italy) for sharing reagents, Roberto Sitia (San Raffaele Scientific Institute, Milan, Italy), Bibek Gooptu (King's College London, London, UK) and James Irving (University College London, London, UK) for helpful discussions, and Nuccia Gatta (Associazione Nazionale Alpha1 AT Onlus, Brescia, Italy) and the Italian Alpha1 Patients' Association for constant encouragement.

References

- 1 Gooptu B, Dickens JA, Lomas DA. The molecular and cellular pathology of α_1 -antitrypsin deficiency. *Trends Mol Med* 2014; 20: 116–127.
- 2 Gooptu B, Lomas DA. Conformational pathology of the serpins: themes, variations, and therapeutic strategies. *Annu Rev Biochem* 2009; 78: 147–176.
- 3 Tan L, Dickens JA, Dimeo DL, *et al.* Circulating polymers in α_1 -antitrypsin deficiency. *Eur Respir J* 2014; 43: 1501–1504.
- 4 Mulgrew AT, Taggart CC, Lawless MW, *et al.* Z α_1 -antitrypsin polymerizes in the lung and acts as a neutrophil chemoattractant. *Chest* 2004; 125: 1952–1957.
- 5 Miranda E, Perez J, Ekeowa UI, *et al.* A novel monoclonal antibody to characterize pathogenic polymers in liver disease associated with alpha1-antitrypsin deficiency. *Hepatology* 2010; 52: 1078–1088.
- 6 Fra AM, Gooptu B, Ferrarotti I, *et al.* Three new alpha1-antitrypsin deficiency variants help to define a C-terminal region regulating conformational change and polymerization. *PLoS One* 2012; 7: e38405.
- 7 Ordoñez A, Snapp EL, Tan L, *et al.* Endoplasmic reticulum polymers impair luminal protein mobility and sensitize to cellular stress in α_1 -antitrypsin deficiency. *Hepatology* 2013; 57: 2049–2060.
- 8 Ordoñez A, Perez J, Tan L, *et al.* A single-chain variable fragment intrabody prevents intracellular polymerization of Z α_1 -antitrypsin while allowing its antiprotease activity. *FASEB J* 2015; 29: 2667–2678.
- 9 Novorodovskaya N, Lee J, Yu ZX, *et al.* Inhibition of intracellular degradation increases secretion of a mutant form of alpha1-antitrypsin associated with profound deficiency. *J Clin Invest* 1998; 101: 2693–2701.

- 10 Granell S, Baldini G, Mohammad S, *et al.* Sequestration of mutated α_1 -antitrypsin into inclusion bodies is a cell-protective mechanism to maintain endoplasmic reticulum function. *Mol Biol Cell* 2008; 19: 572–586.
- 11 Medicina D, Montani N, Fra AM, *et al.* Molecular characterization of the new defective P_{brescia} alpha1-antitrypsin allele. *Hum Mutat* 2009; 30: E771–E781.
- 12 Matlin KS, Simons K. Reduced temperature prevents transfer of a membrane glycoprotein to the cell surface but does not prevent terminal glycosylation. *Cell* 1983; 34: 233–243.
- 13 Alam S, Li Z, Janciauskiene S, *et al.* Oxidation of Z α_1 -antitrypsin by cigarette smoke induces polymerization: a novel mechanism of early-onset emphysema. *Am J Respir Cell Mol Biol* 2011; 45: 261–269.

Eur Respir J 2016; 47: 1005–1009 | DOI: 10.1183/13993003.00940-2015 | Copyright ©ERS 2016