Correspondence regarding the article “The asbestos fibre burden in human lungs: new insights into the chrysotile debate”

To the Editor:

The article by FEDER *et al.* [1] states that the lung asbestos fibre burden in 23 955 patients was analysed to address fibre type and biopersistence; data from 12 patients undergoing two tissue excisions at intervals at least 4 years were considered.

We believe that the article has serious shortcomings, as follows.

1) *Unclear aim.* Contrary to the authors' claim, there is no ongoing debate about the biopersistence of chrysotile asbestos among independent, credible scientists. In support of their claim that such a debate exists, the authors rely on an article commissioned, funded and developed in collaboration with asbestos lobbyists.

2) *Faulty study design.* Significant scientific problems in patient/sample selection and applied methods exist. First, the small sample size: only 12 (0.05%) of the 23 955 cases were analysed with two investigations; only six had electron microscopic examination of tissue. Second, the selection criterion of 500 asbestos bodies per gramme of wet lung is discretionary and arbitrary. Third, relationships between outcome and fibre-years were not examined using a detailed occupational history. No statistical analysis accounting for occupation/exposures, interim exposures and latency periods, and exposure changes over recent decades is reported.

3) *Methods.* The authors do not explain why they used both field energy (FE) scanning electron microscopy (SEM) and transmission electron microscopy (TEM), nor is it clear which data come from which method in the supplementary table. TEM is regarded as the method of choice by impartial and credible pathologists; the limitations of SEM have been discussed previously [2].

The use of FE-SEM and TEM on autopsy specimens alone does not allow the authors to say anything about the change in the number of chrysotile fibres over time.

The term “pulmonary asbestos fibre concentration” defined as “total of asbestos bodies and bare fibres” is incorrectly conflated with asbestos fibre concentration or burden as generally understood.

4) *Results.* First, there is no information on fibre length, which would be crucial for understanding outcome since short fibres are more rapidly cleared from the lung [3, 4].

Second, in the six cases analysed with electron microscopy, the fibre counts reported are generally similar to, or even lower than, asbestos body counts. This unusual finding contradicts the literature where total fibres outnumber asbestos bodies by three orders of magnitude [5].

Third, in a large number of ferruginous bodies, the authors were unable to identify the core material. How did the authors define a ferruginous body as asbestos and non-asbestos in those cases not subjected to electron microscopy?

Last, the authors mention that “asbestos grading followed national and international criteria, *i.e.* primarily the Helsinki criteria”. It is unclear which criteria were used in each case. There are significant differences

@ERSpublications

All asbestos types cause asbestosis (cancer): chrysotile is not biopersistent, so fibre analysis is not diagnostic <http://ow.ly/BOLC30grqYg>

Cite this article as: Oliver LC, Belpoggi F, Budnik LT, *et al.* Correspondence regarding the article “The asbestos fibre burden in human lungs: new insights into the chrysotile debate”. *Eur Respir J* 2017; 50: 1701644 [<https://doi.org/10.1183/13993003.01644-2017>].

between the NIOSH/CAP definition and the definition of Roggli and co-workers and its respective formulation in the Helsinki criteria 2014 [6] (for further details, see [7]).

5) *Data analysis, data interpretation and conclusions.* First, in the six cases with fibre type differentiation, 10% to 95% were reported to represent chrysotile. In Germany, about 94% of asbestos used was chrysotile. The relative paucity of chrysotile fibres in cases 1 and 2 (33% of the cases) with intervals of 14 and 21 years between the first and second examinations is consistent with low biopersistence of chrysotile fibres.

Second, the authors state “Thus fibre clearance and biopersistence are considered the most important factors for diagnostics and risk assessment of malignant and non-malignant diseases.” In fact, diagnosis is based mainly on a thorough occupational history and noninvasive clinical findings; risk assessment is related to fibre concentration in the workplace [8].

Third, the major interpretation of the data by the authors is that chrysotile fibre counts, like those of amphibole fibres, do not change over time in the human lung. Their findings and extensive literature show just the opposite [8].

6) *Discussion.* The authors claim an ongoing debate about the hazardous nature of chrysotile. Publications not cited by FEDER *et al.* [1] (IARC, WHO), and other professional bodies and government agencies contradict this claim, concluding that chrysotile asbestos exposure increases risk for asbestosis, mesothelioma, lung and other cancers [9].

7) *Medico-legal relevance.* There is grave risk that the publication by FEDER *et al.* [1] will influence outcomes in the adjudication of asbestos-related disease in the legal system and that, as a result, the injured worker will suffer unfairly and unjustly. This risk would apply to those with a history of occupational exposure to chrysotile asbestos and in whom few or no asbestos fibres are found in the lung years later. The claim by FEDER *et al.* [1] that chrysotile fibres are biopersistent in the lung could be used in courts of law to deny justice to asbestos-harmed victims.

8) *Conflicts of interest.* The authors fail to disclose significant financial conflicting interests [10].

In conclusion, FEDER *et al.* [1] provide misleading findings that fail to refute the generally accepted tenet that chrysotile asbestos fibres are not biopersistent in the human lung. Neither their clinical nor their statistical analyses, nor the literature, support their claims.

L. Christine Oliver^{1,2}, Fiorella Belpoggi³, Lygia T. Budnik^{4,5}, David Egilman⁶, Arthur L. Frank⁷, Daniele Mandrioli³, Colin L. Soskolne^{8,9}, Benedetto Terracini¹⁰, Laura Welch¹¹ and Xaver Baur⁴

¹Dalla Lana School of Public Health, University of Toronto, Toronto, ON, Canada. ²Occupational Health Initiatives, Inc., Brookline, MA, USA. ³Istituto Ramazzini, Bentivoglio, Italy. ⁴European Society for Environmental and Occupational Medicine, Berlin, Germany. ⁵University of Hamburg, Hamburg, Germany. ⁶Alpert Medical School Dept of Medicine, Brown University, Attleboro, MA, USA. ⁷Drexel University School of Public Health and College of Medicine, Philadelphia, PA, USA. ⁸University of Alberta, Edmonton, AB, Canada. ⁹Health Research Institute, University of Canberra, Canberra, Australia. ¹⁰Unit of Cancer Epidemiology, CPO Piemonte and University of Turin, Turin, Italy. ¹¹George Washington University School of Public Health and Health Sciences Center for Construction Research and Training, Silver Spring, MD, USA.

Correspondence: Xaver Baur, European Society for Environmental and Occupational Medicine (EOM), P.O. Box 370514, 14135 Berlin, Germany. E-mail: baur@eomsociety.org

Received: Aug 12 2017 | Accepted: Oct 25 2017

Conflict of interest: Disclosures can be found alongside this article at erj.ersjournals.com

References

- 1 Feder IS, Tischoff I, Theile A, *et al.* The asbestos fibre burden in human lungs: new insights into the chrysotile debate. *Eur Respir J* 2017; 49: 1602534.
- 2 Health Effects Institute. Asbestos in public and commercial buildings: a literature review and synthesis of current knowledge. Cambridge, Health Effects Institute (HEI); Asbestos Research, 1991.
- 3 Churg A, Wiggs B. Fiber size and number in workers exposed to processed chrysotile asbestos, chrysotile miners, and the general population. *Am J Ind Med* 1986; 9: 143–152.
- 4 Abraham JL Analysis of fibrous and non-fibrous particles. Environmental and occupational medicine. 4th Edn. Philadelphia, Lippincott Williams and Wilkins, 2006; pp. 277–297.
- 5 Dodson RF, O'Sullivan M, Corn CJ, *et al.* Analysis of asbestos fiber burden in lung tissue from mesothelioma patients. *Ultrastruct Pathol* 1997; 21: 321–336.
- 6 Wolff H, Vehmas T, Oksa P, *et al.* Asbestos, asbestosis, and cancer, the Helsinki criteria for diagnosis and attribution 2014: recommendations. *Scand J Work Environ Health* 2015; 41: 5–15.
- 7 Hammar SP, Abraham JL. Commentary on pathologic diagnosis of asbestosis and critique of the 2010 Asbestos Committee of the College of American Pathologists (CAP) and Pulmonary Pathology Society's (PPS) update on the diagnostic criteria for pathologic asbestosis. *Am J Ind Med* 2015; 58: 1034–1039.

- 8 Ramazzini C. Comments on the 2014 Helsinki consensus report on asbestos. *J Occup Health* 2016; 58: 224–227.
- 9 Baur X, Soskolne CL, Lemen RA, *et al.* How conflicted authors undermine the World Health Organization (WHO) campaign to stop all use of asbestos: spotlight on studies showing that chrysotile is carcinogenic and facilitates other non-cancer asbestos-related diseases. *Int J Occup Environ Health* 2015; 21: 176–179.
- 10 Baur X. Response to the letter of R. Merget, I. Feder and A. Tannapfel. *Pneumologie* 2017; 71: 121–123.

Copyright ©ERS 2017

To the Editor:

We read with interest the generally excellent article by FEDER *et al.* [1] published in the *European Respiratory Journal* and we would like to add just a comment. Authors screened the German Mesothelioma Register for patients with asbestos body (AB) counts ≥ 500 per gramme of wet lung (corresponding to approximately ≥ 5000 AB per gramme of dry lung tissue) which had been analysed twice from different tissue excisions at minimum interval of 4 years. In the 12 patients with longitudinal data the asbestos fibre burden in the lung tissue was stable in particular for chrysotile. Authors stated that the study was the first to present intra-individual longitudinal data about the asbestos fibre burden in living human lungs.

The measure of the fibre load of lung tissue using electron microscopy represents the best indicator of retained dose, but this can only be performed after open lung biopsy, lung surgery or death. Mineralogical analysis of bronchoalveolar lavage fluid (BALF) by electron microscopy has been successfully used as a marker of asbestos fibre load in a number of studies [2–6]. Its use to characterise asbestos lung burden is generally accepted [7, 8]. In 2007 we published a study whose purpose was to assess the reliability of asbestos fibre concentration in BALF as a marker of past asbestos exposure by carrying out at different times the mineralogical analysis of BALF in the same patient and comparing the results [9]. Mineralogical analysis of BALF was carried out in 22 patients who underwent diagnostic fibreoptic bronchoscopy twice (the first to assess the past asbestos exposure, the second for different clinical reasons). The mean lag time between the first and the second bronchoalveolar lavage was 4.0 ± 2.3 years (median 4, range 1–10 years). In 16 patients (72.7%), a reduction of concentration in BALF of both chrysotile and amphiboles was observed, but the differences were not statistically significant while a significant decrease in AB concentration between the first and the second bronchoalveolar lavage was found.

Although the article of FEDER *et al.* [1] is not really the first to consider longitudinal individual data, it is of a great interest because it confirms histologically in the lung tissue what has been previously observed in the BALF. Moreover in this case the lag time from the cessation of exposure was much greater. The nonsignificant reduction of chrysotile and amphiboles fibres between the first and the second bronchoalveolar lavage observed in many cases was not apparently attributable to the pulmonary clearance because it was not related to the lag time between the first and the second BAL. Maybe it could be partially explained by the effect of the first lavage when the second was performed in the same lung region. Workers who have been exposed for a long time have concentrations of fibres in BALF which are higher than subjects who have been more recently exposed [4, 5]. Even if this information leads back to the high exposure in the past, it confirms the biopersistence of asbestos fibres in the human lung, in agreement with the results of FEDER *et al.* [1]. We agree with the authors that the sustained presence of both chrysotile and amphibole fibres causes lung diseases even many years after exposure cessation. This could also cause a different latency of asbestos-related lung cancer in comparison to those related to smoking.

Pietro Sartorelli

Dept of Medical Biotechnology, University of Siena, Unit of Occupational Health Azienda Ospedaliera Universitaria Senese, Siena, Italy.

Correspondence: Pietro Sartorelli, Dept of Medical Biotechnology, University of Siena, Unit of Occupational Health Azienda Ospedaliera Universitaria Senese, Viale Bracci 16, 53100 Siena, Italy. E-mail: pietro.sartorelli@unisi.it

@ERSpublications

Longitudinal intra-individual data demonstrate the high biopersistence of both amphiboles and chrysotile in the lung <http://ow.ly/ksnk30guepj>

Cite this article as: Sartorelli P. Correspondence regarding the article “The asbestos fibre burden in human lungs: new insights into the chrysotile debate”. *Eur Respir J* 2017; 50: 1702188 [<https://doi.org/10.1183/13993003.02188-2017>].

Received: Oct 24 2017 | Accepted: Oct 25 2017

Conflict of interest: None declared.

References

- 1 Feder IS, Tischoff I, Theile A, *et al.* The asbestos fibre burden in human lungs: new insights into the chrysotile debate. *Eur Respir J* 2017; 49: 1602534.
- 2 De Vuyst P, Dumortier P, Moulin E, *et al.* Diagnostic value of asbestos bodies in bronchoalveolar lavage fluid. *Am Rev Respir Dis* 1987; 136: 1219–1224.
- 3 Dodson R, Garcia J, O'Sullivan M, *et al.* The Usefulness of bronchoalveolar lavage in identifying past occupational exposure to asbestos: a light and electron microscopy study. *Am J Ind Med* 1991; 19: 619–628.
- 4 Sartorelli P, Scancarello G, Romeo R, *et al.* Asbestos exposure assessment by mineralogical analysis of bronchoalveolar lavage fluid. *J Occup Environ Med* 2001; 43: 872–881.
- 5 Romeo R, Scancarello G, Cassano P, *et al.* Assessment of asbestos exposure via mineralogical analysis of bronchoalveolar lavage fluid. *Med Lav* 2004; 95: 17–32.
- 6 Montomoli L, Romeo R, Sisinni AG, *et al.* Dose and effect biomarkers in the diagnosis of asbestos-related diseases and health surveillance of exposed workers. *G It Med Lav Erg* 2012; 34: Suppl. 3, 577–580.
- 7 Roggli VL, Gibbs AR, Attanoos R, *et al.* Pathology of asbestosis – an update of the diagnostic criteria: Report of the asbestosis committee of the College of American Pathologists and Pulmonary Pathology Society. *Arch Pathol Lab Med* 2010; 134: 462–480.
- 8 Wolff H, Vehmas T, Oksa P, *et al.* Asbestos, asbestosis, and cancer, the Helsinki criteria for diagnosis and attribution 2014: recommendations. *Scand J Work Environ Health* 2015; 41: 5–15.
- 9 Sartorelli P, Romeo R, Scancarello G, *et al.* Measurement of asbestos fibre concentrations in fluid of repeated bronchoalveolar lavages of exposed workers. *Ann Occup Hyg* 2007; 51: 495–500.

Copyright ©ERS 2017

From the authors:

We appreciate the opportunity to respond to the correspondence from X. Baur and colleagues regarding our article recently published in the *European Respiratory Journal* [1].

We added the subtitle “new insights into the chrysotile debate” to the title of our manuscript. In their accompanying editorial, NEMERY *et al.* [2] asked about the nature of the debate and stated among others that the term “chrysotile debate is not neutral”. The correspondence from X. Baur and colleagues corroborates this view. It demonstrates that there is indeed a heavy debate, mainly going on in Germany, but, as suggested by the large number of authors of this correspondence, it is also relevant in an international frame. Of note, people who have fought for the global ban of asbestos were co-authors of the article.

In our view, “debates” and discussions about scientific topics are common and needed. Multiple debates about chrysotile exist with changing aspects over time. The debate about the hazard of chrysotile inducing malignant and non-malignant pleural and lung diseases is over. Still open is the question about biopersistence of chrysotile in the lungs of affected workers.

In contrast to animal data [3], our manuscript shows high biopersistence of chrysotile fibres in human lungs [1], a result that perfectly fits to the well-known characteristics of asbestos: “imperishable”, “rot-proof”, “extremely resistant”. Our data explain the toxicity of the fibre and are a strong argument for a global ban of asbestos.

So what is the motivation of the authors of the correspondence? They fear that our findings may influence compensation claims. It was clearly the aim of our study to assess biopersistence of asbestos fibres, irrespective of any socio-economic implications which may even differ between countries. It is our experience that cases with relevant occupational asbestos exposure and a clinical picture of idiopathic pulmonary fibrosis without histological evidence of asbestosis are extremely rare. It is an international consensus; if one considers the Helsinki criteria as a clinical guideline, it should be clear that the histological diagnosis of asbestosis requires the existence of fibrosis *plus* the demonstration of asbestos

@ERSpublications

Intraindividual longitudinal data display: the asbestos fibre burden in living human lungs is stable over many years <http://ow.ly/Bl9S30gufCM>

Cite this article as: Feder IS, Tischoff I, Theile A, *et al.* Correspondence regarding the article “The asbestos fibre burden in human lungs: new insights into the chrysotile debate”. *Eur Respir J* 2017; 50: 1702204 [https://doi.org/10.1183/13993003.02204-2017].

bodies or uncoated asbestos fibres [4]. According to this consensus, asbestosis with few or no detectable asbestos fibres is speculative. Indeed, our manuscript makes the existence of such rare cases even more speculative. It should be noted that there are much higher numbers of patients with lung “fibrosis” who are diagnosed to have asbestosis by histology [5, 6]. Those patients are then recommended for compensation. Thus we do not see any disadvantages for patients with true asbestosis.

The criticism raised in the correspondence from X. Baur and colleagues is manifold and we cannot give a point to point reply within the space limit of the journal. Concerning methodological problems we refer to our method section and a more detailed view on some of the controversies published recently [7]. Here we would like to comment on the major points raised.

Regarding study design and the allegedly low number of subjects we are strongly convinced that this is not a real weakness of our study. As outlined in our manuscript, such data are difficult to obtain and therefore unique. It is a longitudinal observational study which is largely unaffected by the sample size. This was also recognised by NEMERY *et al.* [2] in their editorial where they concluded that “...it is not unreasonable to assume that similar results would probably be obtained in a study with more patients”. To support their conclusion, NEMERY *et al.* [2] added valuable information about seven patients who underwent repetitive bronchoalveolar lavage. This supplementary analysis confirms our data. There is an additional study by SARTORELLI *et al.* [8] of a further 22 patients, who too obtained similar results from repeated bronchoalveolar lavage fluid samples.

Concerning the chrysotile to amphibole ratio it is well known that the chrysotile fraction in the human lung is lower than at workplaces [9–12], as mentioned in our manuscript. This does not allow any statement about biopersistence of chrysotile in human lungs. It just is an observation, that chrysotile is deposited and accumulated in human lungs to a lower degree than amphibole. This is generally accepted and not challenged by our data.

Our data demonstrate that this accumulated asbestos fibre burden in the human lung is stable after exposure cessation and that this is true also for chrysotile. This was not known before as no one so far had analysed longitudinal intra-individual data from human lung tissue. As an explanation we discussed that the failure of chrysotile to accumulate in human lungs reflects events that occur early after exposure rather than long-term clearance mechanisms, as postulated by others earlier [13]. Thus our findings are not the opposite of but consistent with extensive literature [14].

The correspondence claims undisclosed financial conflicts of interest. We are convinced that putative conflicts of interest were disclosed properly as requested by the *European Respiratory Journal*.

In conclusion, we have shown for the first time that asbestos fibres including chrysotile are highly biopersistent in human lungs. We see these findings *per se* worth reporting.

Inke Sabine Feder¹, Iris Tischoff¹, Anja Theile¹, Inge Schmitz¹, Rolf Merget² and Andrea Tannapfel¹

¹Institute for Pathology of the Ruhr-University Bochum, Bochum, Germany. ²Institute for Prevention and Occupational Medicine of the German Social Accident Insurance, Institute of the Ruhr-University (IPA), Bochum, Germany.

Correspondence: Inke Sabine Feder, Georgius Agricola Stiftung Ruhr, Institut für Pathologie, Ruhr-Universität Bochum, am BG Universitätskliniken Bergmannsheil, Deutsches Mesotheliomregister, Bürkle-de-la-Camp Platz 1, 44789 Bochum, Germany. E-mail: Inke.feder@ruhr-uni-bochum.de

Received: Oct 26 2017 | Accepted: Nov 02 2017

Conflict of interest: Disclosures can be found alongside this article at erj.ersjournals.com

Acknowledgements: The Institute of Pathology gets third-party funds from the German Social Accident Insurance (DGUV) for research of the German Mesothelioma Register. Freedom to design, conduct, interpret, and publish research is not compromised by this.

References

- 1 Feder IS, Tischoff I, Theile A, *et al.* The asbestos fibre burden in human lungs: new insights into the chrysotile debate. *Eur Respir J* 2017; 49: 1602534.
- 2 Nemery B, Nuyts V, Nackaerts K. Quantifying asbestos in lung tissue: what debate? *Eur Respir J* 2017; 49: 1700861.
- 3 Bernstein DM, Cevalier J, Smith P. Comparison of Calidria chrysotile asbestos to pure tremolite: final results of the inhalation biopersistence and histopathology examination following shortterm exposure. *Inhal Toxicol* 2005; 17: 427–449.
- 4 Henderson D, Rantanen J, Barnhart S, *et al.* Asbestos, asbestosis and cancer: the Helsinki criteria for diagnosis and attribution. *Scand J Work Environ Health* 1997; 23: 311–316.
- 5 Everatt RP, Smolianskiene G, Tossavainen A, *et al.* Occupational asbestos exposure among respiratory cancer patients in Lithuania. *Am J Ind Med* 2007; 50: 455–463.

- 6 Attanoos RL, Alchami FS, Pooley FD, *et al.* Usual interstitial pneumonia in asbestos-exposed cohorts – concurrent idiopathic pulmonary fibrosis or atypical asbestosis? *Histopathology* 2016; 69: 492–498.
- 7 Merget R, Feder IS, Tannapfel A. Only One Lesson to be Learned Reply to X. Baur. *Pneumologie* 2016; 70: 405–412.
- 8 Sartorelli P, Romeo R, Scancarello G, *et al.* Measurement of asbestos fibre concentrations in fluid of repeated bronchoalveolar lavages of exposed workers. *Ann Occup Hyg* 2007; 51: 495–500.
- 9 Churg A. Deposition and clearance of chrysotile asbestos. *Ann Occup Hyg* 1994; 38: 625–633.
- 10 Friedrichs KH, Brockmann M, Fischer M, *et al.* Electron microscopy analysis of mineral fibers in human lung tissue. *Am J Ind Med* 1992; 22: 49–58.
- 11 Rasmuson JO, Roggli VL, Boelter FW, *et al.* Cumulative Retrospective Exposure Assessment (REA) as a predictor of amphibole asbestos lung burden: validation procedures and results for industrial hygiene and pathology estimates. *Inhal Toxicol* 2014; 26: 1–13.
- 12 Wagner JC, Pooley FD, Berry G, *et al.* A pathological and mineralogical study of asbestos-related deaths in the United Kingdom in 1977. *Ann Occup Hyg* 1982; 26: 423–431.
- 13 Churg A, DePaoli L. Clearance of chrysotile asbestos from human lung. *Exp Lung Res* 1988; 14: 567–574.
- 14 Wolff H, Vehmas T, Oksa P, *et al.* Asbestos, asbestosis, and cancer, the Helsinki criteria for diagnosis and attribution 2014: recommendations. *Scand J Work Environ Health* 2015; 41: 5–15.

Copyright ©ERS 2017