





Avoiding immortal time bias in observational studies

Samy Suissa^{1,2} and Pierre Ernst^{1,2}

Affiliations: ¹Center for Clinical Epidemiology, Lady Davis Institute - Jewish General Hospital, Montreal, QC, Canada. ²Dept of Epidemiology and Biostatistics and Dept of Medicine, McGill University, Montreal, QC, Canada.

Correspondence: Samy Suissa, Centre for Clinical Epidemiology, Jewish General Hospital, 3755 Cote Ste-Catherine, Montreal, Québec, Canada H3T 1E2. E-mail: samy.suissa@mcgill.ca

@ERSpublications

Raymakers and colleagues used the proper analysis that avoided immortal time bias, though the wide discrepancy in results with other studies remains; further studies are needed http://bit.ly/2Oobgqb

Cite this article as: Suissa S, Ernst P. Avoiding immortal time bias in observational studies. *Eur Respir J* 2020; 55: 2000138 [https://doi.org/10.1183/13993003.00138-2020].

This single-page version can be shared freely online.

From the authors:

We thank A.J.N. Raymakers and co-workers for their letter, which clarifies the method of data analysis used in their study of the association between inhaled corticosteroids (ICS) and lung cancer incidence in COPD [1]. Indeed, as shown by the computing code they provide, the data analysis did consider ICS use as a time-dependent exposure, thus avoiding immortal time bias.

Copyright ©ERS 2020