What constitutes a “clinically significant” bronchodilator response in children?

To the Editor:

We read with great interest the recent paper of DE JONG *et al.* [1] evaluating the contribution of a detailed history and a variety of diagnostic tests, including spirometry and bronchodilator tests, to diagnosing asthma in 111 children. In the methodology section, with regard to their definition of a “clinically significant” bronchodilator responsiveness (BDR), the authors only considered the forced expiratory volume in 1 s (FEV₁) and applied the following two thresholds: $\geq 10\%$ increase (no reference was cited) and $\geq 12\%$ increase (according to the National Institute for Health and Care Excellence (NICE) [2]). Their approach could be a source of confusion for at least three reasons.

First, how did DE JONG *et al.* [1] express the FEV₁ increase (percentage of the initial value (%_{init}) or that of the predicted reference value (%_{pred})?) and what about the increase in absolute change (Δ_A)? Although it is widely used in clinicians’ daily practice, $\Delta_{\%init}$ has three major disadvantages [3]. In fact, it is sex- and size-biased in assessing BDR. Moreover, patients with the lowest initial FEV₁ display the largest BDR. Finally, it is subject to change variability (*e.g.* in the absence of any true change, the difference between two successive measurements is negatively correlated with the initial value). Expressing BDR as a $\Delta_{\%pred}$ allows the comparison of patients whatever the age and height are. However, its use encounters two practical difficulties. On the one hand, it is not applicable in countries where spirometric norms are lacking, and on the other hand, it requires the use of software, which may not be available in low-income countries. ΔFEV_{1A} correlates with both age and height, and the changes are smaller in short children than in tall ones [3].

Secondly, why did DE JONG *et al.* [1] choose FEV₁? What about forced vital capacity (FVC) and/or peak expiratory flow (PEF)? The difference between the flow (*i.e.* FEV₁ and/or PEF) *versus* the volume (*i.e.* FVC) effects is not fully understood in the literature [4–8]. Although FEV₁ has several advantages [9] (*e.g.* well-standardised, highly reproducible), it presents some disadvantages. Most importantly, it is negatively effort-dependent with an overestimation in case of sub-maximal effort at the start of expiration [10]. The use of PEF, an effort-dependent datum [10], is still recommended by the Global Initiative for Asthma (GINA) [5]. Moreover, the PEF meter is proposed by the World Health Organization as an indispensable tool in the package of essential non-communicable disease interventions. In children, the use of FVC is still recommended by some scholarly societies [7, 8]. This reproducible datum [9] provides useful information about hyperinflation, air trapping and distal airways function [11]. Moreover, a decrease in air trapping, often associated with severe exacerbations, is related to an increase in the FVC [11].

Thirdly, which scholarly society recommended the use of a “FEV₁ increase $\geq 10\%$ ” to diagnose a “clinically significant” BDR? The retained aforementioned definition [1], first proposed by the European Respiratory Society (ERS) in 1993 [12], is no longer considered. A literature search found that some scholarly societies (GINA, American Thoracic Society (ATS), ERS, National Asthma Education and Prevention Program (NAEPP), South African Thoracic Society (SATS), NICE, Research Group on Advances in Paediatric Pneumology (GRAPP)) “failed” to adopt a clear consensus about what constitutes a “clinically significant” BDR in children [2, 4–8]. The following several definitions were proposed:

- GINA [4]: $\Delta FEV_{1\%pred} \geq 12\%$
- ATS/ERS [7]: $\Delta FEV_{1\%init} \geq 12\%$ and $\Delta FEV_{1A} \geq 0.2$ L and/or $\Delta FVC_{\%init} \geq 12\%$ and $\Delta FVC_A \geq 0.2$ L, regardless of age
- NAEPP [6]: $\Delta FEV_{1\%init} \geq 12\%$ and $\Delta FEV_{1A} > 0.2$ L

@ERSpublications

The worldwide disagreement with regard to what constitutes a “clinically significant” BDR has long been a source of confusion for clinicians and can hinder an appropriate asthma diagnosis and therefore “disturb” its management <http://bit.ly/2VONQhR>

Cite this article as: Guezguez F, Ben Saad H. What constitutes a “clinically significant” bronchodilator response in children? *Eur Respir J* 2020; 55: 2000207 [<https://doi.org/10.1183/13993003.00207-2020>].

- SATS [8]: $\Delta FEV_{1\%init} \geq 12\%$ or $\Delta FEV_{1A} > 0.2$ L and/or $\Delta FVC_{\%init} \geq 12\%$ or $\Delta FVC_A > 0.2$ L
- NICE [2]: $\Delta FEV_{1\%init} \geq 12\%$
- GRAPP [5]: $\Delta FEV_{1\%init} \geq 12\%$ or a $\Delta PEF_{\%init} \geq 20\%$

To conclude, this worldwide disagreement with regard to what constitutes a “clinically significant” BDR has long been a source of confusion for clinicians. This can hinder an appropriate asthma diagnosis and therefore “disturb” its management. It is high time scholarly societies standardised the interpretation of BDR in children.

Fatma Guezguez^{1,2} and Helmi Ben Saad^{1,2,3}

¹Laboratory of Physiology and Functional Explorations, Farhat HACHED Hospital, Sousse, Tunisia. ²Heart Failure (LR12SP09) Research Laboratory, Farhat HACHED Hospital, Sousse, Tunisia. ³Université de Sousse, Faculté de Médecine de Sousse, Laboratoire de Physiologie, Sousse, Tunisia.

Correspondence: Helmi Ben Saad, Laboratory of Physiology, Faculty of Medicine of Sousse, Street Mohamed Karoui, Sousse 4000, Tunisia. E-mail: helmi.bensaad@rns.tn

Received: 30 Jan 2020 | Accepted after revision: 27 Feb 2020

Acknowledgements: The authors wish to thank Samir Boukattay for his invaluable contribution in the improvement of the quality of the writing in the present article.

Conflict of interest: None declared.

References

- 1 de Jong CCM, Pedersen ESL, Mozun R, *et al.* Diagnosis of asthma in children: the contribution of a detailed history and test results. *Eur Respir J* 2019; 54: 1901326.
- 2 National Institute for Health and Care Excellence (NICE). Guideline Asthma Diagnosis and Monitoring. www.nice.org.uk/guidance/ng80/evidence/full-guideline-asthma-diagnosis-and-monitoring-pdf-4656178047 Date last updated: November 2017. Date last accessed: 24 February, 2020.
- 3 Waalkens HJ, Merkus PJ, van Essen-Zandvliet EE, *et al.* Assessment of bronchodilator response in children with asthma. Dutch CNSLD Study Group. *Eur Respir J* 1993; 6: 645–651.
- 4 Global Initiative for Asthma. Global Strategy for Asthma Management and Prevention. 2019. www.ginasthma.org Date last accessed: 24 February, 2020.
- 5 Groupe de Recherche sur les Avancées en Pneumo-Pédiatrie (GRAPP). Pulmonary function tests for the assessment and monitoring of asthma in children above 3 years of age. *Rev Mal Respir* 2003; 20: 638–643.
- 6 National Asthma Education and Prevention Program. Expert Panel Report 3 (EPR-3): guidelines for the diagnosis and management of asthma-summary report 2007. *J Allergy Clin Immunol* 2007; 120: Suppl. 5, S94–138.
- 7 Pellegrino R, Viegi G, Brusasco V, *et al.* Interpretative strategies for lung function tests. *Eur Respir J* 2005; 26: 948–968.
- 8 Masekela R, Gray D, Verwey C, *et al.* Paediatric spirometry guideline of the South African Thoracic Society: Part 1. *S Afr Med J* 2013; 103: 12 Suppl. 2, 1036–1041.
- 9 Otto W, Friedman M, Easton J. Intrasubject variability of forced expiratory flow rates in children with chronic, stable asthma. *Pediatr Asthma Allergy Immunol* 1987; 1: 153–157.
- 10 Stockley JA, Cooper BG. Breathing out: forced exhalation, airflow limitation. In: *Pulmonary Function Testing*. Springer, 2018; pp. 103–125.
- 11 Mahut B, Bokov P, Delclaux C. Abnormalities of plethysmographic lung volumes in asthmatic children. *Respir Med* 2010; 104: 966–971.
- 12 Standardized lung function testing. Official statement of the European respiratory society. *Eur Respir J* 1993; 6: Suppl. 16, 1–100.

Copyright ©ERS 2020

A clinically significant bronchodilator response in children: how should it be measured?

From the authors:

We thank F. Guezguez and H. Ben Saad for raising important questions on recommendations for assessing a bronchodilator response (BDR) in children. The authors summarise how recommended outcome measures and cut-offs for BDR in children vary between guidelines, and raise questions about our study [1].

To clarify: our study did not focus on bronchodilator reversibility testing in children or compare different outcome measures. Rather, we assessed the contribution of a detailed history and a variety of tests for diagnosing asthma in children aged 6–16 years referred to pulmonary outpatient clinics. We compared the diagnostic performance (sensitivity, specificity) of different commonly performed tests: skin-prick tests, measurement of exhaled nitric oxide fraction (F_{ENO}), spirometry, bronchodilator reversibility and bronchial provocation tests (BPT) by exercise, methacholine and mannitol against an asthma diagnosis from the paediatric pulmonologist. Thus, we used tests, outcomes and cut-offs commonly recommended and used in clinical practice. In addition, for examinations that provided a continuous (rather than a binary) output, such as forced expiratory volume in 1 s (FEV_1), BDR or F_{ENO} , we also assessed which cut-off distinguished best between those with and without asthma. We found that the combined sensitivity and specificity was highest for reported symptoms (frequent wheeze, night-time awakening due to wheeze, and wheeze triggered by pollen or pets). Among the tests, the area under the curve was highest for F_{ENO} and BPT by methacholine or exercise, and lower for spirometry, bronchodilator reversibility and skin-prick tests.

F. Guezguez and H. Ben Saad ask how we had assessed FEV_1 increase: as percentage of the initial value, as percentage of the predicted value, or as absolute increase. We calculated BDR as percentage increase of the initial FEV_1 (in mL) using the following formula: (FEV_1 post-bronchodilator – FEV_1 pre-bronchodilator)/ FEV_1 pre-bronchodilator. This is the most widely used method for calculating reversibility and recommended by most guidelines [2–4]. FEV_1 pre- and post-bronchodilator was measured in triplicate and American Thoracic Society/European Respiratory Society guidelines reproducibility criteria were applied [5]. Second, they indicated that BDR can also be calculated for forced vital capacity and peak expiratory flow. Although this is true, FEV_1 is the most widely recommended outcome, as it is less subject to cooperation and has higher reproducibility [2, 3].

Third, they wondered what evidence we used to base our cut-off levels on. The 12% cut-off is recommended in all recent guidelines, but derives from studies in adults, expert opinion, or studies that compared severe asthmatics with healthy children. Recent population-based studies have questioned this cut-off for the paediatric population [6, 7]. In our study of children seen for evaluation of possible asthma in paediatric outpatient clinics, a cut-off of 10% had the highest combined sensitivity and specificity. However, we want to stress that cut-off levels for diagnostic tests that produce an outcome on a continuous scale are artificial and an oversimplification. We lose information if we force a continuous measure into a binary one. There is a trade-off between sensitivity and specificity, and depending on the clinical question, higher or lower cut-offs can be preferable. In general, the further away a result is from the mean of the frequency distribution in healthy children, the more likely it is that it is pathological. There is no such thing as a “true” cut-off that distinguishes unequivocally between healthy and diseased. For clinical application, a cut-off is often helpful, but factors such as pre-test probability, place of the test in the diagnostic algorithm, and costs must also be considered.

In conclusion, we fully agree with F. Guezguez and H. Ben Saad that more research is needed to evaluate the usefulness of diagnostic tests for asthma in children. We also agree that cut-offs should be critically questioned and defined based on evidence from the patient population of interest (*i.e.* children suspected with asthma) and not based on studies in adults or on expert opinion.

@ERSpublications

More research is needed to evaluate the usefulness of diagnostic tests for asthma in children and cut-offs should be critically questioned and defined based on evidence from the patient population of interest <https://bit.ly/33xjLVW>

Cite this article as: de Jong CCM, Pedersen ESL, Mozun R, *et al.* A clinically significant bronchodilator response in children: how should it be measured? *Eur Respir J* 2020; 55: 2000636 [https://doi.org/10.1183/13993003.00636-2020].

Carmen C.M. de Jong ¹, Eva S.L. Pedersen¹, R. Mozun ¹, Myrofora Goutaki ^{1,2}, Daniel Trachsel³, Juerg Barben⁴ and Claudia E. Kuehni ^{1,2}

¹Institute of Social and Preventive Medicine, University of Bern, Bern, Switzerland. ²Paediatric Respiratory Medicine, Children's University Hospital of Bern, University of Bern, Bern, Switzerland. ³Paediatric Respiratory Medicine, Children's University Hospital of Basel, Basel, Switzerland. ⁴Paediatric Respiratory Medicine, Children's Hospital of Eastern Switzerland, St Gallen, Switzerland.

Correspondence: Claudia E. Kuehni, University of Bern, Institute of Social and Preventive Medicine, Mittelstrasse 43, CH-3012 Bern, Switzerland. E-mail: claudia.kuehni@ispm.unibe.ch

Received: 10 March 2020 | Accepted after revision: 13 March 2020

Conflict of interest: None declared.

References

- 1 de Jong CCM, Pedersen ESL, Mozun R, *et al.* Diagnosis of asthma in children: the contribution of a detailed history and test results. *Eur Respir J* 2019; 54: 1901326.
- 2 NICE. Guideline Asthma Diagnosis and Monitoring. www.nice.org.uk/guidance/ng80/resources/asthma-diagnosis-monitoring-and-chronic-asthma-management-pdf-1837687975621 Date last updated: November 2017, December 2019.
- 3 National Asthma Education and Prevention Program. Expert Panel Report 3 (EPR-3): Guidelines for the Diagnosis and Management of Asthma-Summary Report 2007. *J Allergy Clin Immunol* 2007; 120: Suppl. 5, S94–S138.
- 4 Pellegrino R, Decramer M, van Schayck CP, *et al.* Quality control of spirometry: a lesson from the BRONCUS trial. *Eur Respir J* 2005; 26: 1104–1109.
- 5 Quanjer PH, Stanojevic S, Cole TJ, *et al.* Multi-ethnic reference values for spirometry for the 3-95-yr age range: the global lung function 2012 equations. *Eur Respir J* 2012; 40: 1324–1343.
- 6 Murray C, Foden P, Lowe L, *et al.* Diagnosis of asthma in symptomatic children based on measures of lung function: an analysis of data from a population-based birth cohort study. *Lancet Child Adolesc Health* 2017; 1: 114–123.
- 7 Tse SM, Gold DR, Sordillo JE, *et al.* Diagnostic accuracy of the bronchodilator response in children. *J Allergy Clin Immunol* 2013; 132: 554–559.

Copyright ©ERS 2020