



Reply: BAL lymphocyte percentage is as good as the company it keeps

Nicola Adderley¹, Christopher J. Humphreys², Hayley Barnes ^{3,4}, Zahra A. Premji ⁵ and Kerri A. Johannson ^{1,6}

Affiliations: ¹Dept of Medicine, University of Calgary, Calgary, AB, Canada. ²Dept of Medicine, University of Toronto, Toronto, ON, Canada. ³Dept of Allergy, Immunology and Respiratory Medicine, Alfred Hospital, Melbourne, Australia. ⁴Dept of Medicine, University of California, San Francisco, San Francisco, CA, USA. ⁵Depts of Libraries and Cultural Resources, University of Calgary, Calgary, AB, Canada. ⁶Dept of Community Health Sciences, University of Calgary, Calgary, AB, Canada.

Correspondence: Kerri A. Johannson, 6th floor – Pulmonary Diagnostics, 4448 Front Street S.E., Calgary, AB, Canada T3M-1M4. E-mail: kerri.johannson@ahs.ca

@ERSpublications

No single value of BAL lymphocyte % in fibrotic hypersensitivity pneumonitis will be diagnostic for all, and it must be considered in the context of other clinical, radiological and histopathological features <https://bit.ly/3qSwUDo>

Cite this article as: Adderley N, Humphreys CJ, Barnes H, *et al.* Reply: BAL lymphocyte percentage is as good as the company it keeps. *Eur Respir J* 2021; 57: 2100092 [<https://doi.org/10.1183/13993003.00092-2021>].

This single-page version can be shared freely online.

Reply to O. Moran-Mendoza and M. Khalil: