



SHAREABLE PDF

# $P_{aO_2}/F_{IO_2}$ ratio: the mismeasure of oxygenation in COVID-19

Martin J. Tobin , Amal Jubran and Franco Laghi

**Affiliation:** Division of Pulmonary and Critical Care Medicine, Hines Veterans Affairs Hospital and Loyola University of Chicago Stritch School of Medicine, Hines, IL, USA.

**Correspondence:** Martin J. Tobin, Division of Pulmonary and Critical Care Medicine, Hines Veterans Affairs Hospital and Loyola University of Chicago Stritch School of Medicine, Hines, IL 60141, USA. E-mail: mtobin2@lumc.edu



@ERSpublications

$P_{aO_2}$  is the most precise measurement of patient oxygenation in COVID-19, but its accuracy is corrupted when communicated as  $P_{aO_2}/F_{IO_2}$  ratio <https://bit.ly/3tNipTJ>

**Cite this article as:** Tobin MJ, Jubran A, Laghi F.  $P_{aO_2}/F_{IO_2}$  ratio: the mismeasure of oxygenation in COVID-19. *Eur Respir J* 2021; 57: 2100274 [<https://doi.org/10.1183/13993003.00274-2021>].

This single-page version can be shared freely online.

*Reply to A. Tulaimat:*