



The role of cigarette smoking-derived pollutants in the risk of mortality in idiopathic pulmonary fibrosis

Hee-Young Yoon¹, Sun-Young Kim², Ok-Jin Kim² and Jin Woo Song ¹

¹Dept of Pulmonary and Critical Care Medicine, Asan Medical Center, University of Ulsan College of Medicine, Seoul, Republic of Korea. ²Dept of Cancer Control and Population Health, Graduate School of Cancer Science and Policy, National Cancer Center, Gyeonggi, Republic of Korea.

Jin Woo Song (jwsongasan@gmail.com)



Shareable abstract (@ERSpublications)

The effect of air pollutants on mortality risk in patients with IPF is independent of their smoking status <https://bit.ly/3rLUAvA>

Cite this article as: Yoon H-Y, Kim S-Y, Kim O-J, *et al.* The role of cigarette smoking-derived pollutants in the risk of mortality in idiopathic pulmonary fibrosis. *Eur Respir J* 2021; 58: 2101739 [DOI: 10.1183/13993003.01739-2021].

This single-page version can be shared freely online.

Copyright ©The authors 2021. For reproduction rights and permissions contact permissions@ersnet.org

Received: 19 June 2021
Accepted: 20 July 2021

Reply to T. Yasuma and co-workers: