**Supplementary Table S1.** Response to biological treatment of patients in Table 1

	Mepolizumab	Omalizumab
Blood eosinophils (cells/µl)		
Pre-treatment	$598.6 \pm 694.6$	$278.0 \pm 188.6$
On treatment	$65.7 \pm 31.0  (\mathbf{p=0.02})$	$200.0 \pm 185.1 \ (p=0.43)$
FeNO (ppb)		
Pre-treatment	$52.8 \pm 28.8$	$35.6 \pm 36.6$
On treatment	$49.1 \pm 35.0 \ (p=0.60)$	$22.9 \pm 12.1 \ (p=0.47)$
FEV1 (% predicted)		
Pre-treatment	$77.8 \pm 17.8$	$69.0 \pm 23.2$
On treatment	$71.3 \pm 13.3 \ (p=0.087)$	$69.0 \pm 22.2 (p=1)$
FEV1 postBD (% predicted)		
Pre-treatment	$81.9 \pm 19.2$ )	$74.4 \pm 22.7$
On treatment	$80.8 \pm 13.5 \ (p=0.78)$	$79.5 \pm 20.2 \ (p=0.45)$
Exacerbations/year		
Pre-treatment	$2.60 \pm 1.51$	$2.22 \pm 1.79$
On treatment	$0.13 \pm 0.28  (\mathbf{p=7.1x10^{-4}})$	$0.38 \pm 0.55 \ (\mathbf{p=0.015})$
ACQ		
Pre-treatment	$2.50 \pm 1.76$	$3.41 \pm 0.89$
On treatment	$1.57 \pm 0.68 \ (p=0.21)$	$2.91 \pm 1.22 (p=0.24)$

Results are expressed as mean  $\pm$  SD and p-values are given for comparisons of variables preand post-establishment of the treatment in each patients group (paired t-tests). ACQ: asthma control questionnaire; FeNO: exhaled nitric oxide; FEV1: forced expiratory volume in 1 s; postBD: post-bronchodilator treatment.