

Impact of climate change on ozone related mortality and morbidity in Europe

Hans Orru, *Department of Public Health and Clinical Medicine, Umea University, Umeå, Sweden; Department of Public Health, University of Tartu, Tartu, Estonia*

Camilla Andersson, *Swedish Meteorological and Hydrological Institute, Norrköping, Sweden*

Kristie L. Ebi, *ClimAdapt, Los Altos, California, USA*

Joakim Langner, *Swedish Meteorological and Hydrological Institute, Norrköping, Sweden*

Christofer Åström, *Department of Public Health and Clinical Medicine, Umea University, Umeå, Sweden*

Bertil Forsberg, *Umea University, Department of Public Health and Clinical Medicine, Occupational and Environmental Medicine, Umeå, Sweden*

Abstract

Ozone is a highly oxidative pollutant formed from precursors in the presence of sunlight, associated with respiratory morbidity and mortality. All else being equal, concentrations of ground-level ozone are expected to increase due to climate change.

Ozone-related health impacts under a changing climate are projected using emission scenarios, models and epidemiological data. European ozone concentrations are modelled with MATCH-RCA3 (50x50 km). Projections from two climate models, ECHAM4 and HadCM3, are applied, under greenhouse gas emission scenarios A2 and A1B respectively. We apply a European-wide exposure-response function to gridded population data and country-specific baseline mortality and morbidity.

Comparing the current situation (1990–2009) with the baseline period (1961–1990), the largest increase in ozone-associated mortality and morbidity due to climate change (4–5%) have occurred in Belgium, Ireland, Netherlands and UK. Comparing the baseline period and the future periods (2021–2050 and 2041–2060), much larger increase in ozone-related mortality and morbidity are projected for Belgium, France, Spain and Portugal with the impact being stronger using the climate projection from ECHAM4 (A2). However, in Nordic and Baltic countries the same magnitude of decrease is projected.

The current study suggests that projected effects of climate change on ozone concentrations could differentially influence mortality and morbidity across Europe.

Introduction

Ozone is one of the most important air pollutants. Ozone is formed in photochemical reactions, with concentrations affected by weather and the supply of chemical precursors, including nitrogen oxides (NO_x), volatile organic compounds (VOCs), methane (CH_4) and carbon monoxide (CO). Climate change can affect ozone concentrations and thus influence respiratory health [1] through a number of processes, including chemical production, dilution and deposition of ozone that are regulated by temperature, cloud cover, humidity, wind and precipitation [2-4]. Ozone also is formed from reactions that include natural biogenic emissions; these reactions also depend on temperature and solar radiation, and thereby cloud cover. Although there is high confidence in projected changing temperatures [5], changes in other meteorological parameters, such as precipitation and cloud cover are more uncertain. There is also large uncertainty in how natural vegetation will respond to climate change.

Extreme weather events such as heat waves (observed in Europe in 2003 and 2006) can further increase ground-level ozone concentrations [6, 7]. During the 2003 heat wave, the relative contributions of temperature to adverse health effects in France ranged from 97.5% in Bordeaux to 14.7% in Toulouse, where air pollution and ozone singly and jointly contributed to mortality [6]. Chemical transport models project widespread temperature-related summertime increases of ground-level ozone in polluted regions of Europe, North America and Asia. The sensitivity of ground-level ozone to climate change is particularly high in urban areas, reflecting the concentration of precursors for ozone formation. The frequency of stagnation episodes is projected to increase over northern mid-latitude continents and the ventilation is projected to decrease in Europe, eastern North America and East Asia [8].

Climate change may lead to higher biogenic VOC emissions [9]. Warmer temperatures may lead to increasing soil microbial activity that may cause an increase in NO_x emissions and a consequent increase in ozone amounts [10]. Methane emissions promote tropospheric ozone formation and

global climate change [11]. Climate change and increasing temperature could also affect the risk of wildfires [12]. Due to the emissions of precursors such as CO, the 2007 California fires, for instance, caused a three-fold exceedance of the limit values for ozone in the region [13].

Epidemiological studies have shown a broad range of effects of ground-level ozone on health, leading to excess daily mortality and morbidity. Significant negative health effects have been demonstrated for different causes, mainly for respiratory and (to a lesser extent) cardiovascular diseases [14]. The majority of epidemiological studies concentrated on acute health consequences (due to short-term very high concentrations of ozone). For instance, the association between acute exposures of ozone and short-term mortality was shown in several studies [15-18]. There are several large multi-city studies relating the numbers of hospital admissions for respiratory diseases [19] and COPD [20] to ambient ozone levels. Exposure to ozone has been shown to increase the likelihood of wheeze, chest tightness and asthma [21]. The other main effects include emergency department visits for asthma, respiratory tract infections, exacerbation of existing airway diseases [22] as well as the decline of lung function [23].

Epidemiological evidence of the chronic effects of ground-level ozone is less conclusive. Chronic exposures could induce significant changes in airways and systemic inflammation. A recent study by Jerrett et al. [24] showed significant increases in the risk of death from respiratory causes in association with an increase in ozone concentration. However, the risk of death from cardiovascular causes was not any more significant when the concentration of PM_{2.5} was taken into account. The summer ozone average was associated with survival in four cohorts of persons with chronic conditions that might predispose to ozone effects, such as congestive heart failure [25]. Other studies indicate some cardiovascular effect that may be due to systemic inflammation responses and alterations of the cardiac rhythm. Chuang et al. [26] found increases in inflammatory markers in the blood and a decrease in heart rate variability associated with elevated levels of ozone lagged 1–3

days in young and healthy adults. In some Asian studies, ozone exposure has been associated with increasing mortality due to stroke [27]. The mortality risk exhibited a seasonal variation, being more intense during periods of warmer weather.

A few studies have projected the effects of climate change on ground-level ozone concentrations and the subsequent effects on public health [28-36]. Four of these studies were published since the review by Ebi and McGregor [37]; three focused on the United States and one on the UK. Although using different greenhouse gas emission scenarios and models, the US studies by Chang et al. [33] and Tagaris et al. [34] concluded that climate change is projected to increase ozone concentrations by the 2050s resulting in increased ozone-related mortality, with more premature deaths in the southern states. A third study focused on childhood asthma emergency department visits in the New York City metropolitan area [36], using the same ozone projections as Knowlton et al. [28]. The UK study compared changes in ground-level ozone assuming A2 emissions (+15%) to changes assuming current legislation and air quality standards in the future (+6%) to changes assuming maximum feasible reductions (-5 %) [35]. Ozone concentrations in ten world regions, assuming A2 emissions and current baseline mortality rates, are projected to increase by 9.2 ppb(v) as a global population-weighted daily 8-hour maximum between 2000 and 2030 [32]. For Europe the population weighted mean is projected to increase by 4.7 ppb(v).

Impact assessments have estimated the health impacts of historic vs current ground-level ozone concentrations. Anenberg et al. [38] estimated anthropogenic ground-level ozone concentrations and related health impacts, and found that assuming a low concentration threshold (25 ppb(v)) has a large impact on the estimated impact on mortality. Moreover, they found that comparing 2000 to 1860, ozone concentrations increased by a factor two or three, depending on the region of world. For Europe, the modelled annual average concentration increased from 18.26 to 48.92 ppb(v), with

the current concentrations in Europe expected to cause 0.041 ± 0.021 or 0.023 ± 0.017 million excess respiratory mortalities annually, depending on the threshold assumed.

The current study assesses the impacts of climate change on ozone-related mortality and morbidity in Europe over wider time periods than often used. Further, it illustrates the impact of applied greenhouse gas emission scenarios and global climate models on projected health impacts.

Material and methods

Data on ozone exposure, baseline mortality, morbidity, and population were generated for the health impact assessment.

European ozone concentrations were modelled at a grid size of 50x50 km using the chemistry-transport model MATCH [4, 39] that models about 130 chemical reactions between 70 chemical components, including thermal and photo chemical reactions, wet chemistry and secondary particle formation. Species at the lateral and top boundaries of MATCH were kept at levels representative for year 2000 throughout the simulated years. MATCH simulates biogenic emissions of isoprene based on hourly temperature and solar radiation. Biogenic emissions of isoprene also depended on the geographical distribution of vegetation types, which was held constant throughout the simulations. This is a simplification, but reliable projections of vegetation linked to climate projections are unavailable. Anthropogenic precursor emissions of nitrogen oxides, sulphur oxides, carbon monoxide, non-methane volatile organic compounds and ammonia from EMEP (<http://www.ceip.at>) for the year 2000 were used as input to MATCH. These anthropogenic emissions were emitted with seasonal, weekly and daily variation at 2000 levels for all simulated years. MATCH uses meteorology produced by the regional climate model RCA3 [40, 41]. Projections from two global climate models, ECHAM4 and HadCM3 under greenhouse gas emission scenarios A2 and A1B respectively, were used as input to RCA3, which in turn produced a dynamical downscaling of the climate over Europe at a higher resolution. With ECHAM4 (A2) two previously presented [42, 43] periods were compared: the baseline period as 1961–1990 and future as 2021–2050 (MATCH-RCA3-ECHAM4). With HadCM3 (A1B) [44, 45] two additional periods were included: the current situation as 1990–2009 and further future as 2041–2060 (MATCH-RCA3-HadCM3). These model parameters have been extensively evaluated within the given references.

Different emission scenarios of greenhouse gases were used in the two climate projections downscaled by RCA3. However, the European average temperature changes are quite similar for the two simulations between 1961–1990 and 2021–2050 using the HadCM3 (A1B) and ECHAM4 (A2) projection, 2.1 and 1.9°C respectively. Warming is somewhat stronger in HadCM3 because HadCM3 has a stronger response to greenhouse gas forcing. The resemblance is partly explained by the limited difference in anthropogenic greenhouse gas emissions for the period 2021–2050 between the A1B and A2 scenarios. Projected CO₂ concentrations for 2010 in A1B and A2 are close to current (2010) observed CO₂ concentrations. The projected changes in CO₂ emissions are larger towards the end of the 21st century with a bigger difference between A1B and A2 that results in larger differences in the simulated temperature change between 1961–1990 and 2071–2100, 3.8 and 4.3°C using HadCM3 and ECHAM4 respectively. The two climate projections have the second and fourth highest average temperature change for the period 2040–2070 in an ensemble of 13 different projections downscaled from different GCM runs by RCA3 over Europe using the A1B or A2 scenario [46]. Even though differences in mean temperature change until 2050 are not large there are other differences between the two projections on the regional scale within Europe. In particular the reduction in precipitation in summer is larger in southwestern Europe in ECHAM4 (A2) and consequently cloud cover and soil moisture. These factors are important for ozone production and dry deposition to vegetation.

Ozone exposure measured by SOMO₃₅ (the Sum of Ozone daily 8 hour max Means Over 35 ppb(v) in the calendar year, expressed in µgm⁻³ times hours) values were calculated with a geographical resolution of 50 km. Hence, the health effects were calculated for levels above 35 ppb(v) (70 µgm⁻³). Often the cut-off value of 70 µgm⁻³ is used in risk assessments, as a statistically significant increase in mortality risk estimates has been observed at daily ozone concentrations above 50–70 µgm⁻³ [47, 48]. As a sensitivity analysis we also used cut-off values of SOMO₅₀ and SOMO₂₅. To see the seasonal

impacts, the SOMO₃₅ values and its expected health impacts were calculated separately for summer and winter. The population average exposures were calculated from country impact estimates.

The data on mortality and hospitalization were obtained from WHO European Health for All Database (<http://data.euro.who.int/hfad>). The crude non-standardized all-cause mortality and respiratory hospitalizations rates were used. Average rates for 2000–2005 were calculated for all countries and applied to all grids within the country. A shape of the national borders was used to determine to which country each grid cell belongs.

The gridded population data for Europe in 2000 were taken from the HYDE theme within the Netherlands Environmental Assessment Agency [49].

For the calculation of mortality and morbidity cases (ΔY) in absolute and relative numbers the following equation was used:

$$\Delta Y = (Y_0 \times pop) \times (e^{\beta \times X} - 1),$$

where Y_0 is the baseline mortality or morbidity rate; pop the number of exposed persons; β the exposure-response function (relative risk) and X the estimated excess exposure.

In order to describe the effects of ozone on mortality, the WHO meta-analysis relative risk (RR) 1.003 per 10 $\mu\text{g m}^{-3}$ increase in the maximum daily 8-hour average ozone concentration (95% CI 1.001–1.004) was used as the exposure-response coefficient (ERC) [14]. The same meta-analysis estimated the ERC of respiratory hospitalisations for adults (15–64) as 1.001 (95% CI 0.991–1.012) and for older adults (65+) as 1.005 (95% CI 0.998–1.012) per 10 $\mu\text{g m}^{-3}$ increase in the maximum daily 8-hour average ozone concentration. However, because for most countries only the hospitalization rates for all ages were available, the alternative ERC (1.003) was used. This combined COMEAP [50] unofficial meta-analysis results 1.003 (95% CI 1.000–1.007) and ERC based on proportion of respiratory

hospital admissions in different age groups (15–64 and 65+).

Results

Climate change will affect ground-level ozone concentrations (Figure 1), through a number of processes, as described in the Introduction. Changing ozone concentrations will then affect mortality and respiratory hospital admissions (Table 1). The baseline (recent) and current exposure to ground-level ozone above the applied threshold ($SOMO_{35}$) is associated with a large number of premature deaths (approx. 26–28,000 per year) and hospitalizations in Europe (Table 1, 2). The largest impact is in southern European countries with large populations, such as Italy, Spain and France.

When the current situation (1990–2009) is compared with the baseline period (1961–1990) using the ozone estimates based on MATCH-RCA3-HadCM3, the largest climate change driven relative increase in ozone related mortality and hospitalizations is modelled to have occurred in Ireland, UK, the Netherlands and Belgium (Table 1); an increase up to 5% is estimated. A decrease is estimated for the northernmost countries, with largest decrease, by 5%, in Finland. In absolute numbers, the model suggests 647 more deaths, and 867 excess hospitalizations per year in Europe with the largest numbers in Italy: 100 and 117 respectively.

If we compare the baseline period (1961–1990) with the future (2021–2050), the difference is even more dramatic for several countries (Table 1). The increase in ozone related cases is projected to be largest in Belgium, France, Spain and Portugal (10–14%). However, in most Nordic and Baltic countries, there is a projected decrease in ozone-related mortality of the same magnitude. The change is stronger if we compare the further future (2041–2060) with the baseline period (1961–1990) as simulated using HadCM3 (A1B). The projected impacts are larger using the ECHAM4 (A2) projection, up to 34% increase in Belgium, due to a stronger reduction in summer precipitation in this region and corresponding reductions in cloudiness and soil moisture leading to higher ozone concentrations.

We also investigated how large climate change induced changes in ground-level ozone concentration could be on the total all-cause mortality and on the total rate of respiratory hospitalizations. Because the exposure-response coefficients for total mortality and respiratory hospitalization are of the same magnitude (0.3% per 10 $\mu\text{g m}^{-3}$ increase in the maximum daily 8-hour average ozone concentration), the impact would be the same in relative numbers but different in total numbers. The relative increase of all-cause mortality and respiratory hospitalizations in some areas of Spain, Italy and Portugal are expected to be more than 0.2% in the future compared to base-line period rates (Figure 2). The effect is again larger using the ECHAM4 (A2) projection. Comparing the current period (1990–2009) with the baseline (1961–1990) and the further future (2041–2060) with baseline (1961–1990) using the HadCM3 (A1B) projection suggests that the majority of the impacts in the highest risk areas will happen in the future and only a smaller part has already occurred (Figure 2). However, there is variability on decadal scale in the models as in reality, but the modelled and real world decadal variability do not necessarily correlate; in fact, the decadal variabilities may even be anti-correlated. This means that change over time periods differing by one or a few decades simulated by climate models are not necessarily comparable to reality.

There are regional differences in the climate change projections (1961–1990 vs 2021–2050), depending on which global climate model (ECHAM4 or HadCM3) and CO₂ emission scenario (A2 or A1B) were used as input to RCA3. For most countries, MATCH-RCA3-ECHAM4 (under the A2 scenario) produced larger increases; however, for some countries (e.g. Greece, Bulgaria), the increase is of the same magnitude in the MATCH-RCA3-HadCM3 scenario (under the A1B scenario). There are also differences in SOMO₃₅ values in the base-line period (1961–1990) modelled with MATCH based on the regional downscaling of the two global climate models, despite similar greenhouse gas emissions in the base-line period in the two realisations (Figure 1). This is due to differences in the model realisations of the current climate [51]. For some countries, e.g. Belgium, Netherlands and UK, the MATCH-RCA3-HadCM3 scenario results in more than 25% higher concentrations; whereas for

Southern European countries e.g. Spain and Portugal, the SOMO₃₅ values were more than 10% lower compared to MATCH-RCA3-ECHAM4.

Table 1. Projected annual counts of premature mortality and respiratory hospitalizations due to ozone >SOMO₃₅ in the EU27 countries, Norway and Switzerland. Estimates build on modelled ground-level ozone concentrations based on chemistry-transport calculations using regional climate downscaling (with RCA3) of two different global climate models (ECHAM4 and HadCM3) with two CO₂ emission scenarios (A2 and A1B) in different time periods

	Mortality (absolute number of cases)						Hospitalizations (absolute number of cases)					
	MATCH-RCA3-ECHAM4 (A2)		MATCH-RCA3-HadCM3 (A1B)				MATCH-RCA3-ECHAM4 (A2)		MATCH-RCA3-HadCM3 (A1B)			
	1961–1990	2021–2050	1961–1990	1990–2009	2021–2050	2041–2060	1961–1990	2021–2050	1961–1990	1990–2009	2021–2050	2041–2060
Austria	485	522	533	539	559	558	945	1016	1037	1050	1088	1087
Belgium	381	512	529	551	602	626	537	722	745	777	848	882
Bulgaria	720	744	672	693	716	722	1266	1308	1180	1217	1259	1268
Cyprus	54	54	51	51	52	52	55	54	51	51	52	53
Czech	600	650	664	678	704	704	828	897	917	935	971	971
Denmark	238	255	290	291	295	292	374	400	455	457	463	459
Estonia	55	51	61	60	58	54	86	80	96	94	91	85
Finland	123	113	145	138	132	126	273	250	322	306	294	280
France	2659	3320	3123	3178	3488	3594	3852	4809	4524	4604	5052	5206
Germany	2167	2562	2675	2723	2903	2945	2771	3276	3421	3482	3712	3766
Greece	1007	1052	956	984	1020	1045	1481	1546	1405	1447	1499	1536
Hungary	791	853	802	830	874	866	1226	1322	1244	1287	1355	1343
Ireland	62	72	79	83	78	75	121	139	152	161	151	144
Italy	5737	6491	6003	6103	6553	6630	6712	7594	7024	7141	7667	7757

Latvia	100	94	108	108	106	98	162	152	175	174	171	159
Lithuania	123	113	129	130	127	119	278	255	291	292	287	268
Luxembourg	19	24	24	24	26	27	37	46	46	47	51	51
Malta	35	37	38	38	39	40	26	27	28	28	29	29
Netherlands	496	640	696	729	776	791	375	483	526	551	586	598
Norway	121	115	150	148	137	132	180	172	223	220	204	197
Poland	1771	1825	1939	1990	2028	1957	2617	2697	2865	2940	2997	2892
Portugal	819	972	726	744	823	848	668	792	592	606	671	691
Romania	1481	1500	1397	1435	1486	1473	3849	3898	3632	3730	3862	3830
Slovakia	302	315	312	317	328	325	436	455	450	458	474	469
Slovenia	127	138	132	134	139	139	163	178	171	172	179	179
Spain	3236	3730	2887	2975	3324	3425	3902	4499	3482	3587	4008	4130
Sweden	303	295	360	355	347	337	285	278	339	334	327	317
Switzerland	412	456	485	488	502	507	453	501	534	536	552	558
UK	1489	1954	2045	2143	2191	2215	1640	2152	2252	2361	2413	2439
Total	25915	29458	28012	28658	30414	30723	35596	39998	38178	39045	41313	41645

Table 2. Total annual counts of premature mortality and respiratory hospitalizations in Europe and projected change (%) in future due to ozone exposure change using different seasons and cut-off values

	MATCH-RCA3-ECHAM4 (A2)		MATCH-RCA3-HadCM3 (A1B)			
	1961–1990 (number of cases)	2021–2050 vs 1961–1990 (%)	1961–1990 (number of cases)	1990–2009 vs 1961–1990 (%)	2021–2050 vs 1961–1990 (%)	2041–2060 vs 1961–1990 (%)
	Mortality/hospitalizations					

SOMO ₃₅ annual	25,915/ 35,596	13.7/12.4	28,012/ 38,178	2.3/2.3	8.6/8.2	9.7/9.1
SOMO ₃₅ winter	4,550/ 6,348	9.1/7.2	4,553/ 6,457	-3.2/-4.0	0.6/-1.2	-0.9/-2.9
SOMO ₃₅ summer	21,342/ 29,217	14.6/13.5	23,434/ 31,688	3.4/3.5	10.1/10.1	11.7/11.5
SOMO ₂₅	47,389/ 65,603	8.2/7.3	49,558/ 68,334	0.9/0.8	4.6/4.2	5.1/4.6
SOMO ₅₀	7,108/ 9,275	35.3/34.3	8,289/ 10,705	7.4/7.9	23.8/24.5	27.8/28.1

Discussion

Ground-level ozone is a crucial public health issue. The ambient ozone concentrations in the northern hemisphere generally reach concentrations known to be harmful to health during large parts of every year [4]. The respiratory system has antioxidant defences that can mitigate to some extent the impacts of ozone exposure. Age, pre-existing diseases, social and economic status, habits, genetics, and other factors affect individual thresholds for noticeable impacts. This variability obscures the determination of a clear no-effect exposure concentration at the population level. Individuals with underlying lung or airways diseases are at higher risk of adverse impacts, including wheeze, chest tightness, cough, and asthma attacks. People with asthma and allergic rhinitis are somewhat more susceptible to transient alterations in respiratory function caused by acute exposure to ozone. Lung function decrease is more consistently observed in asthmatic children, especially those with low birth weight. At high concentrations, ozone enhances airway responsiveness also in healthy individuals. In the current assessment, health indicators are premature mortality and respiratory hospitalizations related to short-term exposure because there are available meta-coefficients for these outcomes. The actual health effects could be more severe, including effects on survival, but also affect a larger proportion of the population in terms of respiratory illness.

Exposure-response functions support the use of a cumulative SOMO metric to estimate health impacts (12, 44). The cut-off value used (35 ppb(v)) is below the WHO air quality guideline for ozone of maximum daily 8-hour average $100 \mu\text{g m}^{-3}$ [52] and the EU air quality directive 2008/50/EC of maximum daily 8-hour average $120 \mu\text{g m}^{-3}$, not to be exceeded on more than 25 days per calendar year [53]. The same cut-off value has been used in other health impact assessments [54-56]. As epidemiological studies have shown associations also at lower concentrations (e.g. [48, 57]), by using this cut-off value, the total number of cases attributed to ozone are likely underestimated in all scenarios. Using SOMO_{25} values as a cut-off would approximately double the number of attributed cases, but decrease the projected relative increase (Table 2). However, using the higher cut-off of

SOMO₅₀ would significantly decrease the number of cases, but increase the relative changes (Table 2), since the largest increase appeared among high ozone days (maximum daily 8-hour average more than 100 µg^m⁻³). Most of the projected increase in SOMO₃₅ is during summer, April–September (Table 2).

Our results on the numbers of current ozone induced mortalities for Europe with different cut-off values (Table 2) are very similar to the results from Anenberg et al. [38], even though they used a “long-term” exposure-response relationships [24] while we use a more established “short-term” exposure-response relationships [14]. However, because long-term exposure may also induce chronic disease (such as asthma) and possibly effects on survival, the size and kind of ozone-related health impacts could be more significant than we indicate. Recently, ozone exposure in Sweden and other countries has also been associated with pre-term birth, a risk factor for asthma [58].

We focused on the impacts of climate change-related alterations in ground-level ozone concentrations, holding other factors constant. Our results show that climate change could impact health in the future through higher ozone concentrations in several countries. An increase of up to 13.7% ozone-related mortality in Europe (with SOMO₃₅), that translates into 0.2% in all-cause total mortality, and respiratory hospitalizations would affect the public health sector through increased health-care costs. In some countries (e.g. Northern Europe) reductions of ozone induced mortality and hospitalizations are expected in the future due to climate change. Many processes contribute to the decrease in tropospheric ozone including increasing chemical destruction due to more water vapour, decreasing natural isoprene emissions, increased dry deposition and changing pollution transport patterns [43].

Comparing our current analysis with the Clean Air for Europe (CAFE) programme assessment for 2000 and 2020 [55], we found somewhat larger impacts for most countries. In the CAFE assessment,

two ozone scenarios for 2020 were projected based on current legislation and maximum technically feasible reductions that forecast decreased concentrations with no change in climate. In the current analysis, the possible decrease of ozone precursor emissions and decrease of population in Europe were not taken into account because the goal of the study was to show only the effects of climate change. Our estimates for UK compare well with estimates by the UK Department of Health [54].

Climate change will also affect other air pollutants, such as particulate matter (PM), though the atmospheric chemistry differs. PM has a stronger (inverse) relation than ozone to boundary layer height, which is related to wind speed and convective processes. Temperature dependency varies by particle species, with a positive association with sulphates and a possibly negative association with organic species. PM concentrations decrease with increasing precipitation [8]. The effect of climate change on PM is highly uncertain; even the sign differs between studies [8]. Both climate realisations used here simulate decreased precipitation in the south and increased in the north, increased temperature in all of Europe and decreasing wind speed in southern Europe. This implies that different particulate components will experience future changes of different magnitude and spatial variation.

Several methodological issues may have affected the results. For the time periods studied, the choice of greenhouse gas emission scenario is not crucially important because the differences in emissions between the scenarios are small before 2050. A more important factor is the global climate model used. The downscaling of the two different global climate models gave somewhat different results in different regions of Europe. In most countries, using the HadCM3 global model resulted in higher ground-level ozone baseline values (1961–1990) compared to ECHAM4. This indicates that in assessing local effects, the choice of global model is important. Also, the climatic variables (such as temperature, humidity, etc.) could affect respiratory hospitalization rates and thus the impacts of

ozone. As the rates are low in mild climate areas such as Southern-Europe compared to Northern-Europe, the absolute increase in impacts might be higher than suggested in this study.

These projections can be used in conjunction with projections of changes in emissions under different proposed regulations to understand the magnitude and extent of impacts under a higher temperature future. These projections also can be used by ministries of health and public health organizations to begin planning how to improve current programmes to ensure that vulnerable populations are protected from projected increases in ground-level ozone concentrations in a changing climate.

Conclusions

The projected effects of climate change on ground-level ozone concentrations could differentially influence mortality and morbidity across Europe. There would be an increase in ozone-related mortality in Southern and Central Europe and a slight decrease in Northern Europe. Compared to the baseline period (1961–1990), few climate-related ozone impacts appeared in the last two decades (1990–2009), with more projected in the future (2021–2050 and 2041–2060). The HadCM3 global model projected somewhat higher ozone concentrations for the baseline compared to using ECHAM4 in many countries. ECHAM4 gave generally larger health impacts for 2021–2050.

Acknowledgements

This work was supported by the EU-funded Climate-Trap project (contract EAHC 20081108)) and by the Swedish Environmental Protection Agency through the research programme CLEO – Climate Change and Environmental Objectives. We would also like to acknowledge Estonia's Ministry of Education for providing resources to H. Orru with the grant SF0180060s09.

References

1. Ayres JG, Forsberg B, Annesi-Maesano I, Dey R, Ebi KL, Helms PJ, Medina-Ramon M, Windt M, Forastiere F. Climate change and respiratory disease: European Respiratory Society position statement. *Eur Respir J* 2009; 34(2): 295-302.
2. US EPA. Assessment of the impacts of global change on regional U.S. air quality: A synthesis of climate change impacts on ground-level ozone. Washington, DC: U. S. E. P. Agency; 2009.
3. Andersson C, Engardt M. European ozone in a future climate: Importance of changes in dry deposition and isoprene emissions. *J Geophys Res* 2010; 115(D2): D02303.
4. Andersson C, Langner J, Bergström R. Interannual variation and trends in air pollution over Europe due to climate variability during 1958–2001 simulated with a regional CTM coupled to the ERA40 reanalysis. *Tellus B* 2007; 59(1): 77-98.
5. IPCC. Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2007.
6. Filleul L, Cassadou S, Medina S, Fabres P, Lefranc A, Eilstein D, Le Tertre A, Pascal L, Chardon B, Blanchard M, Declercq C, Jusot JF, Prouvost H, Ledrans M. The relation between temperature, ozone, and mortality in nine French cities during the heat wave of 2003. *Environ Health Perspect* 2006; 114(9): 1344-1347.
7. Leonardi G, Cooper D, Walton H, Catchpole M, Pebody R, Smith G, Bickler G, Bryden C, Stedman J, Kent A, Grice S. Rapid evaluation of 2006 heat wave: epidemiological aspects: Health Protection Agency; 2006.
8. Jacob DJ, Winner DA. Effect of climate change on air quality. *Atmospheric Environment* 2009; 43(1): 51-63.
9. Steiner A, Tonse S, Cohen R, Goldstein A, Harley R. The Effects of Climate Change on Biogenic VOCs and Regional Air Quality in California. American Geophysical Union, Fall Meeting, 2005; pp. A32A-02.
10. Pfeiffer M, Kaplan JO. Response of terrestrial N₂O and NO_x emissions to abrupt climate change. *IOP Conference Series: Earth and Environmental Science*: 9(1): 012001.
11. West JJ, Fiore AM, Horowitz LW, Mauzerall DL. Global health benefits of mitigating ozone pollution with methane emission controls. *Proceedings of the National Academy of Sciences of the United States of America* 2006; 103(11): 3988-3993.
12. Westerling AL, Hidalgo HG, Cayan DR, Swetnam TW. Warming and Earlier Spring Increase Western U.S. Forest Wildfire Activity. *Science* 2006; 313(5789): 940-943.
13. Pfister GG, Wiedinmyer C, Emmons LK. Impacts of the fall 2007 California wildfires on surface ozone: Integrating local observations with global model simulations. *Geophysical Research Letters* 2008; 35(19): CiteID L19814.
14. Anderson H, Atkinson R, Peacock J, Marston L, Konstantinou K. Meta-analysis of time-series studies and panel studies of particulate matter (PM) and ozone (O₃). Report of a WHO task group. Copenhagen: WHO Regional Office for Europe; 2004.
15. Bell ML, Dominici F, Samet JM. A meta-analysis of time-series studies of ozone and mortality with comparison to the national morbidity, mortality, and air pollution study. *Epidemiology* 2005; 16(4): 436-445.
16. Gryparis A, Forsberg B, Katsouyanni K, Analitis A, Touloumi G, Schwartz J, Samoli E, Medina S, Anderson HR, Niciu EM, Wichmann HE, Kriz B, Kosnik M, Skorkovsky J, Vonk JM, Dortbudak Z. Acute effects of ozone on mortality from the "air pollution and health: a European approach" project. *Am J Respir Crit Care Med* 2004; 170(10): 1080-1087.
17. Ito K, De Leon SF, Lippmann M. Associations between ozone and daily mortality: analysis and meta-analysis. *Epidemiology* 2005; 16(4): 446-457.

18. Levy JI, Chemerynski SM, Sarnat JA. Ozone exposure and mortality: an empiric bayes metaregression analysis. *Epidemiology* 2005: 16(4): 458-468.
19. Burnett RT, Brook JR, Yung WT, Dales RE, Krewski D. Association between ozone and hospitalization for respiratory diseases in 16 Canadian cities. *Environ Res* 1997: 72(1): 24-31.
20. Anderson HR, Spix C, Medina S, Schouten JP, Castellsague J, Rossi G, Zmirou D, Touloumi G, Wojtyniak B, Ponka A, Bacharova L, Schwartz J, Katsouyanni K. Air pollution and daily admissions for chronic obstructive pulmonary disease in 6 European cities: results from the APHEA project. *Eur Respir J* 1997: 10(5): 1064-1071.
21. Mortimer KM, Neas LM, Dockery DW, Redline S, Tager IB. The effect of air pollution on inner-city children with asthma. *Eur Respir J* 2002: 19(4): 699-705.
22. Amann M, Bertok I, Cofala J, Gyarmas F, Heyes C, Klimont Z, Schöpp W, Winiwarter W. Baseline scenarios for the Clean Air for Europe (CAFE) programme. Final report. Laxenburg: International Institute for Applied Systems Analysis; 2005.
23. Peters JM, Avol E, Gauderman WJ, Linn WS, Navidi W, London SJ, Margolis H, Rappaport E, Vora H, Gong H, Jr., Thomas DC. A study of twelve Southern California communities with differing levels and types of air pollution. II. Effects on pulmonary function. *Am J Respir Crit Care Med* 1999: 159(3): 768-775.
24. Jerrett M, Burnett RT, Pope CA, 3rd, Ito K, Thurston G, Krewski D, Shi Y, Calle E, Thun M. Long-term ozone exposure and mortality. *N Engl J Med* 2009: 360(11): 1085-1095.
25. Zanobetti A, Schwartz J. Ozone and survival in four cohorts with potentially predisposing diseases. *Am J Respir Crit Care Med* 2011: 184(7): 836-841.
26. Chuang KJ, Chan CC, Su TC, Lee CT, Tang CS. The effect of urban air pollution on inflammation, oxidative stress, coagulation, and autonomic dysfunction in young adults. *Am J Respir Crit Care Med* 2007: 176(4): 370-376.
27. Hong YC, Lee JT, Kim H, Kwon HJ. Air pollution: a new risk factor in ischemic stroke mortality. *Stroke* 2002: 33(9): 2165-2169.
28. Knowlton K, Rosenthal JE, Hogrefe C, Lynn B, Gaffin S, Goldberg R, Rosenzweig C, Civerolo K, Ku JY, Kinney PL. Assessing ozone-related health impacts under a changing climate. *Environ Health Perspect* 2004: 112(15): 1557-1563.
29. Bell M, Goldberg R, Hogrefe C, Kinney P, Knowlton K, Lynn B, Rosenthal J, Rosenzweig C, Patz J. Climate change, ambient ozone, and health in 50 US cities. *Climatic Change* 2007: 82(1): 61-76.
30. Hwang R, Burer M, Bell M. Smog in the forecast: global warming, ozone pollution and health in California. San Francisco: National Resources Defence Council; 2004.
31. Anderson H, Derwent R, Stedman J. Air pollution and climate change. In: McMichael A, Kovats R, eds. Health Effects of Climate Change in the UK. Department of Health, London, 2001; pp. 193-217.
32. West JJ, Fiore AM, Naik V, Horowitz LW, Schwarzkopf MD, Mauzerall DL. Ozone air quality and radiative forcing consequences of changes in ozone precursor emissions. *Geophys Res Lett* 2007: 34(6): L06806.
33. Chang HH, Zhou J, Fuentes M. Impact of climate change on ambient ozone level and mortality in southeastern United States. *Int J Environ Res Public Health* 2010: 7(7): 2866-2880.
34. Tagaris E, Liao KJ, Delucia AJ, Deck L, Amar P, Russell AG. Potential impact of climate change on air pollution-related human health effects. *Environ Sci Technol* 2009: 43(13): 4979-4988.
35. Doherty RM, Heal MR, Wilkinson P, Pattenden S, Vieno M, Armstrong B, Atkinson R, Chalabi Z, Kovats S, Milojevic A, Stevenson DS. Current and future climate- and air pollution-mediated impacts on human health. *Environ Health* 2009: 21(8).

36. Sheffield PE, Knowlton K, Carr JL, Kinney PL. Modeling of regional climate change effects on ground-level ozone and childhood asthma. *Am J Prev Med* 2011; 41(3): 251-257.
37. Ebi KL, McGregor G. Climate change, tropospheric ozone and particulate matter, and health impacts. *Environ Health Perspect* 2008; 116(11): 1449-1455.
38. Anenberg SC, Horowitz LW, Tong DQ, West JJ. An estimate of the global burden of anthropogenic ozone and fine particulate matter on premature human mortality using atmospheric modeling. *Environ Health Perspect* 2010; 118(9): 1189-1195.
39. Robertson L, Langner J, Engardt M. An Eulerian limited-area atmospheric transport model. *Journal of Applied Meteorology* 1999; 38: 190-210.
40. Kjellström E, Bärring L, Gollvik S, Hansson U, Jones C, Samuelsson P, Rummukainen M, Ullerstig A, Willén U, Wyser K. A 140-year simulation of the European climate with the new version of the Rossby Centre regional atmospheric climate model (RCA₃). Swedish Meteorological and Hydrological Institute, Norrköping, 2005.
41. Samuelsson P, Jones CG, Willén U, Ullerstig A, Gollvik S, Hansson ULF, Jansson C, Kjellström E, Nikulin G, Wyser K. The Rossby Centre Regional Climate model RCA₃: model description and performance. *Tellus A* 2011; 63(1): 4-23.
42. Engardt M, Bergstrom R, Andersson C. Climate and emission changes contributing to changes in near-surface ozone in Europe over the coming decades: results from model studies. *Ambio* 2009; 38(8): 452-458.
43. Andersson C, Engardt M. European ozone in a future climate: Importance of changes in dry deposition and isoprene emissions. *Journal of Geophysical Research* 2010; 115(D2): D02303.
44. Langner J, Engardt M, Andersson C, Steyn DGG, Trini Castelli ST. Modelling the Impact of Climate Change on Air Pollution over Europe Using the MATCH CTM Linked to an Ensemble of Regional Climate Scenarios. *Air Pollution Modeling and its Application XXI*. Springer Netherlands, 2010; pp. 627-635.
45. Langner J, Engardt M, Andersson C. We have only working title at the present, but should submit the manuscript before the year is up. We can update the reference with title and journal when your manuscript is back from the reviewers. *Manuscript to be submitted* 2012.
46. Kjellström E, Nikulin G, Hansson ULF, Strandberg G, Ullerstig A. 21st century changes in the European climate: uncertainties derived from an ensemble of regional climate model simulations. *Tellus A* 2011; 63(1): 24-40.
47. Bell ML, Peng RD, Dominici F. The exposure-response curve for ozone and risk of mortality and the adequacy of current ozone regulations. *Environ Health Perspect* 2006; 114(4): 532-536.
48. Amann M, Derwent D, Forsberg B, Haenninen O, Hurley F, Krzyzanowski M, Leeuw Fd, Liu SJ, Mandin C, Schneider J, Schwarze P, Simpson D. Health risks of ozone from long-range transboundary air pollution. Copenhagen: World Health Organization, Regional Office for Europe; 2008.
49. Klein Goldewijk K, Beusen A, Janssen P. Long-term dynamic modeling of global population and built-up area in a spatially explicit way: HYDE 3.1. *The Holocene*; 20(4): 565-573.
50. Committee on the Medical Effects of Air Pollutants. COMEAP Working paper 2002/9a. Is there a threshold for the effect of ozone on health? Is there an effect on mortality and respiratory or circulatory hospital admissions? NB: This is NOT an official publication of COMEAP or the UK Department of Health; 2003.
51. Nikulin G, Kjellström E, Hansson ULF, Strandberg G, Ullerstig A. Evaluation and future projections of temperature, precipitation and wind extremes over Europe in an ensemble of regional climate simulations. *Tellus A*; 63(1): 41-55.
52. WHO. Air quality guidelines: global update 2005. Particulate matter, ozone, nitrogen dioxide and sulfur dioxide. Copenhagen: WHO Regional Office for Europe; 2006

53. EC. Directive 2008/50/EC of the European Parliament and of the Council of 21 May 2008 on ambient air quality and cleaner air for Europe. *Official Journal of the European Union* 2008: L152: 1-44.
54. Anderson H, Derwent R, Stedman J, Hayman G. The health impact of climate change due to changes in air pollution. *In: Kovats R, ed. Health Effects of Climate Change in the UK 2008 An update of the Department of Health report 2001/2002.* Department of Health, London, 2008; pp. 91-105.
55. Watkiss P, Pye S, Holland M. CAFE CBA: baseline analysis 2000 to 2020. Brussels: European Commission; 2005.
56. De Marco A. Assessment of present and future risk to Italian forests and human health: modelling and mapping. *Environ Pollut* 2009: 157(5): 1407-1412.
57. Bell ML, Davis DL, Gouveia N, Borja-Aburto VH, Cifuentes LA. The avoidable health effects of air pollution in three Latin American cities: Santiago, São Paulo, and Mexico City. *Environmental Research* 2006: 100(3): 431-440.
58. Olsson D, Ekström M, Forsberg B. Temporal Variation in Air Pollution Concentrations and Preterm Birth - A Population Based Epidemiological Study. *International Journal of Environmental Research and Public Health* 2012: 9(1): 272-285.

Figure 1. Change in SOMO₃₅ ($\mu\text{g m}^{-3}\text{days}$) values due to climate change induced ozone exposure variations.









Figure 2. Change in all-cause mortality and respiratory hospitalizations (same relative change, %) due to climate change induced ozone exposure variations.

