

Fine and coarse particulate air pollution in relation to respiratory health in Sweden

Saskia M. Willers^{1,2}, Charlotta Eriksson¹, Lars Gidhagen³, Mats E. Nilsson^{1,4}, Göran Pershagen^{1,5}
and Tom Bellander^{1,5}

1. Institute of Environmental Medicine, Karolinska Institutet, Stockholm, Sweden
2. Institute for Risk Assessment Sciences, Utrecht University, Utrecht, The Netherlands
3. Swedish Meteorological and Hydrological Institute, Norrköping, Sweden
4. Department of Psychology, Stockholm University, Stockholm, Sweden
5. Centre for Occupational and Environmental Medicine, Stockholm County Council, Stockholm, Sweden.

Correspondence to:

Tom Bellander, PhD
Karolinska Institutet
Institute of Environmental Medicine
17177 Stockholm
Sweden
Phone: +46 (0)8 524 80003
Email: tom.bellander@ki.se

Abstract (199) (limit is 200 words)

Health effects have repeatedly been associated with residential levels of air pollution. However, it is difficult to disentangle effects of long-term exposure to locally generated and long-range transported pollutants as well as of exhaust emissions and wear particles from road traffic. We aimed to investigate effects of exposure to particulate matter (PM) fractions on respiratory health in the Swedish adult population, using an integrated assessment of sources at different geographical scales.

The study was based on a nationwide environmental health survey performed in 2007, including 25,851 adults aged 18 to 80. Individual exposure to PM at residential addresses was estimated by dispersion modelling of regional, urban and local sources. Associations between different size fractions or source categories and respiratory outcomes were analysed using multiple logistic regression, adjusting for individual and contextual confounding.

Exposure to locally generated wear particles (PM₁₋₁₀) showed associations for blocked nose/hay fever, chest tightness/cough and restricted activity days with odds ratios of 1.5 to 2 per 10 µg/m³. Associations were also seen for locally generated combustion particles, which disappeared following adjustment for exposure to wear particles. In conclusion, our data indicate that long-term exposure to locally generated road wear particles increases the risk of respiratory symptoms in adults.

Keywords

particulate matter, particle size, respiratory symptoms, sources, survey

Introduction

There is a large body of epidemiological evidence on health effects of particulate air pollution, including increased respiratory symptoms, decreased lung function and increased hospital admissions and mortality from cardio-pulmonary causes [1-5]. Mortality and cardiovascular events have been linked primarily to fine particulate matter (PM_{2.5}), possibly because it can penetrate deep into the lungs and cause systemic inflammation. Coarse particulate matter (commonly defined as particles in the 2.5 to 10 µm aerodynamic diameter range) has mostly been associated with non-fatal respiratory effects [5]. However, a review of Brunekreef & Forsberg [6] indicated that coarse PM is associated with cardiovascular disease and increased mortality as well. It is difficult to separate health effects of long-term exposure to exhaust emissions and wear particles from road traffic, and to separate effects of locally generated and long-range transported pollutants. Further epidemiological evidence to disentangle these relationships is needed. Although the background concentration of particulate air pollution in the Nordic countries is lower than in the rest of Europe [7], there are major concerns about health effects of exposure to particulate air pollution from traffic [8-10]. The use of studded winter tyres on cars and the sanding of roads are common in Northern Europe, and road wear constitutes an important source of coarse PM [11]. There are only few large population studies investigating health effects of air pollution from road traffic which include urban as well as non-urban adult populations, and use spatially resolved air pollution data. Within the ROADSIDE project, we have data on air pollution exposure and health outcomes for approximately 26,000 people from 18 to 80 years of age over the whole country of Sweden. The main objectives of the present study were to investigate associations between exposure to total PM₁₀, long-range transport particles, locally generated combustion particles and locally generated road wear particles at the home address and self-reported respiratory health outcomes in a general population sample in Sweden.

Methods

Study population and design

ROADSIDE is a large cross-sectional study based on the National Environmental Health Survey (NEHS 2007) conducted in Sweden in 2007. The aim of this nationwide postal survey was to study exposure to several environmental and lifestyle factors as well as related health outcomes in the adult population (18 to 80 years of age) living in Sweden for at least 5 years. According to the Swedish population register this target population consisted of 6,761,887 persons. Random

sample selection was conducted in two steps. First, a sample of 10,500 persons, equally distributed over Sweden's 21 counties was selected. Second, this sample was enhanced with 33,405 persons from 10 counties because of supplementary funding. In total, the survey was sent to 43,905 persons (0.6% of the target population) of which 25,851 persons (59.4%) responded. The survey was sent out and processed by Statistics Sweden and had been approved by the regional ethical committee in Stockholm [12]. A map of Sweden with its county borders, showing response rates and counties with enhanced sampling is provided in figure 1a.

Exposure assessment

Individual exposure to PM was assessed as average outdoor levels at the home address estimated by dispersion modelling. The home addresses of the respondents to the NEHS 2007 were geocoded by the Swedish Mapping, Cadastre and Land Registration Authority. The SIMAIR modelling system [13, 14] was used to estimate address-specific outdoor levels of PM₁₀. SIMAIR is a national modelling system that combines air pollution contributions on three scales: A regional contribution from simulation on the European scale, an urban contribution calculated on a 1x1 km grid over about 100 cities and towns in Sweden and a local contribution from traffic within a radius of 250 m from the home address. The determination of the local traffic impact is based on a unique national road and traffic database, covering the majority of the Swedish roads, managed by the Swedish Transport Administration [15]. Emission factors for the exhaust part of PM₁₀ are calculated by the Assessment and Reliability of Transport Emission Models and Inventory Systems (ARTEMIS) working group [16]. A semi-empirical model is used for the non-exhaust part of PM₁₀, considering generation, suspension and resuspension of wear particles in the supermicron fraction [17]. The SIMAIR system enables splitting up of the PM₁₀ size fractions for the urban as well as the local traffic contribution of PM₁₀. Exhaust- and combustion-related particles were assumed to have a diameter <1 µm (indicated as PM₁) and the resuspended road and tyre wear particles were assumed to have a diameter ranging from 1 to 10 µm (indicated as PM₁₋₁₀). The long-range regional contribution to PM₁₀ was estimated to be equally distributed between the supermicron (PM₁₋₁₀) and submicron (PM₁) fraction, based on observations from a rural background monitor in central Sweden [18]. The origin and composition of these two fractions in the regional contribution is different from the local traffic and urban contribution. The regional PM₁ concentration contains a larger part of aged inorganic and organic secondary aerosols and to a lesser extent sea salt, while the regional PM₁₋₁₀ concentration mainly contains sea salt and mineral dust transported over larger distances. The SIMAIR system uses meteorological input data relevant for dispersion models – e.g. wind speed and direction,

temperature, cloudiness, global radiation, sensible heat flux, friction velocity, humidity and precipitation. Further details of the SIMAIR modelling system, as well as results of the model validation, have been described previously [13, 14].

In this study, we investigated associations for total PM₁₀ level, long-range transport particles (regional PM₁₀ contribution), locally generated PM₁₀ level (local traffic + urban PM₁₀ contribution), locally generated combustion particles (local traffic + urban PM₁ fraction) and locally generated wear particles (local traffic + urban PM₁₋₁₀ fraction).

Health outcome assessment

The NEHS 2007 survey enquired about annoyance from specific sources, prevalence of respiratory, cardiovascular and allergic symptoms and diseases, as well as quality of life. Furthermore, it contained questions about smoking and sunbathing behaviour, home and living environment characteristics, food and water consumption, and mobile phone use. Respondents returned the questionnaire to Statistics Sweden (SCB). SCB digitized the answers, added registry data on country of origin, civil status, income, education and home geographical coordinates and estimated a survey calibration weight variable. This calibration weight variable was used to calculate results that are representative for the target population. Within a geographical sampling stratum, the differences in calibration weight represent differences in response rate in substrata defined by gender, age, civil status, income, education and country of origin. The weight was on average 260 and ranged from 6 to 1874.

Because of the cross-sectional nature of the survey and the self-reporting, we focused on current and recent respiratory health in the present study. Respiratory health outcomes analysed were asthma symptoms in the last 12 months, the use of asthma medication in the last 12 months, having a blocked nose or hay fever symptoms for more than 4 days a week in the last 12 months, woken by chest tightness or cough in the last 3 months, and having restricted activity days due to respiratory symptoms in the last 3 months.

Statistical analyses

Descriptive statistics were produced for the total study population and stratified by region. The regions were defined as follows: Scania (Skåne county), South-East (Östergötland, Jönköping, Kronoberg, Kalmar and Blekinge counties, and the island Gotland), Mälardalen (Uppsala, Södermanland, Örebro and Västmanland counties), West (Värmland, Västra Götaland and Halland counties), Stockholm (Stockholm county), Central (Jämtland, Västernorrland, Gävleborg, Dalarna counties), and North (Norrbotten and Västerbotten counties), see Figure 1a.

Associations between traffic-related PM₁₀ exposure and respiratory health outcomes were analysed using multivariate logistic regression adjusted for the respondent's gender, age, educational level, household income, smoking, passive smoking, moist damage or mould exposure in the home and country of birth (see Table 1). Furthermore, we used the proportion of higher educated adults and the proportion of unemployed adults in the municipality to additionally adjust for contextual confounding. The data to construct the two contextual variables were derived from internet databases of Statistics Sweden for level of education in municipalities in 2007 [19], and of the Municipality Database (Kommundatabasen) for unemployment rate in 2007 [20]. For total PM₁₀, long-range transport particles, locally generated PM₁₀ and locally generated wear particles, associations were calculated by 10 µg/m³ increase, while for the locally generated combustion particles associations were calculated by 1 µg/m³ increase. To compare the health impact of locally generated combustion particles versus locally generated wear particles, we also calculated associations by an increase from the 5th to the 95th percentile in exposure levels.

In order to explore if differences in response rate in socioeconomic substrata were related to levels of air pollution, and thus potential confounders, we calculated the correlations between the relative calibration weights and total PM₁₀ exposure.

Results

Female and middle-aged respondents were slightly overrepresented in the study population, with little regional variation (Table 1). The largest proportion of persons with higher vocational/university education was in the Stockholm region. Smokers constituted 18% of the total study population, with a lower proportion in region North (14%). Passive smoking was most common in the southern region Scania. Moist damage was present in approximately 15% of the homes, while visible mould was reported for 5%. More than 90% of the study population was of Nordic origin. The highest proportions of participants of non-Nordic origin were in Scania and Stockholm.

The estimated residential outdoor levels of particulate air pollution in Sweden could be modelled for 25,846 of the 25,851 study participants (Table 2 and Figures 1b and 2). Total PM₁₀ levels of our total study population appeared to have a Gaussian distribution with a mean of 11.3 µg/m³, a 5 to 95% range from 8.0 to 16.1 µg/m³, and a few addresses between 20 and 40 µg/m³. Highest mean concentrations of total PM₁₀ were seen in the regions Scania, Stockholm and West, which contain the three largest cities in Sweden: Stockholm, Göteborg (region West) and Malmö (region

Scania). In all regions, total PM₁₀ level was dominated by long-range transport particles, with an overall average of 10 µg/m³ (88% of the total PM₁₀ level). Long-range transport particle levels (regional PM₁₀ contribution) ranged from approximately 4 (region North) to 14 µg/m³ (region Scania; Figure 1b). Locally generated PM₁₀ level had a skewed distribution with an average of 1.3 µg/m³. Overall, the locally generated PM₁₀ level was dominated by wear particles (62%) with an average of 0.8 µg/m³, the rest being traffic exhaust and other combustion particles with an average of 0.5 µg/m³. The highest levels of locally generated combustion particles were found in region Stockholm, Mälardalen and South-East+Gotland while the highest levels of locally generated road wear particles were found in region Stockholm, Mälardalen and West. The differences in response rate across geographic and socioeconomic substrata of the population were not related to estimated total PM₁₀ levels (data not shown).

The prevalence of respiratory outcomes is shown in table 3. Asthma symptoms and the use of asthma medication in the last 12 months seemed to be very similarly reported, by about 8% of the total study population. There were distinct regional differences, with the highest prevalence in region Mälardalen and lowest in Scania and West. Blocked nose/hay fever for more than 4 days a week in the last 12 months was reported by 18.0%, with smaller regional differences. Woken by chest tightness/cough in the last 3 months was reported by 19.4% and restricted activity days due to respiratory symptoms in the last 3 months by 10.9% of the population, both also with smaller regional differences.

Figure 3 shows associations between total PM₁₀ as well as the different fractions and respiratory health outcomes in the total study population. Total PM₁₀ concentration was positively associated with chest tightness/cough and restricted activity days in the last 3 months. The long-range transport part of PM₁₀ was not associated with any outcome except for a non-expected marginally significant inverse association with asthma symptoms (OR per 10 µg/m³ difference 0.69; 95% CI 0.49-0.99). Locally generated PM₁₀ was significantly associated with woken by chest tightness and cough in the last 3 months and with restricted activity days in the last 3 months, but also with blocked nose/hay fever in the last 12 months. When separating the locally generated particles in wear and combustion particles, both fractions showed elevated risks for respiratory symptoms. Locally generated wear particles were significantly associated with blocked nose/hay fever (OR per 10 µg/m³ difference 1.40; 95% CI 1.02-1.90), chest tightness/cough (OR 1.91; 95% CI 1.40-2.60) and restricted activity days (OR 1.50; 95% CI 1.05-2.16), while locally generated combustion particles were significantly associated with blocked nose/hay fever (OR per 1 µg/m³ difference 1.09; 1.02-1.16), and with woken by chest tightness/cough (OR 1.08; 95% CI 1.01-1.15).

In order to compare the effects of locally generated wear and combustion particles we also calculated associations for an exposure difference between the 5th to the 95th percentile of the distribution in the study participants. These exposure differences were 3.22 and 1.74 $\mu\text{g}/\text{m}^3$, respectively. The corresponding odds ratios were fairly similar: blocked nose/hay fever 1.11 (95% CI 1.00-1.23) and 1.15 (1.03-1.29), chest tightness/cough 1.23 (1.11-1.36) and 1.14 (1.02-1.28), and restricted activity days 1.14 (1.01-1.28) and 1.10 (0.97-1.27), respectively. Exposure to locally generated wear and combustion particles were highly correlated ($r=0.75$), and in regression models including both exposure parameters, the effect of the combustion particles disappeared while the effect of the wear particles was strengthened (data not shown). When we performed analyses restricting the population to those resident at the current address for at least five years the results were essentially the same.

Discussion

We have investigated the associations between exposure to different fractions of particulate air pollution and current or recent respiratory symptoms in Swedish adults. Our results indicate significant effects of exposure to total PM_{10} , and locally generated PM_{10} , including wear and combustion particles, on the prevalence of restricted activity days due to respiratory symptoms, woken by chest tightness/cough and blocked nose/hay fever (only statistically significant for locally generated PM_{10}). However, we did not find any consistent effects of PM exposure on asthma outcomes in the present study.

Our results are broadly in line with other studies showing respiratory effects from coarse particles. A systematic review of the effects of coarse airborne particles on health observed that in studies of COPD, asthma and respiratory admissions, coarse PM has a stronger or as strong short-term effect as fine PM, suggesting that coarse PM may lead to adverse responses in the lungs triggering processes leading to hospital admissions [6]. Among recent studies, Chen et al. [21] showed that ambient coarse PM was positively associated with respiratory hospital admissions among the elderly population in Vancouver, Canada. Lin et al. [22] found a detrimental effect of coarse particulate matter on hospitalization for respiratory infections in children. A study conducted in France also showed positive associations between exposure to $\text{PM}_{2.5-10}$ and hospitalization for respiratory infections in children.[23] Another French study, using a dispersion model, found positive associations between PM_{10} exposure and asthma and allergies in children. [24]

It was possible to explore the effects of different particle size fractions, which would have been more problematic in smaller areas because of the locally very high correlation between the fractions. In our study, locally generated wear particles appear less harmful than combustion particles on mass basis, but when the associations were calculated for difference between the 5th to the 95th percentile of the distribution of the two fractions the results were rather similar. When both these types of locally generated particles were entered into the same model, the effect of combustion particles disappeared while the effect of the wear particles was strengthened. It should however be noted that also in our study the two fractions to a large extent come from the same source, and thus are substantially correlated ($r=0.75$), which implies that the results regarding the specific role of each fraction have to be interpreted with caution.

Our study was based on a large sample (nearly 26.000 people) distributed over a large geographical area (449,964 km²), including densely populated areas - the three major cities Stockholm, Göteborg and Malmö - as well as rural areas all over Sweden. This resulted in a greater than usual exposure contrast for a Swedish population, and in a high statistical power of the analyses.

The response rate of the initial environmental health survey was reasonable (59.4%). From a non-response analysis of the question: "How do you assess your general health status, compared to others of your age?", it was concluded that the study population well represented the target population of long-term Swedish residents aged 18 to 80 [12]. The difference in response rates over the socio-economical and geographical strata was not related to air pollution exposure, which implies that no bias was introduced by the lack of total participation in the survey.

A limitation of this study was the cross-sectional study design, which makes it difficult to study effects of long-term exposure. Subjects who have developed respiratory problems may have moved away from areas with high air pollution exposure, which could result in apparent protective effects from air pollution. We therefore focussed on the effects of air pollution exposure current and recent respiratory symptoms (in the last 3 months or in the last year).

Restricting the analyses to subjects living at their address for more than 5 years did not substantially change the results.

The different air quality gradients, as simulated by the SIMAIR model system, vary in quality. The gradient of the regional contribution over Sweden was estimated from a combination of model results and measurement data, and is expected to have high validity for the model year (2004). The validity of the urban contribution of PM is highly dependent on the information on traffic in the 1x1 km gridnet, which was of good quality [13, 14]. The total contribution from residential wood combustion to PM exposure as well as its spatio-temporal distribution is less

well known. As for the local impact of traffic emissions within a radius of 250 m, the spatial distribution of the emissions should be well described for all state-owned roads and for the major city roads. However, traffic on minor city roads was to a great extent simulated, and thus described with poorer quality. The modelling was performed with hourly input of meteorology for the model year 2004. The questionnaire data are from 2007 but very little is expected to have happened with any of the gradients in the years before and after the model year.

SIMAIR model output has been extensively evaluated against monitor data of 12 urban background and 15 street site locations [25]. The results of these evaluations show that the SIMAIR model performs substantially better than the EU Directive's quality criteria for PM₁₀, that require a maximum deviation between the simulated and monitored annual average concentration levels less than 50% for 90 % of individual monitoring points [26]. Furthermore, the results of the comparison of the SIMAIR model with monitored data at six urban background and four street site locations in different parts of Sweden showed that the difference between simulated and monitored annual averages of PM₁₀ levels were below ca 20% for urban background and ca 10% for street level concentrations, with correlation coefficients for the time-series of daily average PM₁₀ varying between 0.56 to 0.76.

An important uncertainty may be found in basing the exposure determination on outdoor levels at the home address only. People spend most of their time indoors, but indoor PM levels in Sweden are similar to outdoor levels [27], and the home address is a common approach found to constitute a fairly good proxy for especially PM_{2.5} exposure [28]. However, other environments that may give significant contributions are school, the workplace and traffic. Future studies could benefit from including estimates of these contributions to individual exposure.

Health outcome variables used in this study were self-reported and derived from a questionnaire containing questions on both living environment and personal health. Self-reporting of health is clearly subject to error [29]. Biased reporting of the outcomes associated with the estimated air pollution is however unlikely as the data on air pollution levels used in our analyses were derived independently from the questionnaire. Our study has been conducted in a very heterogeneous study area varying from very sparsely populated rural areas to the urban areas of Stockholm, Göteborg and Malmö. Furthermore, the exposure to air pollution and the prevalence of the investigated respiratory symptoms vary over the country as well. Earlier studies in Sweden have found that the socio-economic context of the area of residence is associated with levels of air pollution [30] as well as disease risk [31]. Potential contextual confounding by socioeconomic position of the area of residence has been taken into account by conducting multilevel analyses adjusting for percentage of high educated persons and percentage of people unemployed in the

municipality. These two contextual variables were both associated with air pollution levels and the investigated respiratory symptoms and including them in the models slightly attenuated the risk estimates.

The apparent protective effect of long-range transport particles on asthma outcomes is probably due to the fact that asthma is more prevalent in the north, while long-range transport particle concentration shows the opposite gradient. Earlier studies in Sweden have also shown an increased prevalence of asthma and COPD in adults and children in Northern Sweden compared to Southern Sweden [32-35]. Climate factors such as temperature and humidity, lifestyle factors such as nutrition, or indoor air quality may explain these differences in prevalence. However, the exact reason is still unknown, underlining the need of great caution in the interpretation of studies with large-scale differences in both environmental contaminants and health effects. In conclusion, our data indicate that exposure to locally generated road wear particles increases the risk of respiratory symptoms in adults.

Support statement

This study was funded by the Swedish Research Council FORMAS. The National Environmental Health Survey was funded by the Swedish National Board of Welfare together with the regions in which sampling was enhanced.

Acknowledgements

We would like to thank Emma Nordling and Magnus Wickman who were involved in the conception of the study design.

References

1. Schikowski T, Sugiri D, Ranft U, Gehring U, Heinrich J, Wichmann HE, Krämer U. Long-term air pollution exposure and living close to busy roads are associated with COPD in women. *Respir Res* 2005;6:152.
2. Churg A, Brauer M, del Carmen Avila-Casado M, Fortoul TI, Wright JL. Chronic exposure to high levels of particulate air pollution and small airway remodeling. *Environ Health Perspect* 2003;111:714-718.
3. Ackermann-Lieblich U, Leuenberger P, Schwartz J, Schindler C, Monn C, Bolognini G, Bongard JP, Brandli O, Domenighetti G, Elsasser S, Grize L, Karrer W, Keller R, Keller-

- Wossidlo H, Künzli N, Martin BW, Medici TC, Perruchoud AP, Schoni MP, Tschopp JM, Villiger B, Wuthrich B, Zellweger JP, Zemp E. Lung function and long term exposure to air pollutants in Switzerland. Study on Air Pollution and Lung Diseases in Adults (SAPALDIA) Team. *Am J Respir Crit Care Med* 1997;155:122-129.
4. Brunekreef B, Holgate ST. Air pollution and health. *Lancet* 2002;360:1233-1242.
 5. Pope CA, 3rd, Dockery DW. Health effects of fine particulate air pollution: lines that connect. *J Air Waste Manag Assoc* 2006;56:709-742.
 6. Brunekreef B, Forsberg B. Epidemiological evidence of effects of coarse airborne particles on health. *Eur Respir J* 2005;26:309-318.
 7. Horálek J, Denby B, de Smet P, de Leeuw F, Kurfürst P, Swart R, van Noije T. Spatial mapping of air quality for European scale ETC/ACC Technical Paper 2006/6.
http://acm.eionet.europa.eu/reports/ETCACC_TechnPaper_2006_6_Spat_AQ. Date last updated: June 18 2007. Date last accessed: October 10 2011.
 8. Nordling E, Berglind N, Melén E, Emenius G, Hallberg J, Nyberg F, Pershagen G, Svartengren M, Wickman M, Bellander T. Traffic-related air pollution and childhood respiratory symptoms, function and allergies. *Epidemiology* 2008;19:401-408.
 9. Nyberg F, Gustavsson P, Järup L, Bellander T, Berglind N, Jakobsson R, Pershagen G. Urban air pollution and lung cancer in Stockholm. *Epidemiology* 2000;11:487-495.
 10. Rosenlund M, Forastiere F, Porta D, De Sario M, Badaloni C, Perucci CA. Traffic-related air pollution in relation to respiratory symptoms, allergic sensitisation and lung function in schoolchildren. *Thorax* 2009;64:573-580.
 11. Penttinen P, Timonen KL, Tiittanen P, Mirme A, Ruuskanen J, Pekkanen J. Ultrafine particles in urban air and respiratory health among adult asthmatics. *Eur Respir J* 2001;17:428-435.
 12. Svensson H, Egmar A-C, Rodvall Y, Bellander T. National environmental health survey 2007 (Nationella miljöhälsoenkäten 2007. In: Environmental health report 2009 (Miljöhälsorapport 2009). The National Board of Health and Welfare (Socialstyrelsen) Stockholm, Sweden, 2009 (in Swedish). Extended summary in English available at:
www.socialstyrelsen.se/publikationer2009/2009-126-116. Date last updated: April 2009. Date last accessed: October 10 2011.
 13. Gidhagen L, Johansson H, Omstedt G. SIMAIR-Evaluation tool for meeting the EU directive on air pollution limits. *Atmospheric Environment* 2009;43:1029-1036.
 14. Omstedt G, Andersson S, Gidhagen L, Robertson L. New model tools for meeting the target of the EU Air Quality Directive: description, validation and evaluation of local air quality

- improvement due to reduction of studded tyre use on Swedish roads. *Int J Environment and Pollution* 2011; 47(1/2/3/4): 79-96.
15. Swedish Road Administration, 2008. NVDB Contents – overview, NVDB Contents – feature types, NVDB Contents – road network (PDF reports).
<http://www22.vv.se/filer/68850/NVDBContents-Overview.pdf>. Date last updated: October 1, 2008. Date last accessed: October 28, 2011.
16. André M, Rapone M. Analysis and modelling of the pollutant emissions from European cars regarding the driving characteristics and test cycles. *Atmospheric Environment* 2009;43:986-995.
17. Omstedt G, Bringfelt B, Johansson C. A model for vehicle induced non-tailpipe emissions of particles along Swedish roads. *Atmospheric Environment* 2005;39:6088-6097.
18. Johansson C, Andersson C, Bergström R, Krecl P. Exposure to particles due to local and non-local sources in Stockholm – estimates based on modeling and measurements 1997-2006. ITM Report 175 September 2008. http://slb.nu/slb/rapporter/pdf8/itm2008_175.pdf. Date last updated: September 2008. Date last accessed: October 10 2011.
19. Statistics Sweden (SCB). Database: Population 16-74 years of age by municipality, level of education, age and sex. Year 1985-2010. In table: Population 16-74 years of age by highest level of education, age and sex. Year 1985-2010.
<http://www.ssd.scb.se/databaser/makro/Produkt.asp?produktid=UF0506&lang=2>. Date last updated: April 19 2011. Date last accessed: October 10 2011.
20. Kommun- och landstingsdatabasen. Database: Unemployment by municipality in 2007 (Arbetslöshet (%) i kommun 2007). <http://www.kolada.se/portal.php?page=nyckeltalslista>. Date last updated: October 2011. Date last accessed: October 10 2011.
21. Chen Y, Krewski D, Burnett RT, Shi Y, McGrail KM. The effect of coarse ambient particulate matter on first, second, and overall hospital admissions for respiratory disease among the elderly. *Inhalation Toxicology* 2005;17:649-655.
22. Lin M, Stieb DM, Chen Y. Coarse particulate matter and hospitalization for respiratory infections in children younger than 15 years in Toronto: a case-crossover analysis. *Pediatrics* 2005;116:e235-e40.
23. Host S, Larrieu S, Pascal L, Blachard M, Declerq C, Fabre P, Jusot J-F, Chardon B, Le Tertre A, Wagner V, Prouvost V, Lefranc A. Short-term associations between fine and coarse particles and hospital admissions for cardiorespiratory diseases in six French cities. *Occup Environ Med* 2008;65:544-51.

24. Penard-Morand C, Raheison C, Charpin D, Kopferschmitt C, Lavaud F, Caillaud D, Annesi-Maesano I. Long-term exposure to close-proximity air pollution and asthma and allergies in urban children. *Eur Respir J* 2010;36:33-40.
25. Andersson S, Omstedt G. Validering av SIMAIR mot mätningar av PM10, NO2 och bensen. (Validation of SIMAIR against monitor data of PM10, NO2 and benzene). Utvärdering för svenska tätorter och trafikmiljöer avseende år 2004 och 2005. Swedish Meteorological and Hydrological Institute (SMHI) rapport, serie meteorologi 137. Norrköping, Sweden 2009 (in Swedish). <http://www.smhi.se/publikationer/validering-av-simair-mot-matningar-av-pm10-no2-och-bensen-utvardering-for-svenska-tatorter-och-trafikmiljoer-avseende-ar-2004-och-2005-1.7370>. Date last updated: August 2009. Date last accessed: October 10 2011.
26. European Commission Environment. EU Directive 2008/50/EC of the European Parliament and of the Council on ambient air quality and cleaner air for Europe. <http://eur-lex.europa.eu/LexUriServ/LexUriServ.do?uri=OJ:L:2008:152:0001:0044:EN:PDF>. Date last updated: June 11 2008. Date last accessed: October 10 2011.
27. Wichmann J, Lind T, Nilsson MA-M, Bellander T. PM2.5, soot and NO2 indoor- outdoor relationships at homes, pre-schools and schools in Stockholm, Sweden. *Atmospheric Environment* 2010;44:4536-4544.
28. Van Roosbroeck S, Hoek G, Meliefste K, Janssen NAH, Brunekreef B. Validity of residential traffic intensity as an estimate of long-term personal exposure to traffic-related air pollution among adults. *Environ Sci Technol* 2008;42:1337-44.
29. Baker M, Stabile M, Deri C. What do self-reported, objective, measures of health measure? *Journal of Human Resources* 2004;39:1067-1093.
30. Stroh E, Oudin A, Gustafsson S, Pilesjö P, Harrie L, Strömberg U, Jakobsson K. et al. Are associations between socio-economic characteristics and exposure to air pollution a question of study area size? An example from Scania, Sweden. *Int J Health Geogr* 2005;4:30.
31. Kölegård Stjärne M, Diderichsen F, Reuterwall C, Hallqvist J; Sheep Study Group. Socioeconomic context in area of living and risk of myocardial infarction: results from Stockholm Heart Epidemiology Program (SHEEP). *J Epidemiol Community Health* 2002;56:29-35.
32. Forsberg B, Pekkanen J, Clench-Aas J, Mårtensson MB, Stjernberg N, Bartonova A, Timonen KL, Skerfving S. Childhood asthma in four regions in Scandinavia: risk factors and avoidance effects. *Int J Epidemiol* 1997;26:610-619.
33. Bröms K, Norbäck D, Eriksson M, Sundelin C, Svärdsudd K. Effect of degree of urbanisation on age and sex-specific asthma prevalence in Swedish preschool children. *BMC Public Health* 2009;9:303.

34. Lundbäck B, Nyström L, Rosenhall L, Stjernberg N. Obstructive lung disease in northern Sweden: respiratory symptoms assessed in a postal survey. *Eur Respir J* 1991;4:257-266.
35. Montnémary P, Adelroth E, Heuman K, Johannisson A, Johansson SA, Lindholm LH, Lundbäck B, Löfdahl CG.et al. Prevalence of obstructive lung diseases and respiratory symptoms in southern Sweden. *Respir Med* 1998;92:1337-1345.

Tables

Table 1. Characteristics of the total study population included in the Swedish Environmental Survey of 2007, and stratified by region

Characteristic	Total	Scania	SouthEast+ Gotland	Mälardalen	West	Stockholm	Central	North
Total population (N)	25,851	1,122	10,305	1,182	3,941	5,921	2,147	1,233
Gender, n (%)								
Male	11,832 (45.8)	496 (44.2)	4,724 (45.8)	558 (47.2)	1,829 (46.4)	2,632 (44.5)	1,016 (47.3)	577 (46.8)
Female	14,019 (54.4)	626 (55.8)	5,581 (54.2)	624 (52.8)	2,112 (53.6)	3,289 (55.6)	1,131 (52.7)	656 (53.2)
Age category (years), n (%)								
18-29	3,513 (13.6)	167 (14.9)	1,361 (13.2)	167 (14.1)	567 (14.4)	781 (13.2)	256 (11.9)	214 (17.4)
30-39	3,900 (15.1)	187 (16.7)	1,360 (13.2)	190 (16.1)	646 (16.4)	1,066 (18.0)	284 (13.2)	167 (13.5)
40-49	4,661 (18.0)	202 (18.0)	1,824 (17.7)	194 (16.4)	670 (17.0)	1,199 (20.3)	380 (17.7)	192 (15.6)
50-59	4,852 (18.8)	198 (17.7)	1,946 (18.9)	229 (19.4)	754 (19.1)	1,045 (17.7)	441 (20.5)	239 (19.4)
60-69	5,207 (20.1)	210 (18.7)	2,228 (21.6)	249 (21.1)	722 (18.3)	1,127 (19.0)	437 (20.4)	234 (19.0)
70-80	3,718 (14.4)	158 (14.1)	1,586 (15.4)	153 (12.9)	582 (14.8)	703 (11.9)	349 (16.3)	187 (15.2)
Educational level, n (%)								
Primary school	4,659 (18.0)	193 (17.2)	2,131 (20.7)	219 (18.5)	721 (18.3)	779 (13.2)	434 (20.2)	182 (14.8)
High school	11,030 (42.7)	432 (38.5)	4,528 (43.9)	486 (41.1)	1,631 (41.4)	2,415 (40.8)	986 (45.9)	552 (44.8)
Lower vocational education	1,489 (5.8)	75 (6.7)	504 (4.9)	77 (6.5)	254 (6.5)	399 (6.7)	114 (5.3)	66 (5.4)
Higher vocational education/University	6,921 (26.8)	325 (29.0)	2,417 (23.5)	318 (26.9)	1,057 (26.8)	2,003 (33.8)	457 (21.3)	344 (27.9)
Smoking, n (%)								
Non-smoker	12,800 (49.5)	544 (48.5)	5,152 (50.0)	591 (50.0)	1,954 (49.6)	2,828 (47.8)	1,077 (50.2)	654 (53.0)
Ex-smoker	8,182 (31.7)	355 (31.6)	3,190 (31.0)	372 (31.5)	1,195 (30.3)	1,996 (33.7)	688 (32.0)	368 (31.3)
Current smoker	4,559 (17.6)	211 (18.8)	1,836 (17.8)	196 (16.6)	741 (18.8)	1,039 (17.6)	359 (16.7)	177 (14.4)
ETS exposure in home, n (%)	1,865 (7.2)	98 (8.7)	749 (7.3)	80 (6.8)	290 (7.4)	435 (7.4)	133 (6.2)	80 (6.5)
Moist damage in home, n (%)	3,971 (15.4)	159 (14.2)	1,518 (14.7)	199 (16.8)	663 (16.8)	891 (15.1)	379 (17.7)	162 (13.1)
Visible mould in home, n (%)	1,368 (5.3)	66 (5.9)	486 (4.7)	67 (5.7)	250 (6.3)	352 (5.9)	88 (4.1)	59 (4.8)
Country of birth, n (%)								
Nordic country	23,921 (92.5)	993 (88.5)	9642 (93.6)	1,099 (93.0)	3,616 (91.8)	5,282 (89.2)	2,096 (97.6)	1,193 (96.8)
Europe/North America	1,110 (4.3)	79 (7.0)	422 (4.1)	41 (3.5)	196 (5.0)	330 (5.6)	29 (1.4)	13 (1.1)
Other	820 (3.2)	50 (4.5)	241 (2.3)	42 (3.6)	129 (3.3)	309 (5.2)	22 (1.0)	27 (2.2)
% Higher [#] educated in municipality*								
Mean (SD)	16.8 (6.9)	18.5 (7.9)	14.3 (5.6)	15.9 (7.1)	17.4 (5.7)	21.8 (7.5)	13.1 (3.3)	17.5 (7.2)
% Unemployed in municipality*								
Mean (SD)	11.5 (3.9)	12.2 (2.7)	12.8 (3.2)	12.1 (3.0)	11.7 (2.5)	7.3 (3.2)	14.2 (3.3)	13.0 (3.6)

Higher vocational education or university * Contextual variable on municipality level

Table 2. Estimated residential outdoor levels of total PM₁₀ and different PM fractions (µg/m³) for the total study population in the Swedish Environmental Survey in 2007, and stratified by region

Exposure	Total	Scania	SouthEast+ Gotland	Mälardalen	West	Stockholm	Central	North
N	25,846	1,122	10,305	1,182	3,939	5,921	2,147	1,230
PM ₁₀ total								
Mean (SD)	11.3 (2.5)	13.7 (1.2)	10.5 (0.9)	10.0 (1.1)	12.8 (2.5)	13.2 (2.6)	9.0 (1.3)	7.4 (1.6)
Median (IQR)	10.8 (2.8)	13.4 (1.4)	10.5 (1.3)	9.9 (1.6)	12.8 (3.5)	12.8 (3.7)	8.8 (1.7)	7.5 (2.2)
Min-Max	3.6-40.0	11.1-17.6	8.8-14.7	8.1-15.8	6.5-21.4	9.7-40.0	6.1-14.2	3.6-11.3
Long-range transport particles								
Mean (SD)	10.0 (1.5)	13.0 (0.5)	9.8 (0.7)	9.1 (0.5)	11.6 (1.4)	10.2 (0.2)	8.2 (0.7)	6.8 (1.3)
Median (IQR)	10.0 (1.4)	13.2 (0.4)	9.7 (1.2)	9.0 (0.9)	12.3 (1.9)	10.2 (0.2)	8.4 (1.1)	6.6 (2.5)
Min-Max	3.6-13.6	11.0-13.6	8.8-12.0	8.1-10.2	6.5-13.3	9.7-10.7	6.1-9.3	3.6-8.6
Locally generated PM ₁₀								
Mean (SD)	1.3 (1.8)	0.6 (0.8)	0.7 (0.7)	0.9 (1.0)	1.3 (1.6)	3.0 (2.6)	0.8 (0.9)	0.6 (0.7)
Median (IQR)	0.7 (1.9)	0.2 (1.0)	0.5 (1.1)	0.7 (1.6)	0.5 (2.3)	2.5 (3.5)	0.3 (1.3)	0.4 (1.0)
Min-Max	0-29.8	0-4.2	0-5.2	0-6.9	0-8.8	0-29.8	0-5.5	0-3.8
Locally generated wear particles								
Mean (SD)	0.8 (1.3)	0.4 (0.6)	0.4 (0.5)	0.5 (0.7)	0.8 (1.1)	2.0 (2.0)	0.4 (0.5)	0.3 (0.4)
Median (IQR)	0.3 (1.1)	0.1 (0.6)	0.2 (0.5)	0.3 (0.9)	0.3 (1.4)	1.5 (2.3)	0.1 (0.5)	0.2 (0.5)
Min-Max	0-25.2	0-2.9	0-4.3	0-5.4	0-7.0	0-24.2	0-3.8	0-2.8
Locally generated combustion particles								
Mean (SD)	0.5 (0.6)	0.2 (0.3)	0.3 (0.4)	0.4 (0.4)	0.4 (0.5)	1.0 (0.8)	0.4 (0.5)	0.3 (0.4)
Median (IQR)	0.3 (0.8)	0.1 (0.4)	0.2 (0.6)	0.3 (0.7)	0.1 (0.8)	1.1 (1.3)	0.1 (0.7)	0.2 (0.5)
Min-Max	0-5.6	0-1.5	0-1.9	0-1.5	0-3.0	0-5.6	0-2.9	0-3.5

Table 3. Prevalence (%) of self-reported recent respiratory symptoms in the total study population in the Swedish Environmental Survey of 2007, and stratified by region

Outcome	Total	Scania	SouthEast+ Gotland	Mälardalen	West	Stockholm	Central	North
N	25,851	1,122	10,305	1,182	3,941	5,921	2,147	1,233
Asthma symptoms last 12 months	7.5	6.0	7.1	10.2	6.6	8.2	7.6	8.3
Asthma medication use last 12 month	7.8	7.6	7.5	10.1	6.9	8.4	7.6	8.7
Blocked nose/hay fever >4 days a week last 12 months	18.0	19.1	17.4	18.3	16.9	20.3	16.3	17.3
Woken by chest tightness/cough last 3 months	19.4	19.1	19.2	22.4	19.2	20.1	18.6	17.8
Restricted Activity Days due to respiratory symptoms last 3 months	10.9	10.3	10.2	10.7	11.6	12.4	10.1	9.2

Figures




Figure 1a. Map of Sweden showing study regions and main cities, response rates, and areas in which enhanced participant sampling in the Swedish Environmental Survey of 2007 took place
Figure 1b. Map of Sweden showing modelled regional background PM₁₀ (long-range transport particle) concentration




Figure 2. Distribution of modelled outdoor levels of total PM₁₀ and of different particle fractions at the home addresses of the study population in the Swedish Environmental Survey of 2007




Figure 3. Adjusted* associations between different PM fractions and self-reported current respiratory health outcomes in the Swedish Environmental Survey of 2007

* One-pollutant models adjusted for individual confounders gender, age, educational level, smoking, passive smoking, mould and moist exposure in the home and country of birth and the contextual confounders proportion of highly educated and unemployed people in the municipality.

Figure legends

Figure 1a. Map of Sweden showing study regions and main cities, response rates, and areas in which enhanced participant sampling in the Swedish Environmental Survey of 2007 took place

Figure 1b. Map of Sweden showing modelled regional background PM₁₀ (long-range transport particle) concentration

Figure 2. Distribution of modelled outdoor levels of total PM₁₀ and of different particle fractions at the home addresses of the study population in the Swedish Environmental Survey of 2007

Figure 3. Adjusted* associations between different PM fractions and self-reported current respiratory health outcomes in the Swedish Environmental Survey of 2007

* One-pollutant models adjusted for individual confounders gender, age, educational level, smoking, passive smoking, mould and moist exposure in the home and country of birth and the contextual confounders proportion of highly educated and unemployed people in the municipality.