



Early View

Original article

Incidence of pulmonary hypertension and determining factors in patients with systemic sclerosis

Gerry Coghlan, Matthias Wolf, Oliver Distler, Christopher P Denton, Martin Doelberg, Satenik Harutyunova, Alberto M Marra, Nicola Benjamin, Christine Fischer, Ekkehard Grünig

Please cite this article as: Coghlan G, Wolf M, Distler O, *et al*. Incidence of pulmonary hypertension and determining factors in patients with systemic sclerosis. *Eur Respir J* 2018; in press (<https://doi.org/10.1183/13993003.01197-2017>).

This manuscript has recently been accepted for publication in the *European Respiratory Journal*. It is published here in its accepted form prior to copyediting and typesetting by our production team. After these production processes are complete and the authors have approved the resulting proofs, the article will move to the latest issue of the ERJ online.

Copyright ©ERS 2018

Incidence of pulmonary hypertension and determining factors in patients with systemic sclerosis

Gerry Coghlan^{1*}, Matthias Wolf^{2*}, Oliver Distler⁴, Christopher P Denton⁵, Martin Doelberg⁶, Satenik Harutyunova², Alberto M Marra³, Nicola Benjamin², Christine Fischer⁷ and Ekkehard Grünig^{2,8}

¹Cardiology Department, Royal Free Hospital, London; ²Centre for Pulmonary Hypertension, Thoraxclinic at the University Hospital Heidelberg; ³IRRC S.D.N, Naples Italy; ⁴Department of Rheumatology, Faculty of Medicine, University of Zurich; ⁵ Centre of Rheumatology, Royal Free Hospital, London; ⁶Actelion Pharmaceuticals Ltd., Gewerbestrasse 16, CH-4123 Allschwil, Switzerland; ⁸ Translational Lung Research Center (TLRC) of the German Center for Lung Research (DZL), Heidelberg;

* equally contributed

Corresponding author:

Prof. Gerry Coghlan, MD
Cardiology Department, Royal Free Hospital,
Pond Street
London
NW3 2QG
UK.
Phone number: 020 7794 0500 ext 33614/38648/31188
Mail: gerry.coghlan@nhs.net

Summary: This is the first prospective study indicating that borderline elevation of mPAP is associated with higher incidence of PH in high risk SSc-patients using systematic recatheterisation.

Abstract

Objective: The objective of this study was to evaluate the incidence of pulmonary hypertension (PH) and determining factors in patients with systemic sclerosis (SSc) and a DLCO < 60% predicted.

Methods: In this bicentric, prospective cohort study, patients with SSc were assessed at baseline and after 3 years clinically including right heart catheterization (RHC). Analysis of determining factors for development of PH was performed using univariate and multivariate analysis.

Results: Ninety-six patients with mean pulmonary artery pressure (mPAP) < 25 mmHg at baseline were followed 2.95±0.7 (median 3) years. Seventy-one had a second RHC; 18 of the 71 patients (25.3%) developed PH, 5 (7%) a SSc-associated pulmonary arterial hypertension. For patients with mPAP between 21 and 24 mmHg at baseline, the likelihood of presenting with PH as opposed to normal pressures on follow-up was significantly higher (p=0.026). Pulmonary vascular resistance, tricuspid regurgitation velocity, diffusion capacity and size of inferior vena cava at baseline were independent predictors for development of PH during follow-up.

Conclusion: In a selected cohort of SSc patients with a DLCO < 60%, pulmonary pressures appear to rise progressively during follow up. In this population using prospective RHC during follow-up it was possible to identify manifest PH in almost 25% of 44 patients. Therefore, regular clinical assessment including RHC might be useful in SSc-patients.

Most important findings: In a selected cohort of SSc patients pulmonary pressures appear to rise progressively, leading to a development of manifest PH in 25% within 3 years.

Keywords: Systemic sclerosis, pulmonary hypertension, risk stratification

Introduction

Pulmonary hypertension (PH) is a common complication of systemic sclerosis (SSc) which can occur at any stage of the disease and has been observed in 15-27% of symptomatic patients and 8-12% of asymptomatic patients using right heart catheterization (RHC) for screening [1, 2]. If no other underlying disease such as heart or lung disease is the cause of PH, the disease is classified as SSc associated pulmonary arterial hypertension (SSc-PAH). Three-year survival for untreated SSc-PAH patients has been estimated to be 56% compared to 91% in those patients without PAH [3, 4]. At PAH-diagnosis >85% of SSc-patients are already in advanced stages of the disease (WHO functional class (FC) III and IV) [3]. Today, 10 PAH-targeted drugs are available for these patients [5], which have already been shown to improve symptoms, exercise capacity and outcome. Therefore, an early diagnosis of PH/PAH is essential in SSc-patients.

The diagnosis of PAH is defined by a mean pulmonary arterial pressure ≥ 25 mmHg at rest, a pulmonary arterial wedge pressure ≤ 15 mmHg and a pulmonary vascular resistance > 3 wood units, measured by right heart catheterization [6]. Normal mean pulmonary arterial pressures at rest are 14 ± 3 mmHg with an upper limit of approximately 20 mmHg [6, 7]. According to the current guidelines, the clinical significance of a mean pulmonary arterial pressure between 21 and 24 is not known [5].

It is recommended, that patients who are at high risk to develop pulmonary hypertension, e.g. patients with connective tissue disease (CTD), who present with mean pulmonary arterial pressure values within this range, should be carefully monitored [6]. Recent data in SSc-patients have shown that pulmonary arterial pressures of 21-24 mmHg lead to decreased exercise capacity, higher hospitalization and mortality [8-10]. In retrospective studies SSc-patients with mean PAP between 21- and 24 mmHg were more likely to develop PAH than patients with normal pulmonary arterial pressure [11, 12].

In 2014 the DETECT-Algorithm supplied the first evidence-based approach to early detection of SSc-PAH [13]. Visovatti et al. characterized borderline pulmonary arterial pressures as an individual subgroup of systemic sclerosis in a representative post-hoc analysis of the DETECT study cohort and hypothesized that this is an intermediate stage between normal pulmonary arterial pressures and PAH [14].

Determinants of PAH in SSc have already been investigated in several studies [15-19]. In a prospective cohort study Allanore et al. found DLCO/VA and NT-proBNP as being prognostically relevant for the development of PAH in SSc [17]. An association of SSc-PAH with low DLCO has been confirmed by several cohort studies [15, 16, 18, 19]. These studies were however mostly retrospective and did not include a systematic assessment of hemodynamics by right heart catheterization in all patients.

In the DETECT study, echocardiography at rest alone missed about 50% of PH diagnoses. Therefore, a study with a systematic assessment of hemodynamics by right heart catheterization of all patients during follow-up is needed to assess the true incidence and determinants of PAH in SSc.

The aim of this study was to assess the incidence of PH in SSc-patients, characterisation of the clinical course of the patients and investigation of determining factors for PH during follow-up. Furthermore, a specific focus was set on the clinical course of patients who presented with borderline pulmonary arterial pressures at baseline.

Material and methods

Study population and design

Patients who were included in the DETECT study in London and Heidelberg who did not have PH at initial screening using right heart catheterization were systematically followed and reassessed after 3 years. In addition, each centre recruited further 10 DETECT eligible patients without PH, who agreed to follow up.

For study inclusion patients ≥ 18 years of age were considered. SSc was diagnosed according to American College of Rheumatology criteria and a duration > 3 years of non-Raynaud symptoms or mixed connective tissue disease > 3 years was required [20]. SSc-patients receiving endothelin receptor antagonists or other targeted PAH-therapy were not included.

Clinical examinations at baseline and after 3 years comprised of medical history, vital signs, lung function, diffusion capacity, 6-minute walking distance, echocardiography, laboratory including NTproBNP and right heart catheterization. Right heart catheterization was performed according to the current guidelines [5].

After the final assessment including the second right heart catheterization, patients were followed by their hospital visits or contacted via phone for survival analysis.

Significant lung disease was evaluated by lung function test and high resolution computed tomography. Lung involvement of SSc was considered significant when FVC $<60\%$ or HRCT showed severe fibrosis or when FVC was 60%-70% and HRCT was „not available“ or fibrosis „moderate-severe“ or in case of other lung diseases apart from fibrosis by clinical decision of the treating physician. In the case of suspected coronary artery disease and in patients with elevated wedge pressures, patients were referred for left heart catheterization.

The study was conducted in accordance with the declaration of Helsinki. All patients gave written informed consent to the study. The study was approved by the ethics committee of the University of Heidelberg and London which were based on ethics committee approvals for the DETECT-study that has been registered on clinicaltrials.gov (no. NCT00706082).

Statistical analysis

Data were analysed by two statisticians (CF and NB). Values are presented as mean \pm standard deviation or n and percent, respectively. Baseline and follow-up characteristics of patients with borderline (21-24 mmHg) vs. normal (<21 mmHg) mPAP at baseline were compared using the 2*3 Chisquare test with two degrees of freedom. Individual changes during time were analysed by Wilcoxon signed rank test.

Analysis of determining factors of PH was performed by a two-step approach of Pearson regression analysis including univariate analysis for variable selection as first and multivariate stepwise forward procedure with centre as fixed factor as second step. Parameters for univariate analysis were selected according to clinical significance. For uni- and multivariate analysis, only parameters with more than 80% valid values were considered for the analysis.

Analysis of survival was performed by Kaplan-Meier method. The date of initial screening served as baseline date. Patients were regarded as censored at their last date of contact with the study team. The end point for survival was met by death of any cause or lung transplantation. P-values <0.05 were considered statistically significant.

Results

Ninety-six patients (81 female, 75% limited cutaneous SSc, 66% WHO functional class \geq II) with 48 patients from each centre were included in the baseline analysis. In 83 (86.5%) patients clinical follow-up assessment after 2.95 ± 0.7 (median 3) years was performed. Assessment of hemodynamics by right heart catheterization was conducted in 71 (74%) patients during follow-up. Twelve patients refused invasive assessment, of whom one was pregnant at the time of the 3-year follow-up and one had newly diagnosed lung cancer. None of these patients showed clinical signs of PH. Thus, our final study group consisted of 71 patients who were assessed by a second RHC within follow-up. Lung involvement of SSc developed in 14 patients during the course of the study, 11 patients had a FVC $<60\%$, three patients showed FVC 60%-70% and HRCT „moderate-severe“ lung disease. Further 9 patients were considered as significant lung disease according to the treating physician. Patient characteristics at baseline are given in Table 1. Extended description of all patients at baseline and in several subgroups is given in the supplementary tables.

Incidence of pulmonary hypertension

In 18 patients (25.3%, 95% CI: 15.7%-37.1%) pulmonary hypertension with an mPAP ≥ 25 mmHg was detected during follow-up. For patients with mPAP between 21 and 24 mmHg at baseline, the likelihood of presenting with PH as opposed to normal pressures on follow-up was significantly higher ($p=0.026$; Figure 1). The incidence for PH in the cohort of 71 patients

who had a second right heart catheterization was 6.11/100 patient years (95% CI 3.67/100 – 9.5/100). Of the 18 patients with PH at the second RHC, 5 had PH due to left heart disease, 8 due to lung disease. In 5 patients (7%, 95% CI 2.3%-15.7%) SSc-APAH was diagnosed during follow-up.

Progression of hemodynamics and clinical parameters during follow-up

The study cohort showed a significant worsening in 6-minute walking distance, NTproBNP-levels, lung function parameters (forced vital capacity, FEV1), diffusion capacity (DLCO, DLCO VA, DLCO %, DLCO VA%), echocardiography (Tricuspid regurgitation velocity/systolic pulmonary artery pressure) and invasive hemodynamics (mPAP, PVR, Table 2). During the course of the study, mean RAP significantly increased by 1.3 ± 3.5 mmHg ($p=0.001$). Change in RAP (baseline to follow-up) between normal and borderline patients did not significantly differ ($p=0.076$). The rate of progression to PAH was 3 of 21 (14%) with mPAP 21 mmHg - 24 mmHg at baseline versus 2 of 50 (4%) with normal mPAP at baseline. When looking at PH, the rate of progression was 7 of 21 (33%) for patients with mPAP 21 mmHg – 24 mmHg at baseline and 11 of 50 (22%) for patients with normal mPAP. In this population of SSc patients with a DLCO < 60%, the change of mPAP from baseline to 3 years did not significantly differ between patients presenting with normal mPAP ($+4.26 \pm 6.01$ mmHg) and those with borderline pressures at baseline ($+2.81 \pm 3.98$ mmHg).

One outlier was detected in the NTproBNP values, probably due to measurement errors. This patient with NTproBNP baseline value of 7000 ng/mL developed lung cancer within the study period and showed normal right ventricular function at baseline, creatinine of 1.15 mg/dL and uric acid of 4.0 mg/dL. As both right ventricular function and renal function do not explain this value, the NTproBNP was excluded from the analysis. Within the whole cohort, NTproBNP showed a significant increase ($p=0.005$ Wilcoxon rank test) throughout the study. The increase in NTproBNP did not significantly differ between patients with normal mPAP at baseline and those with pressures between 21 and 24 mmHg (<21 mmHg 195.9 ± 1199.5 median 13 vs. 21-24 mmHg 168.8 ± 404.0 mmHg median 42.5).

Comparison of mPAP < 21 mmHg and 21-24 mmHg at baseline

Patients presenting with mPAP between 21 and 24 mmHg at baseline showed significantly lower 6WMD, DLCO %, cardiac output and significantly higher tricuspid regurgitation velocity, systolic pulmonary arterial pressure, transpulmonary gradient and pulmonary vascular resistance both at baseline and during follow-up (all $p<0.05$; Table 3). Furthermore, lung function parameters at baseline were significantly worse in patients with mPAP 21-24 mmHg for forced vital capacity, forced expiratory volume in one second (FEV1), FEV1%, total lung capacity % and residual volume % (all $p<0.05$). For some parameters baseline values did not

differ, however at follow-up right atrial area was significantly larger ($p=0.037$), and tricuspid annular plane systolic excursion showed significantly lower values ($p=0.004$) in patients with mPAP 21-24 mmHg compared to mPAP <21 mmHg at baseline.

Determining factors of mPAP during follow-up

Results of univariate and multivariate analyses are given in Table 4. High pulmonary vascular resistance at baseline was independent predictor of the development of PH during follow-up ($p=0.002$, $r=0.460$). When only parameters of noninvasive assessments were included in the analysis elevated tricuspid regurgitation velocity measured by echocardiography, low diffusion capacity and enlarged size of inferior vena cava were further independent predictors of PH during follow-up (final model $p<0.001$).

Prognostic factors of survival

Eight patients died during follow-up due to the following reasons: pulmonary fibrosis ($n=2$), PH ($n=2$; 1 PAH, 1 postcapillary PH), cancer ($n=2$), primary biliary cholangitis ($n=1$), left heart failure ($n=1$). While the earliest death occurred after 1.0 year, the latest death occurred after 5.6 years of follow-up ($m = 3.2$ years, $M = 3.1$). One further patient with lung cancer was lost to follow-up three years after baseline.

Survival was not significantly different between patients with mPAP of 21-24 mmHg at baseline compared to patients with mPAP <21 mmHg ($p=0.217$, Figure 2a). While survival curves show congruency in patients with and without significant lung disease in the beginning, patients presenting with significant lung disease at baseline showed an impaired survival compared to patients without significant lung disease after >40 months ($p=0.029$, Figure 2b).

Discussion

This is the first prospective study to evaluate incidence and determining factors of pulmonary hypertension (PH) in patients with systemic sclerosis (SSc) using a systematic screening assessment including right heart catheterisation at baseline and after 3 years. The high incidence of PH (25.3%) and PAH (7%) within this time suggests that it is useful to perform regular clinical assessment with a low threshold for RHC in at risk SSc-patients. Likewise, the development of cardiac and pulmonary diseases should be monitored with particular attention, as in 5 patients, significant left heart disease and in 8 patients significant lung disease developed simultaneously to the progression to PH within the 3 years study period. In our study on average pulmonary pressure tended to rise over time in this population. High pulmonary vascular resistance at baseline, elevated tricuspid regurgitation velocity, low diffusion capacity and enlarged size of inferior vena cava were independent predictors for the development of PH during follow-up. This provides further evidence that borderline pulmonary arterial pressure is a possible intermediate stage in the development of pulmonary hypertension.

Incidence of PH/PAH

The incidence for PH in our cohort was 6.11/100 patient years, when only entering the 71 RHC-controlled patients into calculation. Among those the incidence of PAH was 7% similar to the findings of Valerio et al. where progression to PAH for all patients was 8.3% after 3 years. Of note, in the Valerio et al. study patients with pulmonary fibrosis were excluded from follow up, while in the DETECT cohort, patients with mild to moderate pulmonary fibrosis were included.

Our results show a higher rate of development of PH when compared with two further previous studies that analyzed the incidence of PH [16, 21]. The estimated incidence of PH over a period of 3 years, which was observed in a French nationwide study, was 1.37cases/100 patient years; incidences did not differ between PAH and postcapillary PH [21]. In an Italian study, PH incidence was 1.85/100 patient years [16]. In both studies, only patients who presented with suspected PH by clinical presentation or TRV were selected for RHC. In a more recent study Kovacs et al. reported an incidence of 0.75/100 patient years [22]. In all of these studies right heart catheterisation was only performed in those suspected clinically or on non-invasive investigation of having developed PAH.

In contrast to Hachulla et al., Ludici et al. and Kovacs et al. we used a systematic assessment via right heart catheterization in all patients, capturing all incident cases.

In addition, our cohort was preselected for possible PAH, as patients with impaired DLCO were selected. Mean DLCO % was 73.2 ± 18 in Hachulla et al. and 71 ± 21 in Ludici et al. and 82.2% (range 64.5-93.9) in Kovacs et al. [22], at baseline, while our cohort had a DLCO % of 48.9 ± 10.8 [16, 21]. The low DLCO appears to be a major reason for the higher apparent

incidence of pulmonary hypertension in our cohort. A low DLCO may indicate the need to perform closer clinical and invasive follow-up in patients with SSc.

Comparison of mPAP groups – are “borderline” pulmonary pressure an interim stage?

In a retrospective analysis of the DETECT cohort patients with borderline pulmonary arterial pressures showed significantly higher NT-proBNP, larger left atrium diameter, and greater tricuspid regurgitation velocity than patients with normal pulmonary hemodynamics [14]. 6MWD was not significantly different in this cohort [14].

In our study patients with borderline pulmonary arterial pressures at baseline showed significantly lower 6MWD, DLCO %, cardiac output, higher TRV, sPAP, TPG and PVR at baseline and follow-up examination. TAPSE was significantly lower and right atrial area significantly larger at follow-up. These findings are consistent with two studies that reported lower exercise capacity among patients with borderline pulmonary arterial pressure and suggested borderline PH as being indicative of early cardiopulmonary impairment (9,11).

In our cohort, patients with mPAP between 21 and 24 mmHg showed significantly poorer lung function at baseline than patients with mPAP <21 mmHg. This suggests pulmonary comorbidity is prevalent among those with mildly elevated pressures as shown in the PHAROS registry, reporting a higher prevalence of pulmonary fibrosis and abnormal lung physiology in patients with mPAP 21-24 mmHg [23]. Kovacs et al. also described a higher prevalence of cardiac comorbidity and decreased lung function [9] in patients with borderline pulmonary arterial pressures. Thus, the nature of the PH identified among populations during follow-up may also depend on the rigor with which cardiac and pulmonary co-morbidity were excluded.

Determining factors of developing PH

A reduction of DLCO is a frequent finding in systemic sclerosis, and in PH [24]. Compared to other PAH subgroups CTD-APAH patients showed lower DLCO [25, 26]. In our study DLCO/VA% was a significant predictor of developing PH along with enlarged size of inferior vena cava and tricuspid regurgitation velocity, when only noninvasive parameters were taken into account (final model $p < 0.001$). Nevertheless the effect size was small. Our findings are consistent with several previous studies who confirmed a strong association of DLCO and SSc-aPAH [15, 16, 18, 19]. However these studies were mostly retrospective and partially based diagnosis on echocardiography or did not use systematic right heart catheterization of all patients.

Mukerjee et al. found the relationship between mPAP and DLCO to be weak and suggested DLCO being an indicator of advanced rather than early PH as had been suggested by Steen et al. [19, 27].

In analysis of echocardiographic parameters only, tricuspid regurgitation velocity and size of inferior vena cava showed to be significant predictors. TRV has already been identified as independently associated factor in one study [14], Inferior vena cava has not previously been reported as predictor of mPAP or PH.

Progression of hemodynamics, regardless of the baseline stage (mPAP group)

Our study cohort showed a significant worsening in lung function parameters (forced vital capacity, FEV1), diffusion capacity (DLCO, DLCO VA, DLCO %, DLCO VA%), 6-minute walking distance, echocardiography (sPAP / TR-jet) and invasive hemodynamics (mPAP, PVR) during the course of the study. The change of mPAP from baseline to 3 years did not significantly differ between patients presenting with normal mPAP ($+4.26 \pm 6.01$ mmHg) and those with borderline pressures at baseline ($+2.81 \pm 3.98$ mmHg).

Our patients showed an increase in mPAP of 3.8 ± 5.5 mmHg during a three year period, which was also observed in a recent study with 1.1 mmHg/year [12]. This supports previous observations [24] that patients with a reduced DLCO will tend to show worsening of pulmonary haemodynamics over time, however without a catheter-based study of patients with normal gas transfers, we cannot be certain that this is not a general phenomenon among patients with SSc. The data from Kovacs et al. [22] suggests that in patients with a normal DLCO (mean 82%), progressive elevation of pulmonary pressures is not likely to occur, since in that study among those selected for repeat catheterisation no trend toward increasing pressures was observed.

Limitations

Due to the DETECT inclusion criteria, this cohort is preselected for SSc-patients with $DLCO < 60\%$, which can limit its generalisability to an unselected SSc-population. In the analysis of determining factors the study centre was included as fixed factor to take centre effects into account. However, we cannot rule out difference between centres as a contributor to the findings. Right heart catheterization was performed in only 71 out of 96 patients (74%) after 3 years. We do not know, whether the other patients developed pulmonary hypertension within three years. However, 83 patients (87%) were assessed during follow-up by non-invasive assessments. In those patients who were not assessed by right heart catheterization, no clinical signs of pulmonary hypertension were detected. The size of the cohort does not allow independent assessment of the rate of progression to PH (7 of 21 with mPAP 21 -24, vs. 11 of 50 with normal mPAP at baseline).

Conclusion

The results of this prospective study performing RHC at baseline and during follow-up in patients with SSc and reduced gas transfer indicate that progressive elevation of pulmonary pressure occurs in these patients over time. This would be expected to translate into an increased risk of PH and PAH in this population. As of the 18 patients with PH at the second RHC, 5 had PH due to left heart disease (27.8%) and 8 due to lung disease (44.4%), concomitant diseases have to be taken into account during clinical follow-up. We also provide further evidence that borderline pulmonary arterial pressure is a possible intermediate stage in the development of pulmonary hypertension. Using RHC during follow-up assessment it was possible to identify manifest PH in almost 25% of patients, PVR was an independent risk factor to develop manifest disease. Therefore, it seems to be useful to perform regular clinical assessment including RHC in SSc-patients with reduced gas transfer until more reliable noninvasive tools are developed.

Acknowledgements

Actelion Pharmaceuticals Ltd was responsible for designing the DETECT study protocol under the leadership of a company-funded, yet independent, Study Scientific Committee. The authors would like to thank Actelion Pharmaceuticals Ltd, all members of the DETECT Steering Committee and all patients involved in this study. Furthermore, we thank the patients' organization Skleroderma Liga eV for their support of the study.

Support statement.

The study was funded by a grant of Actelion.

References

1. Mukerjee D, St George D, Coleiro B, Knight C, Denton CP, Davar J, Black CM, Coghlan JG. Prevalence and outcome in systemic sclerosis associated pulmonary arterial hypertension: application of a registry approach. *Ann Rheum Dis* 2003; 62(11): 1088-1093.
2. Wigley FM, Lima JA, Mayes M, McLain D, Chapin JL, Ward-Able C. The prevalence of undiagnosed pulmonary arterial hypertension in subjects with connective tissue disease at the secondary health care level of community-based rheumatologists (the UNCOVER study). *Arthritis Rheum* 2005; 52(7): 2125-2132.
3. Hachulla E, Carpentier P, Gressin V, Diot E, Allanore Y, Sibilia J, Launay D, Mouthon L, Jegou P, Cabane J, de Groote P, Chabrol A, Lazareth I, Guillemin L, Clerson P, Humbert M, Itiner AIRSSI. Risk factors for death and the 3-year survival of patients with systemic sclerosis: the French ItinerAIR-Sclerodermie study. *Rheumatology (Oxford)* 2009; 48(3): 304-308.
4. Nihtyanova SI, Schreiber BE, Ong VH, Rosenberg D, Moinzadeh P, Coghlan JG, Wells AU, Denton CP. Prediction of pulmonary complications and long-term survival in systemic sclerosis. *Arthritis Rheumatol* 2014; 66(6): 1625-1635.
5. Galie N, Humbert M, Vachiery JL, Gibbs S, Lang I, Torbicki A, Simonneau G, Peacock A, Vonk Noordegraaf A, Beghetti M, Ghofrani A, Gomez Sanchez MA, Hansmann G, Klepetko W, Lancellotti P, Matucci M, McDonagh T, Pierard LA, Trindade PT, Zompatori M, Hoeper M, Aboyans V, Vaz Carneiro A, Achenbach S, Agewall S, Allanore Y, Asteggiano R, Paolo Badano L, Albert Barbera J, Bouvaist H, Bueno H, Byrne RA, Carerj S, Castro G, Erol C, Falk V, Funck-Brentano C, Gorenflo M, Granton J, Iung B, Kiely DG, Kirchhof P, Kjellstrom B, Landmesser U, Lekakis J, Lionis C, Lip GY, Orfanos SE, Park MH, Piepoli MF, Ponikowski P, Revel MP, Rigau D, Rosenkranz S, Voller H, Luis Zamorano J. 2015 ESC/ERS Guidelines for the diagnosis and treatment of pulmonary hypertension: The Joint Task Force for the Diagnosis and Treatment of Pulmonary Hypertension of the European Society of Cardiology (ESC) and the European Respiratory Society (ERS): Endorsed by: Association for European Paediatric and Congenital Cardiology (AEPC), International Society for Heart and Lung Transplantation (ISHLT). *Eur Heart J* 2016; 37(1): 67-119.
6. Hoeper MM, Bogaard HJ, Condliffe R, Frantz R, Khanna D, Kurzyna M, Langleben D, Manes A, Satoh T, Torres F, Wilkins MR, Badesch DB. Definitions and diagnosis of pulmonary hypertension. *J Am Coll Cardiol* 2013; 62(25 Suppl): D42-50.
7. Kovacs G, Berghold A, Scheidl S, Olschewski H. Pulmonary arterial pressure during rest and exercise in healthy subjects: a systematic review. *Eur Respir J* 2009; 34(4): 888-894.
8. Maron BA, Hess E, Maddox TM, Opatowsky AR, Tedford RJ, Lahm T, Joynt KE, Kass DJ, Stephens T, Stanislawski MA, Swenson ER, Goldstein RH, Leopold JA, Zamanian RT, Elwing JM, Plomondon ME, Grunwald GK, Baron AE, Rumsfeld J, Choudhary G. Association of Borderline Pulmonary Hypertension With Mortality and Hospitalization in a Large Patient Cohort: Insights From the VA-CART Program. *Circulation* 2016.
9. Kovacs G, Avian A, Tscherner M, Foris V, Bachmaier G, Olschewski A, Olschewski H. Characterization of patients with borderline pulmonary arterial pressure. *Chest* 2014; 146(6): 1486-1493.

10. Heresi GA, Minai OA, Tonelli AR, Hammel JP, Farha S, Parambil JG, Dweik RA. Clinical characterization and survival of patients with borderline elevation in pulmonary artery pressure. *Pulm Circ* 2013; 3(4): 916-925.
11. Kovacs G, Maier R, Aberer E, Brodmann M, Scheidl S, Troster N, Hesse C, Salmhofer W, Graninger W, Gruenig E, Rubin LJ, Olschewski H. Borderline pulmonary arterial pressure is associated with decreased exercise capacity in scleroderma. *American journal of respiratory and critical care medicine* 2009; 180(9): 881-886.
12. Valerio CJ, Schreiber BE, Handler CE, Denton CP, Coghlan JG. Borderline mean pulmonary artery pressure in patients with systemic sclerosis: transpulmonary gradient predicts risk of developing pulmonary hypertension. *Arthritis Rheum* 2013; 65(4): 1074-1084.
13. Coghlan JG, Denton CP, Grunig E, Bonderman D, Distler O, Khanna D, Muller-Ladner U, Pope JE, Vonk MC, Doelberg M, Chadha-Boreham H, Heinzl H, Rosenberg DM, McLaughlin VV, Seibold JR, group Ds. Evidence-based detection of pulmonary arterial hypertension in systemic sclerosis: the DETECT study. *Ann Rheum Dis* 2014; 73(7): 1340-1349.
14. Visovatti SH, Distler O, Coghlan JG, Denton CP, Grunig E, Bonderman D, Muller-Ladner U, Pope JE, Vonk MC, Seibold JR, Torres-Martin JV, Doelberg M, Chadha-Boreham H, Rosenberg DM, McLaughlin VV, Khanna D. Borderline pulmonary arterial pressure in systemic sclerosis patients: a post-hoc analysis of the DETECT study. *Arthritis Res Ther* 2014; 16(6): 493.
15. Hsu VM, Chung L, Hummers LK, Wigley F, Simms R, Bolster M, Silver R, Fischer A, Hinchcliff ME, Varga J, Goldberg AZ, Derk CT, Schiopu E, Khanna D, Shapiro LS, Domsic RT, Medsger T, Mayes MD, Furst D, Csuka ME, Molitor JA, Alkassab F, Steen VD. Development of pulmonary hypertension in a high-risk population with systemic sclerosis in the Pulmonary Hypertension Assessment and Recognition of Outcomes in Scleroderma (PHAROS) cohort study. *Semin Arthritis Rheum* 2014; 44(1): 55-62.
16. Iudici M, Codullo V, Giuggioli D, Riccieri V, Cuomo G, Breda S, Manfredi A, Iannace N, D'Alto M, Ghio S, Rossi R, Vizza CD, Caporali R, Valesini G, Ferri C, Valentini G. Pulmonary hypertension in systemic sclerosis: prevalence, incidence and predictive factors in a large multicentric Italian cohort. *Clin Exp Rheumatol* 2013; 31(2 Suppl 76): 31-36.
17. Allanore Y, Borderie D, Avouac J, Zerkak D, Meune C, Hachulla E, Mouthon L, Guillevin L, Meyer O, Ekindjian OG, Weber S, Kahan A. High N-terminal pro-brain natriuretic peptide levels and low diffusing capacity for carbon monoxide as independent predictors of the occurrence of precapillary pulmonary arterial hypertension in patients with systemic sclerosis. *Arthritis Rheum* 2008; 58(1): 284-291.
18. Chang B, Schachna L, White B, Wigley FM, Wise RA. Natural history of mild-moderate pulmonary hypertension and the risk factors for severe pulmonary hypertension in scleroderma. *J Rheumatol* 2006; 33(2): 269-274.
19. Steen V, Medsger TA, Jr. Predictors of isolated pulmonary hypertension in patients with systemic sclerosis and limited cutaneous involvement. *Arthritis Rheum* 2003; 48(2): 516-522.
20. Preliminary criteria for the classification of systemic sclerosis (scleroderma). Subcommittee for scleroderma criteria of the American Rheumatism Association Diagnostic and Therapeutic Criteria Committee. *Arthritis Rheum* 1980; 23(5): 581-590.

21. Hachulla E, de Groote P, Gressin V, Sibilia J, Diot E, Carpentier P, Mouthon L, Hatron PY, Jegou P, Allanore Y, Tiev KP, Agard C, Cosnes A, Cirstea D, Constans J, Farge D, Viallard JF, Harle JR, Patat F, Imbert B, Kahan A, Cabane J, Clerson P, Guillevin L, Humbert M, Itiner AIRSSG. The three-year incidence of pulmonary arterial hypertension associated with systemic sclerosis in a multicenter nationwide longitudinal study in France. *Arthritis Rheum* 2009; 60(6): 1831-1839.
22. Kovacs G, Avian A, Wutte N, Hafner F, Moazedi-Furst F, Kielhauser S, Aberer E, Brodmann M, Graninger W, Foris V, Olschewski A, Olschewski H. Changes in pulmonary exercise haemodynamics in scleroderma: a 4-year prospective study. *Eur Respir J* 2017; 50(1).
23. Bae S, Sagggar R, Bolster MB, Chung L, Csuka ME, Derk C, Domsic R, Fischer A, Frech T, Goldberg A, Hinchcliff M, Hsu V, Hummers L, Schioppa E, Mayes MD, McLaughlin V, Molitor J, Naz N, Furst DE, Maranian P, Steen V, Khanna D. Baseline characteristics and follow-up in patients with normal haemodynamics versus borderline mean pulmonary arterial pressure in systemic sclerosis: results from the PHAROS registry. *Ann Rheum Dis* 2012; 71(8): 1335-1342.
24. Steen VD, Graham G, Conte C, Owens G, Medsger TA, Jr. Isolated diffusing capacity reduction in systemic sclerosis. *Arthritis Rheum* 1992; 35(7): 765-770.
25. Jing ZC, Xu XQ, Badesch DB, Jiang X, Wu Y, Liu JM, Wang Y, Pan L, Li HP, Pu JL, Zhang ZL, Yang YJ. Pulmonary function testing in patients with pulmonary arterial hypertension. *Respir Med* 2009; 103(8): 1136-1142.
26. Ramjug S, Hussain N, Hurdman J, Billings C, Charalampopoulos A, Elliot CA, Kiely DG, Sabroe I, Rajaram S, Swift AJ, Condliffe R. Idiopathic and Systemic Sclerosis-Associated Pulmonary Arterial Hypertension: A Comparison of Demographic, Hemodynamic, and MRI Characteristics and Outcomes. *Chest* 2017; 152(1): 92-102.
27. Mukerjee D, St George D, Knight C, Davar J, Wells AU, Du Bois RM, Black CM, Coghlan JG. Echocardiography and pulmonary function as screening tests for pulmonary arterial hypertension in systemic sclerosis. *Rheumatology (Oxford)* 2004; 43(4): 461-466.

Table 1. Demographics of study cohort at baseline n=96

Characteristics	n (%) or mean (SD)	
gender		
female	81	(84.4%)
Demography		
Age, years	56.2	± 12.0
Body height, cm	164.5	± 8.6
Body weight, kg	68.1	± 14.5
BMI, kg/m ²	25.2	± 4.8
Vital Signs		
Blood Pressure, systolic, mmHg	117.8	± 17.4
Blood Pressure, diastolic, mmHg	72.0	± 10.8
Heart rate, /min	76.8	± 12.1
Systemic sclerosis characteristics		
Modified Rodnan Skin Score	11.9	± 8.8
Duration of SSc, months	11.5	± 9.6
Type of systemic sclerosis		
Diffuse cutaneous SSc	15	(15.6%)
Limited cutaneous SSc	71	(74.0%)
Mixed connective tissue disease	10	(10.4%)
WHO-functional class		
I	22	(22.9%)
II	33	(34.4%)
III	31	(32.3%)

BMI = Body Mass Index, SSc = Systemic Sclerosis, Type of systemic sclerosis as diagnosed by treating rheumatologist, Modified Rodnan Skin Score and WHO-Functional class as obtained in clinical examination.

Table 2. Clinical data at baseline and during follow-up

	baseline			follow-up			changes		p-value
	n	mean	SD	n	mean	SD	mean	SD	
6-MWD	91	403.2 ± 111.4		66	388 ± 125		-21.8 ± 79.3		0.039
NT-proBNP, pg/ml	95	216 ± 266		77	396 ± 1068		188 ± 1030		0.005
pulmonary function testing									
FVC, l	94	2.9 ± 1.0		75	2.7 ± 1.0		-0.1 ± 0.3		0.005
FVC, %	96	91.2 ± 23.7		76	89.2 ± 24.6		-1.1 ± 11.9		n.s.
FEV1, l	94	2.2 ± 0.8		74	2.0 ± 0.8		-0.1 ± 0.3		<0.001
FEV1, %	95	85.7 ± 22.9		75	82.6 ± 21.7		-1.3 ± 11.8		n.s.
DLCO,	93	7.5 ± 4.0		68	3.9 ± 1.5		-4.0 ± 4.0		<0.001
mmol/min/kPa									
DLCO, %	95	48.9 ± 10.8		70	46.3 ± 11.8		-3.1 ± 8.2		<0.001
DLCO/VA,	93	2.5 ± 1.5		69	1.5 ± 1.1		-1.3 ± 1.9		<0.001
mmol/min/kPa/l									
DLCO/VA, %	93	71.4 ± 16.9		69	67.7 ± 17.1		-5.1 ± 23.0		0.011
TLC, %	83	85.8 ± 22.3		70	89.7 ± 22.9		2.7 ± 11.4		n.s.
RV, %	84	85.8 ± 33.1		63	94.4 ± 34.2		5.5 ± 23.1		n.s.
echocardiography									
LA, mm	93	29.1 ± 6.4		72	30 ± 7		0.5 ± 6.4		n.s.
IVC, mm	82	14.0 ± 3.7		64	13 ± 6		-2.0 ± 6.8		n.s.
IVS, mm	92	10.1 ± 1.8		40	10 ± 2		0.0 ± 3.2		n.s.
RA, cm ²	89	12.1 ± 3.7		72	12.7 ± 4.0		0.2 ± 4.2		n.s.
RVD, mm	85	29.1 ± 5.6		42	31 ± 7		3.8 ± 7.6		0.01
RV, cm ²	88	14.5 ± 4.3		72	13.2 ± 3.7		-1.2 ± 4.1		0.043
LV-EDD, mm	92	44.1 ± 5.5		42	42 ± 8		-1.5 ± 8.5		n.s.
LV-ESD, mm	91	27.0 ± 5.1		42	27 ± 7		0.0 ± 7.0		n.s.
TRV, m/s	88	2.4 ± 0.4		63	2.6 ± 0.4		0.1 ± 0.5		0.019
TAPSE, mm	89	22.5 ± 4.6		77	22 ± 5		-0.4 ± 6.1		n.s.
sPAP, mmHg	87	29.1 ± 7.1		63	32 ± 9		3.1 ± 9.1		0.017
right heart catheterization									
mPAP, mmHg	96	17.1 ± 4.0		71	22 ± 6		3.8 ± 5.5		<0.001
PAWP, mmHg	96	8.5 ± 3.3		71	11 ± 3		1.9 ± 4.3		<0.001
TPG, mmHg	96	8.5 ± 2.9		71	9 ± 3		0.1 ± 2.6		n.s.
CO, l/min	96	5.3 ± 1.2		71	5.1 ± 1.1		-0.1 ± 0.9		n.s.
PVR, dynes	96	135 ± 55.4		71	182 ± 118		43.5 ± 96.1		0.001

SD = standard deviation, follow-up = follow-up after 3 years, 6-MWD = 6-minute walking distance, NTproBNP = N-terminal end of pro brain natriuretic peptide, FVC = forced vital capacity, FEV1 = forced expiratory volume in one second, DLCO = diffusion capacity of the lung for carbon monoxide, VA = alveolar volume, TLC = total lung capacity, RV = residual volume, LA = left atrium, IVC = inferior vena cava, IVS = interventricular septum, RA = right atrium, RVD = right ventricular diameter, RV = right ventricle, LV = left ventricle, EDD = end-diastolic diameter, ESD = end-systolic diameter, TRV = tricuspid regurgitation velocity, TAPSE = tricuspid annular plane systolic excursion, sPAP = systolic pulmonary arterial hypertension, mPAP = mean pulmonary arterial pressure, PAWP = pulmonary arterial wedge pressure, TPG = transpulmonary gradient, CO = cardiac output, PVR = pulmonary vascular resistance.

Table 3. Comparison of patients presenting with mean pulmonary arterial pressure <21 mmHg vs. 21-24 mmHg at baseline

	mPAP <21mmHg						mPAP 21-24mmHg						differences		
	baseline			follow-up			baseline			follow-up			p-value		
	n	mean	SD	n	mean	SD	n	mean	SD	n	mean	SD	baseline	follow-up	
6-MWD	69	431 ± 93		49	419 ± 107		22	317 ± 122		17	298 ± 131		<0.001 *	0.002	*
NT-proBNP, pg/ml	72	206 ± 263		54	401 ± 1243		23	245 ± 278		23	384 ± 452		0.255	0.181	
pulmonary function testing															
FVC, l	71	2.99 ± 1.00		56	2.76 ± 1.06		23	2.43 ± 0.75		19	2.41 ± 0.68		0.023 *	0.172	
FVC, %	72	92.90 ± 24.09		55	88.92 ± 25.69		24	86.02 ± 21.97		21	90.07 ± 21.92		0.233	0.963	
FEV1, l	71	2.39 ± 0.88		55	2.14 ± 0.87		23	1.78 ± 0.51		19	1.75 ± 0.49		0.004 *	0.113	
FEV1, %	72	88.43 ± 24.16		55	83.32 ± 23.25		23	77.02 ± 15.73		20	80.65 ± 16.90		0.050 *	0.545	
DLCO, mmol/min/kPa	70	7.26 ± 4.18		50	3.97 ± 1.01		23	8.06 ± 3.58		18	3.57 ± 2.41		0.228	0.003	*
DLCO, %	71	50.78 ± 9.56		50	49.44 ± 10.13		24	43.18 ± 12.54		20	38.55 ± 12.15		0.013 *	<0.001	*
DLCO/VA, mmol/min/kPa/l	70	2.35 ± 1.59		51	1.44 ± 1.09		23	2.93 ± 1.10		18	1.53 ± 1.31		0.061	0.280	
DLCO/VA, %	70	72.80 ± 16.55		51	69.14 ± 16.93		23	67.33 ± 17.64		18	63.65 ± 17.28		0.226	0.170	
TLC, %	63	88.80 ± 21.83		52	92.52 ± 21.68		20	76.21 ± 21.57		18	81.58 ± 24.96		0.028 *	0.070	
RV, %	63	91.56 ± 32.66		45	99.68 ± 33.38		21	68.55 ± 28.82		18	81.36 ± 33.68		0.002 *	0.084	
Echocardiography															
LA, mm	71	28.14 ± 6.48		52	30.07 ± 6.22		22	32.10 ± 4.95		20	30.45 ± 7.38		0.007 *	0.262	
IVC, mm	65	13.83 ± 3.61		47	12.50 ± 6.01		17	14.50 ± 3.96		17	12.59 ± 5.38		0.432	0.681	
IVS, mm	70	10.12 ± 1.81		23	9.66 ± 2.51		22	9.96 ± 1.80		17	10.79 ± 2.19		0.832	0.196	
RA, cm2	68	11.76 ± 3.37		52	12.15 ± 3.77		21	13.14 ± 4.64		20	14.11 ± 4.48		0.317	0.037	*
RVD, mm	65	29.89 ± 5.70		22	31.10 ± 7.41		20	26.36 ± 4.16		20	30.92 ± 5.68		0.003 *	0.830	
RV, cm2	68	14.25 ± 4.05		52	13.19 ± 3.59		20	15.19 ± 5.18		20	13.38 ± 3.94		0.495	0.692	
LV-EDD, mm	70	43.54 ± 5.58		23	41.51 ± 9.50		22	46.03 ± 4.98		19	42.86 ± 6.83		0.062	0.889	
LV-ESD, mm	69	26.64 ± 5.13		23	25.90 ± 7.84		22	28.18 ± 5.05		19	27.45 ± 5.51		0.238	0.486	
TRV, m/s	65	2.35 ± 0.36		23	2.48 ± 0.36		23	2.65 ± 0.30		16	2.88 ± 0.45		<0.001 *	0.003	*
TAPSE, mm	68	22.82 ± 3.92		23	23.02 ± 4.98		21	21.64 ± 6.37		22	19.75 ± 3.78		0.157	0.004	*
sPAP, mmHg	64	27.54 ± 6.78		23	30.16 ± 7.31		23	33.43 ± 6.27		16	38.97 ± 10.56		<0.001 *	0.003	*
right heart catheterization															
mPAP, mmHg	72	15.40 ± 3.03		50	20.12 ± 5.86		24	22.17 ± 0.96		21	24.95 ± 4.12		<0.001 *	<0.001	*
PAWP, mmHg	72	7.65 ± 2.99		50	10.74 ± 3.83		24	11.08 ± 2.67		21	11.24 ± 2.53		<0.001 *	0.407	

TPG, mmHg	72	7.69 ± 2.63	50	7.84 ± 2.22	24	10.83 ± 2.60	21	10.90 ± 2.43	<0.001 *	<0.001 *
CO, l/min	72	5.44 ± 1.17	50	5.24 ± 1.22	24	4.90 ± 1.13	21	4.86 ± 0.88	0.048 *	0.048 *
PVR, dynes	72	117.48 ± 39.04	50	151.48 ± 101.02	24	188.39 ± 63.37	21	253.70 ± 126.82	<0.001 *	<0.001 *
RAP, mmHg	72	3.76 ± 2.33	48	5.65 ± 2.89	24	5.00 ± 2.50	21	5.43 ± 2.23	0.03 *	0.76

* denotes statistically significant differences

* SD = standard deviation, follow-up = follow-up after 3 years, 6-MWD = 6-minute walking distance, NTproBNP = N-terminal end of pro brain natriuretic peptide,

* FVC = forced vital capacity, FEV1 = forced expiratory volume in one second, DLCO = diffusion capacity of the lung for carbon monoxide, VA = alveolar volume,

* TLC = total lung capacity, RV = right ventricle, LV = left ventricle, EDD = end-diastolic diameter, ESD = end-systolic diameter, TRV = tricuspid regurgitation velocity, TAPSE = tricuspid annular plane systolic excursion, sPAP = systolic pulmonary arterial hypertension, mPAP = mean pulmonary arterial pressure, PAWP = pulmonary arterial wedge pressure, TPG = transpulmonary gradient, CO = cardiac output, PVR = pulmonary vascular resistance.

Table 4

Baseline Parameters predictive of mean pulmonary arterial pressure during follow-up

Variable	n	p-value	pearson's R
Univariate Analysis			
Age	71	0.016	0.286
duration of systemic sclerosis	70	0.954	0.007
WHO functional class	70	0.485	0.084
Lung function			
FVC	69	0.051	-0.236
FEV1	69	0.017	-0.286
FEV1 %	70	0.057	-0.229
DLCO %	71	0.025	-0.265
DLCO/VA %	69	0.028	-0.265
NTproBNP	69	0.755	0.38
6-minute walking distance	66	0.097	0.206
Echocardiography			
Inferior vena cava	58	0.04	0.271
right atrial area	64	0.167	0.175
right ventricular area	63	0.828	0.028
Tricuspid regurgitation velocity	66	0.003	0.360
systolic pulmonary arterial pressure	65	0.004	0.351
Right heart catheterization			
mean pulmonary arterial pressure	71	0.001	0.402
Transpulmonary gradient	71	<0.001	0.430
pulmonary vascular resistance	71	<0.001	-0.456
Multivariate Analysis with centre as fixed factor			
Including invasive hemodynamics			
pulmonary vascular resistance	56	0.002	0.460
Only noninvasive parameters			
model 1 Tricuspid regurgitation velocity	56	0.003	0.439
model 2+ DLCO/VA %	56	0.046	0.512
model 3+ Inferior vena cava	56	0.02	0.577

WHO = World Health Organization, FVC = forced vital capacity, FEV 1 = forced expiratory volume in one second, DLCO = diffusion capacity of the lung for carbon monoxide, VA = alveolar volume, NTproBNP = N-terminal end of pro brain natriuretic peptide,

Figure Legends


Figure 1

The figure displays the clinical course and classification of the patients throughout the study. Distribution of patients during follow-up significantly differed between patients with baseline mPAP 21-24 mmHg and patients with mPAP <21 mmHg (mPAP<21mmHg: 11 manifest, 29 normal vs. mPAP 21-24mmHg: 7 manifest 1 normal; χ^2 $p<0.026$).


Figure 2

The two figures (a and b) show survival analyses of a) patients with mPAP 21-24 mmHg vs. mPAP <21 mmHg and b) patients with significant lung disease vs. no significant lung disease.


While mPAP at baseline did not affect survival, patients with significant lung disease presented with worse survival than patients without lung disease ($p=0.029$).



a) pulmonary arterial pressure at baseline



b) Significant lung disease



Patients at risk

mPAP 21-24mmHg	22	21	15	6
mPAP <21mmHg	66	62	41	11

Patients at risk

No lung disease	67	63	42	11
Lung disease	21	20	14	6

Supplementary table 1**Characteristics of borderline patients at baseline**

	n (%) or mean (SD)		median	Q25%-Q75%
gender				
female	19	79.2%		
Demography				
Age, years	61.0 ± 9.3		61.0	56.5 - 67.5
Body height, cm	160.8 ± 7.4		160.3	155.5 - 165.5
Body weight, kg	66.2 ± 13.3		64.9	57.0 - 72.4
BMI	25.5 ± 4.5		25.2	-
Vital Signs				
Blood Pressure, systolic, mmHg	118 ± 17		114	110 - 132
Blood Pressure, diastolic, mmHg	70 ± 11		68	60 - 80
Heart rate, /min	77.8 ± 12.9		77	72 - 80
Systemic sclerosis characteristics				
Modified Rodnan Skin Score	10 ± 8		8	6 - 12
Duration of SSc, months	11.2 ± 6.5		9.5	6.5 - 15.0
Type of systemic sclerosis				
Diffuse cutaneous SSc	6	25.0 %		
Limited cutaneous SSc	15	62.5 %		
Mixed connective tissue disease	3	12.5 %		
WHO-functional class				
I	1	4.2 %		
II	4	16.7 %		
III	12	50.0 %		

BMI = Body Mass Index, SSc = Systemic Sclerosis

Supplementary table 2

Characteristics of borderline patients at baseline developing and not developing PH

	borderline patients not developing PH						patients developing manifest PH						
	n	mean	SD	median	Q25%-Q75%		n	mean	SD	media n	Q25%-Q75%		p-value
6-MWD	15	320	± 111	375	231	- 393	7	311	± 153	364	198	- 472	
NTproBNP	14	170	± 457	37	-51	- 269	7	195	± 387	61	40	- 104	
pulmonary function													
FVC, l	17	2.23	± 0.7	2.2	1.8	- 2.5	6	3.0	± 0.6	3.1	2.8	- 3.3	*
FVC, %	17	84.45	± 20.0	88.3	67.2	- 95.9	7	89.8	± 27.6	82.9	71.0	- 121.0	
FEV ₁ , l	17	1.6	± 0.5	1.6	1.4	- 1.8	6	2.2	± 0.4	2.3	2.1	- 2.5	*
FEV ₁ , %	17	74.62	± 13.4	77.0	63.0	- 84.2	6	83.8	± 21.0	86.1	71.8	- 104.3	
DL _{CO} , mmol/min/kPa	17	7.1	± 3.4	7.5	4.6	- 10.0	6	10.8	± 2.8	10.9	8.5	- 13.4	
DL _{CO} , %	17	43.3	± 12.8	42.9	37.0	- 50.5	7	42.9	± 12.8	36.8	31.9	- 55.0	*
DL _{CO} /VA, mmol/min/kPa/l	17	3.0	± 1.2	3.1	2.6	- 3.6	6	2.7	± 0.5	2.5	2.5	- 3.0	
DL _{CO} /VA, %	17	69.3	± 18.7	70.7	61.4	- 78.0	6	61.7	± 14.0	60.7	52.3	- 69.7	
TLC, %	14	73.4	18.5	70.6	58.5	- 82.0	6	82.7	28.3	83.0	59.1	101.8	
RV, %	15	66.3	± 26.6	62.6	52.4	- 74.5	6	74.3	± 35.8	72.1	39.3	- 96.8	
Echocardiography													
LA, mm	16	31.7	± 4.9	32.0	27.3	- 34.6	6	33.2	± 5.4	31.5	30.0	- 33.0	
IVC, mm	12	14.9	± 4.2	15.6	12.3	- 18.7	5	13.4	± 3.3	13.0	12.1	- 13.1	
IVS, mm	16	10.2	± 1.6	10.6	9.8	- 11.1	6	9.3	± 2.4	9.0	8.0	- 10.7	
RA, cm ²	15	12.1	± 4.3	11.6	9.0	- 14.3	6	15.6	± 4.8	16.8	11.3	- 19.4	
RVD, mm	14	26.2	± 3.7	26.2	22.6	- 30.0	6	26.8	± 5.5	27.5	21.0	- 32.0	
RV, cm ²	14	13.3	± 4.9	12.5	10.6	- 15.3	6	19.6	± 2.7	19.0	18.5	- 21.0	*
LV-EDD, mm	16	44.5	± 4.3	44.8	42.5	- 46.9	6	50.0	± 4.8	51.0	46.0	- 54.1	*
LV-ESD, mm	16	27.0	± 4.1	26.0	25.0	- 29.5	6	31.3	± 6.4	30.3	29.0	- 35.0	
TRV, m/s	17	2.7	± 0.3	2.7	2.5	- 2.8	6	2.6	± 0.4	2.6	2.5	- 2.7	
TAPSE, mm	15	20.4	± 5.6	19.1	15.5	- 24.0	6	24.7	± 7.8	22.5	22.0	- 28.2	

sPAP, mmHg	17	33.5 ± 5.5	33.7	30.0 - 36.4	6	33.1 ± 8.7	32.4	29.2 - 34.2
right heart catheterisation								
mPAP, mmHg	17	22.0 ± 0.9	22.0	21.0 - 23.0	7	22.6 ± 1.1	22.0	22.0 - 24.0
PAWP, mmHg	17	10.9 ± 2.4	11.0	10.0 - 12.0	7	11.6 ± 3.4	12.0	8.0 - 15.0
TPG, mmHg	17	10.8 ± 2.3	12.0	9.0 - 12.0	7	11.0 ± 3.4	11.0	9.0 - 14.0
CO, l/min	17	4.9 ± 1.3	4.4	3.9 - 5.9	7	4.8 ± 0.6	4.9	4.4 - 5.2
PVR, dynes	17	190.2 ± 64.9	181.0	140.0 - 223.3	7	183.9 ± 64.3	169.2	146.9 - 231.0
RAP, mmHg	17	4.8 ± 2.4	4.0	3.0 - 3.46	7	5.6 ± 2.8	5.0	4.0 - 7.0

PH = pulmonary hypertension, Q25% - Q75% = 25% and 75% quantiles, 6-MWD = 6-minute walking distance, NTproBNP = N-terminal pro Brain Natriuretic Peptide, FVC = forced vital capacity, FEV1 = forced expiratory volume in one second, DLCO = diffusion capacity of the lung for carbon monoxide, VA = alveolar volume, TLC = total lung capacity, RV = residual volume, LA = left atrium, IVC = inferior vena cava, IVS = interventricular septum, RA = right atrium, RVD = right ventricular diameter, RV = right ventricle, LV = left ventricle, EDD = end-diastolic diameter, ESD = end-systolic diameter, TRV = tricuspid regurgitation velocity, TAPSE = tricuspid annular plane systolic excursion, sPAP = systolic pulmonary arterial hypertension, mPAP = mean pulmonary arterial pressure, PAWP = pulmonary arterial wedge pressure, TPG = transpulmonary gradient, CO = cardiac output, PVR = pulmonary vascular resistance.

Supplementary table 3
baseline characteristics, all patients - extended description

	n	mean	SD	median	Q25%-Q75%
6-MWD	91	403 ± 111		419	335 - 477
NT-proBNP, pg/ml	95	216 ± 266		124	67 - 256
pulmonary function testing					
FVC, l	94	2.85 ± 1.0		2.7	2.08 - 3.54
FVC, %	96	91.2 ± 23.7		90.9	73.2 - 110
FEV ₁ , l	94	2.2 ± 0.8		2.1	1.52 - 2.81
FEV ₁ , %	95	85.7 ± 22.9		84.2	68.3 - 101
DL _{CO} , mmol/min/kPa	93	7.5 ± 4.0		5.5	4.57 - 10.31
DL _{CO} , %	95	48.9 ± 10.8		51.1	41.3 - 57.1
DL _{CO} /VA, mmol/min/kPa/l	93	2.5 ± 1.5		2.54	1.11 - 3.4
DL _{CO} /VA, %	93	71.4 ± 16.9		69.7	60.2 - 82.9
TLC, %	83	85.8 ± 22.3		87.2	68.0 - 101.8
RV, %	84	85.8 ± 33.1		82.5	62.8 - 106.4
echocardiography					
LA, mm	93	29.1 ± 6.4		29	25.5 - 32.0
IVC, mm	82	14.0 ± 3.7		14	11.8 - 16.0
IVS, mm	92	10.1 ± 1.8		10	9.0 - 11.0
RA, cm ²	89	12.1 ± 3.7		11.8	9.3 - 14.0
RVD, mm	85	29.1 ± 5.6		30	26.0 - 33.0
RV, cm ²	88	14.5 ± 4.3		14.7	11.7 - 17.0
LV-EDD, mm	92	44.1 ± 5.5		44	40.5 - 47.0
LV-ESD, mm	91	27.0 ± 5.1		26	23.5 - 30.0
TRV, m/s	88	2.4 ± 0.4		2.41	2.2 - 2.6
TAPSE, mm	89	22.5 ± 4.6		22	19.8 - 25.0
sPAP, mmHg	87	29.1 ± 7.1		28.4	24.4 - 33.1
right heart catheterisation					
mPAP, mmHg	96	17.1 ± 4.0		17	15.0 - 20.5

PAWP, mmHg	96	8.5 ± 3.3	9	6.0 - 11.0
TPG, mmHg	96	8.5 ± 2.9	8	6.0 - 11.0
CO, l/min	96	5.3 ± 1.2	5.2	4.4 - 6.2
PVR, dynes	96	135.2 ± 55.4	124.2	100.1 - 163.0
RAP, mmHg	96	4.1 ± 2.4	4	2.0 - 5.0

PH = pulmonary hypertension, Q25% - Q75% = 25% and 75% quantiles, 6-MWD = 6-minute walking distance, NTproBNP = N-terminal pro Brain Natriuretic Peptide, FVC = forced vital capacity, FEV1 = forced expiratory volume in one second, DLCO = diffusion capacity of the lung for carbon monoxide, VA = alveolar volume, TLC = total lung capacity, RV = residual volume, LA = left atrium, IVC = inferior vena cava, IVS = interventricular septum, RA = right atrium, RVD = right ventricular diameter, RV = right ventricle, LV = left ventricle, EDD = end-diastolic diameter, ESD = end-systolic diameter, TRV = tricuspid regurgitation velocity, TAPSE = tricuspid annular plane systolic excursion, sPAP = systolic pulmonary arterial hypertension, mPAP = mean pulmonary arterial pressure, PAWP = pulmonary arterial wedge pressure, TPG = transpulmonary gradient, CO = cardiac output, PVR = pulmonary vascular resistance.

Supplementary table 4

Additional baseline characteristics of PH Groups

	PH left heart disease					PH lung disease				
	n	mean	SD	median	Q25%-Q75%	n	mean	SD	median	Q25%-Q75%
6-MWD	7	416 ± 188		445	394 - 560	10	392 ± 101		376	330 - 477
NT-proBNP, pg/ml	7	382 ± 111		256	92 - 310	10	115 ± 87		107	34 - 160
pulmonary function testing										
FVC, l	7	2.9 ± 0.9		3.1	2.6 - 3.7	9	2.7 ± 1.2		2.0	1.9 - 3.9
FVC, %	7	94.3 ± 33.3		102.0	77.2 - 113.2	10	77.1 ± 27.3		69.5	55.8 - 90.8
FEV ₁ , l	7	2.3 ± 0.7		2.6	1.9 - 2.8	9	2.4 ± 1.2		1.9	1.6 - 3.8
FEV ₁ , %	7	87.2 ± 25.5		92.2	85.3 - 100.9	9	77.3 ± 31.0		60.6	59.0 - 80.8
DL _{CO} , mmol/min/kPa	7	8.2 ± 4.1		4.8	4.7 - 12.7	9	6.0 ± 2.3		5.5	4.9 - 7.5
DL _{CO} , %	7	55.3 ± 2.1		55.1	54.5 - 56.5	10	38.6 ± 12.5		34.0	27.3 - 48.1
DL _{CO} /VA, mmol/min/kPa/l	7	1.9 ± 0.8		1.4	1.1 - 3.0	9	2.6 ± 2.3		2.5	1.0 - 2.5
DL _{CO} /VA, %	7	69.0 ± 14.7		65.1	57.8 - 85.3	9	60.6 ± 16.4		58.7	52.8 - 61.7
TLC, %	6	96.5 ± 30.9		96.5	83.0 - 123.1	8	66.8 ± 27.7		57.7	49.9 - 76.3
RV, %	6	11.4 ± 33.3		116.6	78.9 - 137.5	8	57.9 ± 33.2		52.3	37.3 - 78.9
echocardiography										
LA, mm	7	28.9 ± 6.9		30.0	20.0 - 32.0	9	26.8 ± 8.4		27.0	22.0 - 30.0
IVC, mm	7	13.0 ± 3.5		12.0	11.0 - 15.3	8	15.0 ± 3.7		15.6	11.6 - 18.5
IVS, mm	7	10.4 ± 0.7		9.0	8.5 - 10.0	9	10.1 ± 1.8		10.0	9.0 - 11.0
RA, cm ²	7	11.9 ± 3.4		11.6	8.8 - 13.0	9	12.6 ± 4.8		12.0	9.8 - 16.0
RVD, mm	5	31.6 ± 3.2		35.0	29.0 - 35.0	9	28.7 ± 4.8		28.0	26.0 - 33.0
RV, cm ²	7	13.7 ± 5.7		15.0	12.2 - 16.0	9	14.4 ± 4.9		14.9	11.5 - 18.0
LV-EDD, mm	7	43.8 ± 5.9		46.0	37.0 - 49.0	9	41.4 ± 6.7		42.6	39.0 - 43.0
LV-ESD, mm	7	28.3 ± 2.7		29.0	25.0 - 31.0	8	27.3 ± 6.3		25.6	23.5 - 29.0
TRV, m/s	6	2.5 ± 0.1		2.5	2.3 - 2.6	9	2.4 ± 0.4		2.3	2.2 - 2.6
TAPSE, mm	7	22.6 ± 5.7		23.0	17.0 - 27.0	9	21.2 ± 4.2		22.0	19.0 - 22.0

sPAP, mmHg	6	30.3 ± 2.5	29.8	26.2 - 31.6	9	28.2 ± 7.1	26.5	24.4 - 32.7
right heart catheterisation								
mPAP, mmHg	7	17.0 ± 4.5	20.0	13.0 - 20.0	10	18.5 ± 3.3	18.5	16.0 - 22.0
PAWP, mmHg	7	8.9 ± 5.1	8.0	6.0 - 12.0	10	8.9 ± 1.9	9.0	7.0 - 10.0
TPG, mmHg	7	7.3 ± 1.3	8.0	6.0 - 8.0	10	10.7 ± 3.6	11.0	10.0 - 13.0
CO, l/min	7	5.1 ± 0.8	5.0	4.4 - 5.6	10	5.0 ± 1.0	5.0	4.1 - 5.9
PVR, dynes	7	128.4 ± 29.5	111.0	105.7 - 167.4	10	161.3 ± 74.8	153.0	123.1 - 231.0
RAP, mmHg	7	4.7 ± 3.6	5.0	3.0 - 6.0	10	4.6 ± 2.2	4.5	4.1 - 5.0

PH left heart disease = pulmonary hypertension secondary to left heart disease, PH lung disease = pulmonary hypertension secondary to lung disease, Q25% - Q75% = 25% and 75% quantiles, 6-MWD = 6-minute walking distance, NTproBNP = N-terminal pro Brain Natriuretic Peptide, FVC = forced vital capacity, FEV1 = forced expiratory volume in one second, DLCO = diffusion capacity of the lung for carbon monoxide, VA = alveolar volume, TLC = total lung capacity, RV = residual volume, LA = left atrium, IVC = inferior vena cava, IVS = interventricular septum, RA = right atrium, RVD = right ventricular diameter, RV = right ventricle, LV = left ventricle, EDD = end-diastolic diameter, ESD = end-systolic diameter, TRV = tricuspid regurgitation velocity, TAPSE = tricuspid annular plane systolic excursion, sPAP = systolic pulmonary arterial hypertension, mPAP = mean pulmonary arterial pressure, PAWP = pulmonary arterial wedge pressure, TPG = transpulmonary gradient, CO = cardiac output, PVR = pulmonary vascular resistance.

Supplementary table 5

Baseline characteristic of patients according to follow-up (patients who did perform RHC vs. patients who only performed noninvasive follow-up)

	noninvasive follow-up						right heart catheterisation						P-value
	n	mean	SD	median	Q25%-Q75%		n	mean	SD	median	Q25%-Q75%		
6-MWD	13	456	± 108	480	420	- 537	66	384	± 110	397	334	- 451	*
NT-proBNP, pg/ml	13	205	± 340	92	60	- 193	69	216	± 249	128	68	- 256	
pulmonary function testing													
FVC, l	13	2.9	± 1.2	2.9	2.1	- 4.0	69	2.8	± 0.9	2.5	2.0	- 3.3	
FVC, %	13	88.5	± 20.5	84.0	73.0	- 103.7	71	90.2	± 23.5	90.0	71.0	- 110.0	
FEV ₁ , l	13	2.4	± 1.0	2.6	1.4	- 3.0	69	2.1	± 0.8	2.0	1.5	- 2.6	
FEV ₁ , %	13	89.0	± 21.5	85.3	70.6	- 109.5	70	83.0	± 22.6	83.7	64.6	- 99.8	
DL _{CO} , mmol/min/kPa	13	4.2	± 1.6	4.4	3.5	- 4.8	69	8.1	± 3.9	7.5	4.7	- 11.0	*
DL _{CO} , %	13	45.8	± 13.1	47.4	41.8	- 55.2	71	48.7	± 10.8	50.6	40.1	- 57.1	
DL _{CO} /VA, mmol/min/kPa/l	13	1.4	± 0.9	1.1	0.9	- 1.4	69	2.8	± 1.5	2.9	1.5	- 3.6	*
DL _{CO} /VA, %	13	70.1	± 18.8	67.5	61.0	- 82.9	69	71.4	± 17.3	70.7	59.0	- 83.0	
TLC, %	11	87.1	± 17.6	93.0	75.7	- 100.0	61	83.0	± 21.4	83.0	67.9	- 97.0	
RV, %	11	96.9	± 30.9	102.9	64.0	- 125.0	62	81.2	± 32.5	79.2	62.1	- 102.0	
echocardiography													
LA, mm	13	25.0	± 6.1	25.0	22.0	- 29.0	68	30.3	± 6.1	30.0	26.0	- 33.0	*
IVC, mm	12	12.3	± 4.5	12.3	9.5	- 14.0	58	14.1	± 3.4	14.0	12.0	- 16.0	
IVS, mm	13	10.5	± 1.5	10.6	9.0	- 11.0	67	10.0	± 1.9	10.0	9.0	- 11.0	
RA, cm ²	13	9.9	± 2.4	9.5	8.4	- 11.2	64	12.7	± 4.0	12.0	10.0	- 15.5	*
RVD, mm	11	30.8	± 4.2	31.0	29.0	- 33.0	62	28.5	± 6.0	29.5	25.0	- 33.3	
RV, cm ²	13	14.1	± 3.4	14.0	13.3	- 15.0	63	14.8	± 4.4	15.0	11.7	- 18.5	
LV-EDD, mm	13	44.3	± 6.0	44.0	39.0	- 49.0	67	44.2	± 5.4	44.0	41.2	- 47.0	
LV-ESD, mm	13	27.8	± 5.3	26.0	25.0	- 31.0	66	26.8	± 5.1	26.0	23.0	- 30.0	
TRV, m/s	12	2.5	± 0.4	2.5	2.4	- 2.6	66	2.5	± 0.4	2.5	2.3	- 2.7	
TAPSE, mm	13	22.8	± 4.5	23.0	20.0	- 26.0	64	22.5	± 5.0	22.0	19.0	- 26.0	
sPAP, mmHg	12	29.6	± 6.9	29.0	27.3	- 32.8	65	29.7	± 7.4	30.0	25.3	- 33.7	

right heart catheterisation

mPAP, mmHg	13	16.4 ± 4.1	16.0	14.0 - 19.0	71	17.7 ± 3.7	18.0	15.0 - 21.0	
PAWP, mmHg	13	6.9 ± 2.8	7.0	5.0 - 8.0	71	9.0 ± 3.0	9.0	7.0 - 11.0	*
TPG, mmHg	13	9.4 ± 3.5	8.0	7.0 - 12.0	71	8.6 ± 2.9	8.0	6.0 - 11.0	
CO, l/min	13	5.3 ± 1.3	5.3	4.4 - 6.2	71	5.3 ± 1.2	5.2	4.4 - 6.1	
PVR, dynes	13	148.4 ± 73.1	12.7	103.2 - 171.4	71	138.2 ± 52.9	126.0	104.9 - 163.3	
RAP, mmHg	13	3.5 ± 2.0	4.0	2.0 - 5.0	71	4.2 ± 2.3	4.0	3.0 - 5.0	

Q25% - Q75% = 25% and 75% quantiles, 6-MWD = 6-minute walking distance, NTproBNP = N-terminal pro Brain Natriuretic Peptide, FVC = forced vital capacity, FEV1 = forced expiratory volume in one second, DLCO = diffusion capacity of the lung for carbon monoxide, VA = alveolar volume, TLC = total lung capacity, RV = residual volume, LA = left atrium, IVC = inferior vena cava, IVS = interventricular septum, RA = right atrium, RVD = right ventricular diameter, RV = right ventricle, LV = left ventricle, EDD = end-diastolic diameter, ESD = end-systolic diameter, TRV = tricuspid regurgitation velocity, TAPSE = tricuspid annular plane systolic excursion, sPAP = systolic pulmonary arterial hypertension, mPAP = mean pulmonary arterial pressure, PAWP = pulmonary arterial wedge pressure, TPG = transpulmonary gradient, CO = cardiac output, PVR = pulmonary vascular resistance.

* denotes a significant difference between groups.

Supplementary table 6

Additional follow-up characteristics of PH Groups

	normal					PH														
						PAH					PH left heart disease					PH lung disease				
	n	mean	SD	median	Q25%-Q75%	n	mean	SD	median	Q25%-Q75%	n	mean	SD	median	Q25%-Q75%	n	mean	SD	median	Q25%-Q75%
6-MWD	47	399 ± 121		400	327 - 472	4	325 ± 133		292	225 - 425	7	376 ± 159		390	340 - 434	8	366 ± 122		330	261 - 461
NT-proBNP, pg/ml	57	276 ± 377		132	74 - 237	5	418 ± 538		237	107 - 328	6	466 ± 717		177	114 - 328	9	1142 ± 2963		123	107 - 262
pulmonary function testing																				
FVC, l	58	2.6 ± 0.9		2.6	1.9 - 3.34	3	3.0 ± 0.3		2.9	2.8 - 3.3	6	3.3 ± 0.6		3.3	2.9 - 3.7	8	2.4 ± 1.6		1.7	1.2 - 4.1
FVC, %	57	88.9 ± 22		92.1	68.2 - 108.2	4	105.4 ± 27.7		115.4	87.3 - 123.4	6	106.2 ± 16.7		111.8	99 - 118.7	9	73 ± 34		56.4	53.5 - 106.6
FEV ₁ , l	57	2.01 ± 0.8		2.0	1.4 - 2.5	3	2.0 ± 0.1		2.0	1.9 - 2.1	6	2.5 ± 0.5		2.6	2.2 - 3.0	8	2.0 ± 1.2		1.5	1.0 - 3.3
FEV ₁ , %	57	81.8 ± 20,5		80.1	64.9 - 95.9	3	95.2 ± 10.9		90.5	87.4 - 107.7	6	96.2 ± 12.3		95.2	84.1 - 109.4	9	74.2 ± 31.7		67.8	53.8 - 96.5
DL _{co} , mmol/min/kPa	55	2.9 ± 1.6		3.9	3.0 - 4.77	3	3.0 ± 0.4		3.2	2.6 - 3.3	5	4.4 ± 0.5		4.5	4.5 - 4.6	5	3.1 ± 1.3		3.5	2.2 - 3.6
DL _{co} , %	55	47.4 ± 11.2		50.4	41.3 - 55.7	4	38.8 ± 8.2		38.8	31.9 - 45.7	5	54.6 ± 8.9		54	48.1 - 62	6	34.8 ± 13.1		31	24.3 - 49.6
DL _{co} /VA, mmol/min/kPa/l	55	1.38 ± 0.9		1.05	0.9 - 1.4	3	0.7 ± 0.1		0.7	0.6 - 0.8	5	2.5 ± 1.9		1.5	1.1 - 3.6	6	1.7 ± 2.2		0.8	0.6 - 1.6
DL _{co} /VA, %	55	69.0 ± 15.6		68.1	60.3 - 82.8	3	48.8 ± 6.8		52.7	41 - 52.7	5	79.1 ± 18.5		75	71.2 - 93.1	6	56.2 ± 22.6		50.8	45.7 - 78
TLC, %	55	88.7 ± 21.9		89	68.1 - 106.2	3	101.3 ± 6.4		102.1	94.5 - 107.2	6	101.8 ± 19.1		104.1	85.1 - 111.9	6	81.1 ± 36		72.9	54 - 122.4
RV, %	49	95.16 ± 34.6		97.9	79 - 113	3	93.8 ± 7.9		89.9	88.6 - 102.9	5	100.5 ± 35.4		94.8	89.1 - 105.3	6	38.9 ± 42.5		65.2	54.1 - 107.9
echocardiography																				
LA, mm	55	30.4 ± 6.5		30	28 - 34	4	29.2 ± 1.9		29	28 - 30.4	7	31.6 ± 8.1		32	8.5 - 35	6	26.8 ± 7.3		26.5	22 - 32
IVC, mm	50	12.2 ± 5.9		14	8.6 - 16	2	18.5 ± 5.0		18.5	15 - 22	5	11.1 ± 8.0		10	8,5 - 13	7	13.9 ± 2.8		14	11 - 16
IVS, mm	30	10.7 ± 2.0		10.4	9 - 12	3	8.7 ± 1.6		9.2	7 - 10	2	10.1 ± 0.1		10.1	10 - 10.2	5	7.5 ± 3.8		8	8 - 9.3
RA, cm ²	54	12.6 ± 4.1		12	10.6 - 13.8	4	15.8 ± 6.5		13.8	11.5 - 20	6	12.5 ± 3.0		12.8	10 - 14	8	11.7 ± 3.3		12.4	9.2 - 14.5
RVD, mm	30	31 ± 7		30	27 - 36	4	36 ± 8		34	30 - 41	3	33 ± 3		33	30 - 36	5	28 ± 4		26	26 - 30
RV, cm ²	53	13.24 ± 3.63		13.5	10.3 - 15.5	4	15.9 ± 7.1		15.5	9.8 - 22	6	12.9 ± 2.0		13.0	11.3 - 15	9	12.3 ± 2.7		12.5	10.3 - 15
LV-EDD, mm	31	42.3 ± 5.7		43	39 - 46	4	47.2 ± 5.4		46	42.9 - 51.5	2	45.7 ± 0.4		45.7	45.4 - 46	5	35.5 ± 19.0		40	34 - 44
LV-ESD, mm	31	26.4 ± 5.5		27	21.8 - 31	4	29.6 ± 2.1		29.8	28.3 - 31	2	31.2 ± 1.2		31.2	30.3 - 32	5	23.5 ± 14.8		24	21 - 24
TRV, m/s	47	2.5 ± 0.4		2.5	2.3 - 2.8	3	2.9 ± 0.6		2.8	2.4 - 3.6	6	2.4 ± 0.4		2.4	2.1 - 2.6	7	2.9 ± 0.4		2.8	2.5 - 3.2

TAPSE, mm	56	22.1 ± 4.2	22	19 - 25	5	22 ± 3.5	24	22 - 24	7	22.3 ± 8.2	22	19 - 27	9	21.8 ± 6.9	20	18 - 26
sPAP, mmHg	47	31.4 ± 8.4	30	25.8 - 35.7	3	40.3 ± 14.9	35.9	28.0 - 56.8	6	28.8 ± 7.0	27.5	22.8 - 31.6	7	38.7 ± 9.7	36.8	29.6 - 47.0
right heart catheterisation																
mPAP, mmHg	52	19 ± 3.7	19	16 - 22	5	30.6 ± 3.2	31	28 - 32	5	27.8 ± 2	27	26 - 30	9	27.7 ± 6.6	27	26 - 32
PAWP, mmHg	52	10.5 ± 3	11	8.5 - 13	5	11 ± 3	12	11 - 12	5	17.6 ± 3.3	16	16 - 20	9	9.1 ± 2.3	9	8 - 11
TPG, mmHg	52	8.4 ± 2.6	8	6.5 - 11	5	10.4 ± 2.3	10	9 - 11	5	7.6 ± 1.7	8	7 - 9	9	10.2 ± 3.1	11	10 - 11
CO, l/min	52	5.2 ± 1.2	5	4.2 - 5.9	5	5.4 ± 0.7	5.3	5 - 5.4	5	5.4 ± 1.3	6	5.4 - 6.1	9	4.7 ± 0.8	4.6	4.3 - 4.7
PVR, dynes	52	144.3 ± 79.0	147.2	98.3 - 180.5	5	300.3 ± 81.5	281	250 - 345	5	167.6 ± 85.6	132.7	131 - 163	9	340.1 ± 171.1	330.4	238 - 483
RAP, mmHg	50	5 ± 3	5	3 - 6	5	6 ± 1	7	5 - 7	5	10 ± 3	9	8 - 10	9	5 ± 2	4	4 - 6

PAH = pulmonary arterial hypertension, PH left heart disease = pulmonary hypertension secondary to left heart disease,
 PH lung disease = pulmonary hypertension secondary to lung disease, Q25% - Q75% = 25% and 75% quantiles, 6-MWD = 6-minute walking distance,
 NTproBNP = N-terminal pro Brain Natriuretic Peptide, FVC = forced vital capacity, FEV1 = forced expiratory volume in one second,
 DLCO = diffusion capacity of the lung for carbon monoxide, VA = alveolar volume, TLC = total lung capacity, RV = residual volume, LA = left atrium, IVC = inferior vena cava,
 IVS = interventricular septum, RA = right atrium, RVD = right ventricular diameter, RV = right ventricle, LV = left ventricle, EDD = end-diastolic diameter,
 ESD = end-systolic diameter, TRV = tricuspid regurgitation velocity, TAPSE = tricuspid annular plane systolic excursion, sPAP = systolic pulmonary arterial hypertension,
 mPAP = mean pulmonary arterial pressure, PAWP = pulmonary arterial wedge pressure, TPG = transpulmonary gradient, CO = cardiac output,
 PVR = pulmonary vascular resistance.