



Early View

Research letter

Serum mitochondrial DNA predicts the risk of acute exacerbation and progression of IPF

Koji Sakamoto, Taiki Furukawa, Yasuhiko Yamano, Kensuke Kataoka, Ryo Teramachi, Anjali Walia, Atsushi Suzuki, Masahide Inoue, Yoshio Nakahara, Changwan Ryu, Naozumi Hashimoto, Yasuhiro Kondoh

Please cite this article as: Sakamoto K, Furukawa T, Yamano Y, *et al.* Serum mitochondrial DNA predicts the risk of acute exacerbation and progression of IPF. *Eur Respir J* 2020; in press (<https://doi.org/10.1183/13993003.01346-2020>).

This manuscript has recently been accepted for publication in the *European Respiratory Journal*. It is published here in its accepted form prior to copyediting and typesetting by our production team. After these production processes are complete and the authors have approved the resulting proofs, the article will move to the latest issue of the ERJ online.

**Serum mitochondrial DNA predicts the risk of acute exacerbation and progression of
IPF**

Koji Sakamoto^{1*}, Taiki Furukawa^{2*}, Yasuhiko Yamano³, Kensuke Kataoka³, Ryo Teramachi¹, Anjali Walia⁴, Atsushi Suzuki¹, Masahide Inoue¹, Yoshio Nakahara¹, Changwan Ryu⁴, Naozumi Hashimoto¹, Yasuhiro Kondoh³

¹Department of Respiratory Medicine, Nagoya University Graduate School of Medicine

²Department of Medical IT Center, Nagoya University Hospital, Nagoya, Japan

³Department of Respiratory and Allergy, Tosei General Hospital, Seto, Japan

⁴Section of Pulmonary, Critical Care, and Sleep Medicine, Yale University School of Medicine, New Haven, Connecticut

*These authors contributed equally to this work.

Corresponding author: Koji Sakamoto MD PhD.

Department of Respiratory Medicine

Nagoya University Graduate School of Medicine

Address: 65 Tsurumai-cho, Showa-ku, Nagoya 466-8550. Japan

email: sakakoji@med.nagoya-u.ac.jp

phone: +81527442167 Fax: +81527442176

Author contributions: Study design: KS, TF and YK. Data collection: KS, TK, YY, KK, and RT. Data analysis and interpretation: KS, TF, AS, and NH. Manuscript draft: KS, TF AW, CR and YK. Supervision and coordination of the study: NH and YK. Critical revision and final decision to submit: all authors.

Funding sources: This work was supported by the funds from Japan Society for the Promotion of Science (JSPS) KAKENHI [grant numbers 18K15948], and grants from the Satoshi Okamoto Memorial Foundation of Pulmonary Fibrosis.

Character count title: 86

Total word count of text file: 1161

Reference count: 10, Number of figures: 1.

Take home message

Patients with IPF often suffer from acute exacerbation, an unpredictable and deadly complication. Our study elucidated the potential of circulating mitochondrial DNA as a predictor of acute exacerbation and disease progression of IPF.

To the Editor

Idiopathic pulmonary fibrosis (IPF) is a progressive, fatal interstitial lung disease with a median survival of 3-5 years [1]. Its disease course is highly variable, as some

patients experience rapid deterioration in lung function while others experience more gradual decline [2]. The development of acute exacerbation of IPF (AE-IPF), a highly lethal complication of unknown etiology, has been shown to accelerate disease progression [3]. Presently, there are no accepted biomarkers that predict clinical deterioration [4], thus indicating an important, unmet need in the management of this devastating disease.

The pathogenic role of innate immunity has been an emerging field of study, with recent work highlighting the endogenous innate immune agonist mitochondrial DNA (mtDNA), which acts as a danger associated molecular pattern (DAMP) for pathogen recognition receptors (PRRs). Elevated concentrations of extracellular mtDNA have been observed in multiple diseases, including IPF, where it was found to be predictive of poor survival [5]. Given that the role of mtDNA as a biomarker for disease progression has not been fully characterised, we hypothesised that elevated concentrations of serum mtDNA reflect the mechanisms underlying poor clinical outcomes in IPF, including the development of AE-IPF.

This study was approved by the Institutional Review Board at Tosei General Hospital and performed with written informed consent from participants. IPF diagnosis was made upon multidisciplinary discussion based on consensus guidelines [1]. Our IPF cohort included 70 subjects with a median age of 66 (interquartile range (IQR): 63-71) that included 56 males (80.0%) and 46 ever-smokers (65.7%). Relevant clinical characteristics included 84 percent predicted forced vital capacity (%FVC, IQR 69-95%), 58 percent predicted diffusion capacity for carbon monoxide (%DLCO, IQR 42-73%), and 3 Gender-Age-Physiology (GAP) score [6] (IQR 2-4). Subjects were followed for a median of 52

months (Kaplan-Meier estimate), during which time 17 cases of AE-IPF (24.2%) and 30 deaths (42.9%) were observed. The diagnosis of AE-IPF was confirmed retrospectively according to published criteria [3]. Blood samples drawn at each subject's initial clinical encounter, prior to the initiation of anti-fibrotic therapy, were used in the analysis. Serum mtDNA concentrations were determined as previously described [7]; briefly, total cell-free DNA was extracted from 240 μ l of serum, and mtDNA concentrations were expressed as copy numbers of the NADH dehydrogenase 1 gene per microlitre via quantitative PCR.

We first quantified serum mtDNA copy numbers in our IPF cohort, where the median concentration was 801 copies/ μ l. When subjects were stratified by this threshold value, subjects with serum mtDNA concentrations \geq 801 copies/ μ l exhibited higher incidence of AE-IPF (3.1 vs 14.5 per person-months, $p=0.005$ by Fisher's exact test), greater annual decline in %FVC (-1.1 vs -7.5, $p=0.033$ by Mann-Whitney U test), and mortality (median survival; 36.2 vs 76.6 months, $p=0.0001$ by Log-rank test). Moreover, no significant differences were observed in age, gender, smoking status, or pulmonary function between the low (<801 copies/ μ l) and high (\geq 801 copies/ μ l) mtDNA groups. These findings show that high serum mtDNA copy numbers are associated with poor outcomes in IPF.

To further characterise the role of serum mtDNA concentrations as a biomarker for the development of AE-IPF, we conducted time-dependent receiver operating characteristic (ROC) curve analysis (Figure 1A), which revealed that a serum mtDNA concentration of 1371.5 copies/ μ l can reliably stratify subjects at risk for developing AE-IPF within one year (sensitivity of 85.4%, specificity of 80.6%, area under the curve (AUC):

0.90). Following adjustments for mortality as a competing risk [8], we observed a substantially higher risk for developing AE-IPF (median incidence time 5.8 vs 24.3 months, $p=0.010$ by Gray's test, Figure 1B). These data show that elevated circulating mtDNA levels can identify IPF subjects at high risk for developing AE-IPF.

Because current biomarkers poorly predict IPF disease progression [4], we then sought to determine if mtDNA copy numbers could provide insight into overall disease trajectory. Using serum mtDNA concentration as a continuous variable, we identified a significant association with composite IPF outcomes, which we defined as >10% absolute decline in %FVC, acute exacerbation, or death within one year (AUC: 0.708, 95% CI: 0.577 - 0.939, Figure 1C). More excitingly, serum mtDNA levels predicted this composite outcome better than other well-known IPF biomarkers, including KL-6 (AUC: 0.452, Figure 1C) and SP-D (AUC: 0.535, Figure 1C). Taken together, these findings suggest that serum mtDNA assessment in patients with stable IPF may predict future disease behavior.

Because we also found a significant association between serum mtDNA concentration and mortality, we used the above serum mtDNA concentration cutoff value of 1371.5 copies/ μ l to confirm prior reports [5] showing that an elevated concentration of circulating mtDNA is associated with increased mortality (median survival: 76.6 months vs 36.2 months, $p=0.002$ by Log-rank test, Figure 1D). Additionally, we performed Cox Regression analysis with this serum mtDNA cut point and GAP Stage, a validated clinical biomarker of IPF survival [6], to confirm that mtDNA levels (Hazard Ratio (HR) 2.970, 95% confidence interval (95% CI) 1.418 – 6.218, $p=0.004$) and GAP stage (HR 1.821 (95%CI 0.822-4.033), $p=0.140$ for stage 2; HR 8.052 (95%CI 2.219-29.23), $p=0.002$ for stage 3,

with stage 1 as a reference) are independently associated with mortality. These data further validate the role of circulating mtDNA as a prognostic biomarker in IPF.

Lastly, we performed exploratory longitudinal analysis with serum samples obtained at the time of AE-IPF diagnosis for 7 subjects that died from this complication within 3 months following diagnosis. We found a substantial increase in their serum mtDNA concentrations relative to their respective baseline concentrations ($p=0.031$ by Wilcoxon rank test, Figure 1E). Moreover, serum concentrations of KL-6 and SP-D exhibited only modest changes in these subjects (Figure 1E). While this analysis is preliminary, these findings suggest that serial changes in mtDNA levels could function as an indicator of prognosis in AE-IPF.

Overall, our results robustly support the role of circulating mtDNA as a biomarker of poor clinical outcomes in IPF. Specifically, we found that elevated serum mtDNA levels were associated with the development of acute exacerbation, disease progression, and worse survival. While additional work is needed to fully characterise the immunopathogenic role of extracellular mtDNA in IPF, studies show that differential expression of Toll-like receptor 9 [9] and cGAS-STING [10], PRRs that recognise mtDNA, is associated with IPF outcomes. This constellation of findings suggests that interactions between mtDNA and these PRRs determine individual susceptibility to the development of AE and rapid IPF disease progression.

Our work has several limitations. We have not determined the cell of origin of this extracellular mtDNA, which would require access to matched bronchoalveolar lavage and

serum samples. In addition, our work is limited to a single cohort at one institution; validation studies, including determination of optimal threshold values, through multicenter efforts are needed. Nonetheless, our work shows that circulating mtDNA is a promising biomarker of poor prognosis in IPF. Further work in this arena can lead to new approaches in the management of this poorly understood, fatal disease.

Acknowledgements

The authors would like to thank Kana Shimazaki and Yoshie Sato (Department of Respiratory Medicine, Nagoya University) for technical assistance.

References

1. Raghu G, Remy-Jardin M, Myers JL, et al. Diagnosis of Idiopathic Pulmonary Fibrosis. An Official ATS/ERS/JRS/ALAT Clinical Practice Guideline. *Am J Respir Crit Care Med*. 2018;198(5): e44-e68. doi:10.1164/rccm.201807-1255ST
2. Ley B, Collard HR, Talmadge E, King J. Clinical Course and Prediction of Survival in Idiopathic Pulmonary Fibrosis. *Am J Respir Crit Care Med* 2011; 183: 431-440.
3. Collard HR, Ryerson CJ, Corte TJ, Jenkins G, Kondoh Y, Lederer DJ, Lee JS, Maher TM, Wells AU, Antoniou KM, Behr J, Brown KK, Cottin V, Flaherty KR, Fukuoka J, Hansell DM, Johkoh T, Kaminski N, Kim DS, Kolb M, Lynch DA, Myers JL, Raghu G, Richeldi L, Taniguchi H, Martinez FJ. Acute Exacerbation of Idiopathic Pulmonary Fibrosis. An International Working Group Report. *Am J Respir Crit Care Med* 2016; 194: 265-275.
4. Ley B, Bradford WZ, Vittinghoff E, Weycker D, Bois RMd, Collard HR. Predictors of Mortality Poorly Predict Common Measures of Disease Progression in Idiopathic Pulmonary Fibrosis. *Am J Respir Crit Care Med* 2016; 194: 711-718.
5. Ryu C, Sun H, Gulati M, Herazo-Maya JD, Chen Y, Osafo-Addo A, Brandsdorfer C, Winkler J, Blaul C, Faunce J, Pan H, Woolard T, Tzouvelekis A, Antin-Ozerkis DE, Puchalski JT, Slade M, Gonzalez AL, Bogenhagen DF, Kirillov V, Feghali-Bostwick C, Gibson K, Lindell K, Herzog RI, Cruz CSD, Mehal W, Kaminski N, Herzog EL, Trujillo G. Extracellular Mitochondrial DNA Is Generated by Fibroblasts and Predicts Death in Idiopathic Pulmonary Fibrosis. *Am J Respir Crit Care Med* 2017; 196: 1571-

1581.

6. Ley B, Ryerson CJ, Vittinghoff E, Ryu JH, Tomassetti S, Lee JS, Poletti V, Buccioli M, Elicker BM, Jones KD, King TE, Jr., Collard HR. A multidimensional index and staging system for idiopathic pulmonary fibrosis. *Ann Intern Med* 2012; 156: 684-691.
7. Nakahira K, Kyung S-Y, Rogers AJ, Gazourian L, Youn S, Massaro AF, Quintana C, Osorio JC, Wang Z, Zhao Y, Lawler LA, Christie JD, Meyer NJ, Causland FRM, Waikar SS, Waxman AB, Chung RT, Bueno R, Rosas IO, Fredenburgh LE, Baron RM, Christiani DC, Hunninghake GM, Choi AMK. Circulating Mitochondrial DNA in Patients in the ICU as a Marker of Mortality: Derivation and Validation. *PLOS Med* 2014; 10: e1001577.
8. Scrucca L, Santucci A, Aversa F. Competing risk analysis using R: an easy guide for clinicians. *Bone Marrow Transplant* 2007; 40: 381-387.
9. Trujillo G, Meneghin A, Flaherty KR, Sholl LM, Myers JL, Kazerooni EA, Gross BH, Oak SR, Coelho AL, Evanoff H, Day E, Toews GB, Joshi AD, Schaller MA, Waters B, Jarai G, Westwick J, Kunkel SL, Martinez FJ, Hogaboam CM. TLR9 Differentiates Rapidly from Slowly Progressing Forms of Idiopathic Pulmonary Fibrosis. *Sci. Transl. Med.* 2010; 2: 57ra82-57ra82.
10. Qiu H, Weng D, Chen T, Shen L, Chen S-S, Wei Y-R, Wu Q, Zhao M-M, Li Q-H, Hu Y, Zhang Y, Zhou Y, Su Y-L, Zhang F, Lu L-Q, Zhou N-Y, Li S-L, Zhang L-L, Wang C, Li H-P. Stimulator of Interferon Genes Deficiency in Acute Exacerbation of Idiopathic Pulmonary Fibrosis. *Front Immunol* 2017; 8.

Figure Legend

Figure 1: Assessment of mitochondrial DNA levels in sera as a predictor of multiple

clinical outcomes in IPF patients. (A) Time-dependent area under the curve (AUC)

changes over time according to various cutoff mtDNA levels assessing predictive value for

acute exacerbation of IPF. (B) Cumulative incidence curves of acute exacerbation (red)

and competing all-cause mortality (black) stratified by serum mtDNA level with the cut-off of

1371.5 copies/ul. In the comparison of the cumulative incidence rate of acute exacerbation

between the low mtDNA group and high group, the p-value for Gray's test, as well as the

hazard ratio (HR) and its 95% confidence interval (95% CI) for Cox regression analysis are

shown. (C) Receiver operating characteristic (ROC) curves for evaluating serum

biomarkers as predicting composite outcome (>10% absolute decline of %FVC, acute

exacerbation, or death) within a year. Area under the curves (AUC) and their 95%

confidence intervals (95% CI) for each biomarker were shown. (D) Kaplan Meier curve

estimating the probability of overall survival in the cohort stratified by circulating mtDNA

levels (Cutoff 1371.5 copies/ μ L). P-value, HR and its 95% CI for log-rank test in the

comparison of 2 groups are shown. (E) Changes in levels of serum markers obtained from

7 patients who developed AE in their clinical course and subsequently died within 3

months. Changes of serum mtDNA, serum KL-6, and serum SP-D in each subject were

depicted. * $p = 0.031$ by Wilcoxon rank test.

