Early View

Research letter

Practices of therapeutic drug monitoring in tuberculosis - an international survey

Ioana Margineanu, Onno Akkerman, Dario Cattaneo, Delia Goletti, Deborah J E Marriott, Giovanni Battista Migliori, Fuad Mirzayev, Charles A Peloquin, Ymkje Stienstra, Jan-Willem Alffenaar

Please cite this article as: Margineanu I, Akkerman O, Cattaneo D, *et al.* Practices of therapeutic drug monitoring in tuberculosis - an international survey. *Eur Respir J* 2022; in press (<https://doi.org/10.1183/13993003.02787-2021>).

This manuscript has recently been accepted for publication in the *European Respiratory Journal*. It is published here in its accepted form prior to copyediting and typesetting by our production team. After these production processes are complete and the authors have approved the resulting proofs, the article will move to the latest issue of the ERJ online.

Practices of therapeutic drug monitoring in tuberculosis - an international survey

Ioana Margineanu PhDc, MD 1, Onno Akkerman PhD MD 2,3, Dario Cattaneo PhD PharmD 4*, Delia Goletti Prof, MD 5*, Deborah J E Marriott Prof MD 6*, Giovanni Battista Migliori Prof MD 7*, Fuad Mirzayev MD 8*, Charles A Peloquin Prof PharmD 9*, Ymkje Stienstra Prof MD 10, Jan-Willem Alffenaar Prof PharmD 11

*in alphabetical order

1 - Department of Clinical Pharmacy and Pharmacology, Rijksuniversiteit Groningen, University Medical Centrum Groningen, the Netherlands

2 - University Of Groningen, University Medical Centrum Groningen, Department of Pulmonary Diseases and Tuberculosis, Groningen, the Netherlands

3 - University Of Groningen, University Medical Centrum Groningen, TB center Beatrixoord, Groningen, the Netherlands

4 - Unit of Clinical Pharmacology, ASST Fatebenefratelli Sacco University Hospital, Milan, Italy

5 - Translational Research Unit, National Institute for Infectious Diseases (INMI), Rome Italy

6 - Department of Clinical Microbiology and Infectious Diseases, St Vincent's Hospital, Sydney, New South Wales, Australia

7 - Istituti Clinici Scientifici Maugeri IRCCS, Tradate, Italy

8 - Global TB Programme, World Health Organization, Geneva, Switzerland

9 - University of Florida, College of Pharmacy and Emerging Pathogens Institute, United States

10 - Department of Internal Medicine, Rijksuniversiteit Groningen, University Medical Centrum Groningen, the Netherlands

11 - School of Pharmacy, Faculty of Medicine and Health, University of Sydney, Australia
Correspondence

i.s.margineanu@umcg.nl and ismargineanu@gmail.com

Tuberculosis (TB), is still one of the top 10 causes of death in low and lower-middle income countries [1]. TB's long and complex treatment, side-effects, and development of resistant bacteria compromise treatment success. To improve treatment outcomes, therapeutic drug monitoring (TDM) has been included in TB treatment guidelines [2–4] to be considered for specific situations in which there is documented or expected poor response to treatment, drug toxicity, or a lower drug concentration. Several strategies for implementation of TDM for programmatic use have been proposed to overcome barriers to widespread use of TDM [5,6], including more accessible techniques such as dried-blood spot analysis or saliva and urine testing, [7] but uptake in programmatic care is still limited [8].

To understand current use and barriers and facilitators to implementing TDM in TB management we performed an electronic survey which can be found at this webpage:

<https://is.gd/TDMSurvey>

The survey consisted of six sections asking participants if and how TDM is used in their setting. Several questions had multiple-choice answers. The survey was piloted by three of the co-authors (GB, DG, CP) and it took on average 15 minutes to complete. The survey was built in RedCap Systems and distributed through the Global Tuberculosis Network (160 email addresses), the International Association for Therapeutic Drug Monitoring and Clinical Toxicology (638 email addresses), and the Global Drug-resistant TB Initiative members (400 email addresses).

Responses were received from 86 participants, spanning 46 countries across all seven continents (figure 1).

Participants worked as clinicians (19, 22%), pharmacists (31, 36%), researchers (34, 40%), and public health staff (8, 9%). Half worked as academic professionals (43, 50%). By type of medical facility, most participants worked in an university hospital (36, 42%), followed by tuberculosis expert center (22, 26%), general hospital (18, 21%), private hospital (9, 10%), infectious diseases center (8, 9%), outpatient clinic (6, 7%), with several working in more than one facility (12, 13.9%). A majority answered on behalf of their institution only (60, 70%), with 11 (13%) and 15 (17%) responding for region and country respectively.

TDM was performed by 43 out of the total of 86 survey participants (50%), within a total of 18 countries. Those who undertook TDM, indicated the following reasons for initiation: gastrointestinal abnormalities were the most common indication (24/43, 56%), followed by chronic kidney disease, cirrhosis, and HIV+ (22/43, 51%), acute kidney injury (20/43, 47%), diabetes mellitus (19/43, 44%), pregnancy (18/43, 42%), malnutrition (17/43, 40%), and age < 18 (9/43, 21%). Two participants mentioned that all patients were eligible for TDM and one included "COVID-19" as a comorbidity.

Approximately half of all participants reported performing TDM for all TB patients irrespective of drug susceptibility (18/43, 42%), followed by performing TDM if resistant to rifampicin +/- other first line agents (7/43, 16%), MDR-TB (3/43, 7%), XDR-TB (1/43, 2%), or applying additional criteria such as slow-response or rapid relapse (3/43, 7%) or other, undisclosed criteria (11/43, 26%).

Considering drugs for which TDM is performed, 12/43 (28%) performed TDM on all first line drugs and 27/43 (63%) performed on at least rifampicin. The second most frequent drug class analysed was aminoglycosides (17, 40%), followed by linezolid (16/43, 37%), fluoroquinolones (12/43, 27%), carbapenems (11/43, 26%), bedaquiline and cycloserine (9/43, 21%).

TDM was most often performed in pharmacology laboratories (21/43, 49%), followed by university laboratories (7/43, 16%), microbiology (4/43, 9%), public health institutions (4/43, 9%), private laboratories (4/43, 9%), chemistry laboratories (3/43, 7%). Sampling strategies varied, with 13 (30%) performing peak and trough concentrations, sampling at different time points for each drug (12/43, 28%), and collecting a full pharmacokinetic curve (6/43, 14%). The samples most often utilised for TDM were plasma and serum (20/43, 47%), with a few participants reporting they performed TDM on dried blood spot samples (3/43, 7%), saliva or cerebro-spinal fluid (2/43, 5%). More than half of responses (24, 56%) indicated that drug assays were validated according to FDA, CLSI or EMA guidelines.

Participants indicated that TDM results were received in under five days in 15(35%), in five-ten days (9/43, 21%), or more than 10 days (3/43, 7%) and that they would use TDM results as advice for dose adjustment (25/43, 58%), changing the drug (12/43, 28%), or change in route of administration (11/43, 26%).

Participants utilised a variety of guidelines to perform TDM: national guidelines (10/43, 23%), specific articles (9/43, 21%), international guidelines (8/43, 19%), facility guidelines (5/43, 12%).

TDM was most often paid for by public healthcare or health insurance (21/43, 49%).

Alternative sources of funding were TB-specific research grants (6/43, 14%), hospital funds (6/43, 14%) or the patients themselves (5/43, 12%).

The following responses arose from questions open to all 86 participants, regardless of whether or not they performed TDM. Concerning costs, approximately half the participants considered TDM to be cost-effective (39/86, 45%), with some (8/86, 9%) stating it depended on several factors, such as long term dosing, length of hospital stay, and choosing the right patients for TDM. A minority considered TDM not cost-effective (8/86, 9%) and the rest responded that they do not know the answer to this question (31/86, 36%). Furthermore, participants would predominantly spend a maximum of 20 euros (14/86, 16%) or 10 euros (14/86, 16%) per assay, with some reporting they would go up to 50 euros (5/86, 6%) or even 100 euros (5/86, 6%).

Participants reported the main barriers to TDM usage to be a lack of knowledge among the medical staff (32/86, 37%), followed by lack of funding and lack of guideline usage (30/86, 35%), no political/governmental will (17/86, 20%), TDM implementation costs were too high (11/86, 13%), and TDM implementation too complicated (6/86, 7%). One participant specifically mentioned that “many clinicians don’t necessarily agree with TDM”, and another explained that the workload is very high as it is, without introducing a new technique.

Participants indicated training efforts should focus on clinical/laboratory staff (29/86, 34%), then educate stakeholders (26/86, 30%), and finally, hospital administrators (18/86, 22%).

The present study is the first to investigate TDM use in TB through a worldwide survey.

Our survey showed that progress in TDM is being made, with half of the survey participants using TDM. Cheaper and easy to use assays [9–11], clinical guidelines [2–4] and laboratory quality control standards [12] are all increasing in availability. However, the main challenge is a lack of widespread information about TDM, with progress being recognised by participants who look towards guidelines and research when performing TDM.

Half of our responders work in an academic field, indicating that responses arrived from experts already using or interested in using the technique. One potential limiting factor is that even through choosing dissemination networks relevant to the field, we might not have captured a representative sample of the target population of potential TDM users.

Responses to our survey indicate that there is concern surrounding cost-effectiveness and the resources available in different settings to implement TDM. Robust research in diverse settings is needed through high quality studies which investigate and can better inform stakeholders of the potential long-term cost-savings TDM can provide [13].

The electronic survey is open for further data collection and, if you would like to contribute towards further research in this field, you are invited to complete it: <https://is.gd/TDMSurvey>

Acknowledgements

This project is part of the scientific activities of the Global Tuberculosis Network and of the WHO Collaborating Centre for TB and Lung Diseases, Tradate, Italy

Funding

IM is funded from a doctoral project funded from the European Union Horizon 2020 research and innovation programme, under the Marie-Sklódowska Curie grant agreement 713660.

The funding source had no impact on any decision-making regarding this paper. The authors report no conflict of interest.

1. The top 10 causes of death. [cited 15 May 2021]. Available: <https://www.who.int/news-room/fact-sheets/detail/the-top-10-causes-of-death>
2. Nahid P, Dorman SE, Alipanah N, Barry PM, Brozek JL, Cattamanchi A, et al. Official American Thoracic Society/Centers for Disease Control and Prevention/Infectious Diseases Society of America Clinical Practice Guidelines: Treatment of Drug-Susceptible Tuberculosis. *Clin Infect Dis*. 2016;63: e147–e195. doi:10.1093/cid/ciw376
3. Nahid P, Mase SR, Migliori GB, Sotgiu G, Bothamley GH, Brozek JL, et al. Treatment of Drug-Resistant Tuberculosis. An Official ATS/CDC/ERS/IDSA Clinical Practice Guideline. *Am J Respir Crit Care Med*. 2019;200: e93–e142. doi:10.1164/rccm.201909-1874ST
4. Organization WH, Others. WHO operational handbook on tuberculosis: module 4: treatment: drug-resistant tuberculosis treatment. 2020. Available: <https://apps.who.int/iris/bitstream/handle/10665/332398/9789240006997-eng.pdf>
5. Ghimire S, Bolhuis MS, Sturkenboom MGG, Akkerman OW, de Lange WCM, van der Werf TS, et al. Incorporating therapeutic drug monitoring into the World Health Organization hierarchy of tuberculosis diagnostics. *Eur Respir J*. 2016;47: 1867–1869. doi:10.1183/13993003.00040-2016
6. Burgt EPM van der, van der Burgt EPM, Sturkenboom MGG, Bolhuis MS, Akkerman OW, Kosterink JGW, et al. End TB with precision treatment! *European Respiratory Journal*. 2016. pp. 680–682. doi:10.1183/13993003.01285-2015
7. Alffenaar J-WC, Heysell SK, Mpagama SG. Therapeutic Drug Monitoring: The Need for Practical Guidance. *Clinical infectious diseases: an official publication of the Infectious Diseases Society of America*. 2019. pp. 1065–1066. doi:10.1093/cid/ciy787
8. Kim HY, Heysell SK, Mpagama S, Marais BJ, Alffenaar J-W. Challenging the management of drug-resistant tuberculosis. *The Lancet*. 2020. p. 783. doi:10.1016/S0140-6736(20)30049-0
9. Kim HY, Ruiter E, Jongedijk EM, Ak HK, Marais BJ, Pk B, et al. Saliva-based linezolid monitoring on a mobile UV spectrophotometer. *J Antimicrob Chemother*. 2021;76: 1786–1792. doi:10.1093/jac/dkab075
10. Szipszky C, Van Aartsen D, Criddle S, Rao P, Zentner I, Justine M, et al. Determination of Rifampin Concentrations by Urine Colorimetry and Mobile Phone Readout for Personalized Dosing in Tuberculosis Treatment. *J Pediatric Infect Dis Soc*. 2021;10:

104–111. doi:10.1093/jpids/piaa024

11. Mave V, Kadam D, Gaikwad S, Kinikar A, Aguilar D, Chavan A, et al. Measuring TB drug levels in the hair in adults and children to monitor drug exposure and outcomes. *Int J Tuberc Lung Dis*. 2021;25: 52–60. doi:10.5588/ijtld.20.0574
12. Aarnoutse RE, Sturkenboom MGG, Robijns K, Harteveld AR, Greijdanus B, Uges DRA, et al. An interlaboratory quality control programme for the measurement of tuberculosis drugs. *Eur Respir J*. 2015;46: 268–271. doi:10.1183/09031936.00177014
13. Kim HY, Ulbricht E, Ahn YK, Gillooly IS, Lee KJ, Lieu J, et al. Therapeutic drug monitoring practice in patients with active tuberculosis: assessment of opportunities. *Eur Respir J*. 2021;57. doi:10.1183/13993003.02349-2020

Numbers represent the number of replies for each category from a specific country. For participants with multiple roles, the first one chosen is represented on this map.