Early View

Research letter

Lung function during and after acute respiratory infection in COVID-19 positive and negative outpatients

Pekka Tamminen, Dominik Kerimov, Hanna Viskari, Janne Aittoniemi, Jaana Syrjänen, Lauri Lehtimäki

Please cite this article as: Tamminen P, Kerimov D, Viskari H, *et al.* Lung function during and after acute respiratory infection in COVID-19 positive and negative outpatients. *Eur Respir J* 2022; in press (<https://doi.org/10.1183/13993003.02837-2021>).

This manuscript has recently been accepted for publication in the *European Respiratory Journal*. It is published here in its accepted form prior to copyediting and typesetting by our production team. After these production processes are complete and the authors have approved the resulting proofs, the article will move to the latest issue of the ERJ online.

Copyright ©The authors 2022. This version is distributed under the terms of the Creative Commons Attribution Non-Commercial Licence 4.0. For commercial reproduction rights and permissions contact permissions@ersnet.org

Title: **Lung function during and after acute respiratory infection in COVID-19 positive and negative outpatients**

Authors: Pekka Tamminen^{a,b}, Dominik Kerimov^c, Hanna Viskari^{a,e}, Janne Aittoniemi^c, Jaana Syrjänen^{a,e} and Lauri Lehtimäki^{d,e}

Affiliations:

a. Department of Internal Medicine, Tampere University Hospital, Elämänaukio 2, 33520 Tampere, Finland

b. Department of Otorhinolaryngology – Head and Neck Surgery, Tampere University Hospital, Elämänaukio 2, 33520 Tampere, Finland

c. Department of Clinical Microbiology, Fimlab Laboratories, Arvo Ylpön Katu 4, 33520, Tampere, Finland

d. Allergy Centre, Tampere University Hospital, Elämänaukio 2, 33520 Tampere, Finland

e. Faculty of Medicine and Health Technology, Tampere University, Arvo Ylpön Katu 34, 33520 Tampere, Finland

Corresponding author: Pekka Tamminen

Corresponding author email: pekka.tamminen@tuni.fi

Conflict of interest: None declared.

Authors' contributions:

Pekka Tamminen: Conceptualization, Investigation, Methodology, Formal analysis, Data Curation, Writing - Original Draft, Writing - Review & Editing, Visualization.

Dominik Kerimov: Conceptualization, Investigation, Writing - Original Draft , Writing - Review & Editing.

Hanna Viskari: Conceptualization, Investigation, Writing - Review & Editing, Supervision.

Janne Aittoniemi: Conceptualization, Writing - Review & Editing, Supervision.

Jaana Syrjänen: Writing - Review & Editing, Supervision.

Lauri Lehtimäki: Conceptualization, Methodology, Validation, Formal analysis, Writing - Review & Editing, Supervision, Project administration, Funding acquisition.

Funding: This work was supported by funding from the Tampere Tuberculosis Foundation; the Research Foundation of the Pulmonary Diseases; the Competitive State Research Financing of the Expert Responsibility area of Tampere University Hospital, and the Tampere University Hospital Support Foundation. These funders did not have any role in any part of the study.

Keywords

SARS-CoV-2; COVID-19; airwave oscillometry (AOS); forced oscillation technique (FOT); pulmonary function; spirometry; post-acute sequelae

Take Home

At acute phase, outpatients with mild COVID-19 had more symptoms, higher small airway resistance and poorer lung elasticity compared to outpatients with other respiratory infections, but there was no difference between the groups after two months.

To the Editor:

Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) has been associated with prolonged post-acute symptoms in at least 10 % of patients [1, 2]. The majority of published data evaluates hospitalized patients with severe SARS-CoV-2 disease (COVID-19) with symptoms and pulmonary function defects several months after discharge [1]. Most of the infected subjects develop mild symptoms and are treated as outpatients. Though they are also reported to suffer from prolonged symptoms, their lung function is studied far less. Furthermore, the prolonged symptoms and objectively measurable findings are usually not compared to a group suffering from airway infection caused by other pathogens [2]. As spirometry and other aerosol-producing procedures are minimized during the pandemic, there are no reports on lung function during acute COVID-19.

Forced oscillation technique (FOT) or airwave oscillometry (AOS) is a non-invasive and effort-independent method to measure airway resistance and airway reactance, enabling safe and admissible evaluation of pulmonary function [3] without producing possibly infective aerosols. FOT has been used in three previous studies to show small airway impairment one to six months after discharge in subjects who have recovered from severe COVID-19 [4–6].

Our aim was to evaluate and compare symptoms and pulmonary function at the time of acute infection (visit one) and after two months of follow-up (visit two) in outpatients with mild respiratory infection that were either RT-PCR-positive (cases) or RT-PCR-negative (controls) for SARS-CoV-2.

The final study population consisted of two clinical series, namely 43 cases and 39 controls, who took part in the CANNAS-study in Tampere, Finland from January 15th to August 18th, 2021 (R20090, NCT04728919). Cases were recruited during contact tracing phone call. Age-matched ($\pm$ 5 years) and gender-matched controls were recruited among subjects tested for SARS-CoV-2 due to acute respiratory infection but who later received a negative test result and gave their contact information (n=138) at local testing sites; 102 people were contacted, 43 matched controls were recruited and after drop-outs 39 subjects were included. The exclusion criteria were absence of

respiratory symptoms, onset of symptoms over ten days ago, pregnancy or breastfeeding, previous vaccination against COVID-19 and inability to arrive at the study site without exposing others.

A standard nasopharyngeal swab sample was obtained and three target genes were analysed by the SARS-CoV-2 RT-PCR protocol (Seegene Inc., Seoul, South Korea). Lung function was measured with AOS (TremoFlo C-100, Thorasys, Montreal, Canada) at both visits and with spirometry (Medikro Pro, Medikro Oy, Kuopio, Finland) at the follow-up visit, according to international standards [7–9], using international and Finnish reference values [10, 11]. The visual analogue scale (VAS) from “none” (0 mm) to “worst possible” (100 mm) was used to score different symptoms at both visits. Due to skewed distributions (Kolmogorov-Smirnov test), medians (interquartile range) were used to describe the data.

The results are presented in the Table 1. Cases and controls did not differ in their demographics. None of the subjects used leukotriene antagonists. Systemic corticosteroids were used only by one subject for a nonidentifiable reason before the control visit (normal AOS and spirometry). Inhaled corticosteroids were used by 5% of cases (only one of them had the diagnosis of asthma) and 3% of controls ($p=1$). The median time from visit one to follow-up was 59 (56–64) vs 56 (53–67) days, $p=0.496$, in cases and controls, respectively.

Small airways resistance (R5-20) and area of reactance (AX) were significantly higher in cases during infection (visit one) in comparison to controls, indicating impairment of the peripheral lung. In both cases and controls, there was a small but statistically significant impairment in these parameters from visit one to two (Table 1). Proportions of subjects with abnormal results in each oscillometry variable (z -score < -1.65) were small, and there were no differences between the cases and controls at either visit. Spirometry results at follow-up were similar between the groups. Some of the subjects in both groups had spirometry values below the lower limit of normal (z -score < 1.65) (7.7% vs 14.3% in FVC, $p=0.477$; 12.8% vs 17.1% in FEV₁, $p=0.654$; 20.5% vs 14.2% in FEV₁/FVC, $p=0.554$, in cases and controls respectively) but there was no difference between the groups.

During acute infection, cases had higher symptom scores in terms of decreased senses of smell and/or taste, fatigue, coughing, the need to blow their noses and dyspnoea (Table 1). There was a significant decrease in symptoms in both groups at the follow-up and no significant difference between the groups.

This is the first study to evaluate outpatient lung function at the acute phase of COVID-19 infection and we had a control group suffering from respiratory infection caused by other pathogens. At the acute phase, cases had on average poorer peripheral lung function (R5-R20 and AX) in comparison to controls, yet the proportion of abnormal results was low. Although subjects recovering from severe COVID-19 infection may have impaired diffusing capacity, restriction or obstruction [12], our results suggest that mild COVID-19 infection does not impair lung function after two months of follow-up when compared to other mild respiratory infections. There were some subjects with lung function below the lower limit of normal, but this is understandable since, by definition, 5% of healthy subjects have abnormal values of each parameter, and our study also included subjects with asthma, a history of smoking (possible undiagnosed chronic obstruction) and obese people with pulmonary restriction. However, the proportion of cases with FEV_1/FVC below lower limit of normal was unexpectedly high (20.5%), but as we do not know their lung function before infection, we do not know if this is because of COVID-19 or undiagnosed prior lung disease. The important finding is that the proportion of abnormal values did not differ between the cases and controls and none of the subjects had $FEV_1/FVC < 0.70$.

Interestingly, the trend of change in our data was slightly worse in R5, R5-20, AX and X5 in both groups. This might implicate airway damage after infection [4]. Again, there were no differences between cases and controls, suggesting that mild COVID-19 does not differ from other airway infections in this regard.

COVID-19 patients were more symptomatic than the controls at the acute phase, with the greatest differences being a decreased sense of taste and/or smell, and fatigue, which align with previous results [13, 14]. Both groups had very low levels of symptoms at control visit and there was no

difference between the groups suggesting that long-lasting sequelae are not frequent in subjects with mild COVID treated as outpatients.

The study sample is representative of non-vaccinated adult Finnish outpatients with mild RTI, albeit the participation might have been more tempting to people with pulmonary or other health concerns. Measurements at visit one were taken days after the onset of symptoms, not necessarily depicting the culmination of the disease. However, the groups were comparable by background characteristics. The sample size was based on a 90% power with an alpha-error of 5% to find a between-group difference in any continuous variable that was at least 75% of the standard deviation. Thus, we cannot rule out minor differences between the groups.

In conclusion, outpatients with mild COVID-19 had poorer peripheral lung function in comparison to those with other acute respiratory infections, but there were no differences in lung function or symptoms after two months of follow-up, suggesting that possible long-standing symptoms after mild COVID-19 may not be due to physical changes in the airways.

Acknowledgements

The authors would like to thank study nurses Merja Rumpunen and Tiina Mäki as well as the contact tracing unit from Tampere University Hospital for patient management and guidance. The authors would also like to thank statistician Mika Helminen from Tampere University for his advice on statistical analysis.

References

1. Akbarialiabad H, Taghrir MH, Abdollahi A, et al. Long COVID, a comprehensive systematic scoping review. *Infection*. 2021 Jul 28:1–24.
2. van Kessel SAM, Olde Hartman TC, Lucassen PLBJ, et al. Post-acute and long-COVID-19 symptoms in patients with mild diseases: a systematic review. *Fam Pract*. 2021 Jul 16.
3. Bickel S, Popler J, Lesnick B, Eid N. Impulse oscillometry: Interpretation and practical applications. *Chest American College of Chest Physicians*; 2014; 146: 841–847.
4. Huang Y, Tan C yan, Wu J, et al. Impact of Coronavirus Disease 2019 on Pulmonary Function in Early Convalescence Phase. *Respiratory Research*; 2020; : 1–10.
5. Lindahl A, Reijula J, Malmberg LP, et al. Small airway function in Finnish COVID-19 survivors. *Respiratory Research* 2021; 22: 237.
6. Lopes AJ, Mafort TT, da Cal MS, et al. Impulse Oscillometry Findings and Their Associations With Lung Ultrasound Signs in COVID-19 Survivors. *Respir Care*. 2021 Nov;66(11):1691-1698.
7. King GG, Bates J, Berger KI, et al. Technical standards for respiratory oscillometry. *European Respiratory Journal* 2020; 55: 1–21.
8. Oostveen E, MacLeod D, Lorino H, et al. The forced oscillation technique in clinical practice: Methodology, recommendations and future developments. *European Respiratory Journal* 2003; 22: 1026–1041.
9. Graham BL, Steenbruggen I, Barjaktarevic IZ, et al. Standardization of spirometry 2019 update an official American Thoracic Society and European Respiratory Society technical statement. *American Journal of Respiratory and Critical Care Medicine* 2019; 200: E70–E88.
10. Oostveen E, Boda K, van der Grinten CPM, et al. Respiratory impedance in healthy subjects: Baseline values and bronchodilator response. *European Respiratory Journal* 2013; 42: 1513–1523.
11. Kainu A, Timonen KL, Toikka J, et al. Reference values of spirometry for Finnish adults. *Clinical Physiology and Functional Imaging Blackwell Publishing Ltd*; 2016; 36: 346–358.
12. Torres-Castro R, Vasconcello-Castillo L, Alsina-Restoy X, et al. Respiratory function in patients post-infection by COVID-19: a systematic review and meta-analysis. *Pulmonology* 2021. p. 328–337.

13. Nielsen KJ, Vestergaard JM, Schlünssen V, et al. Day-by-day symptoms following positive and negative PCR tests for SARS-CoV-2 in non-hospitalized healthcare workers: A 90-day follow-up study. *International Journal of Infectious Diseases* Elsevier B.V.; 2021; 108: 382–390.
14. Søråas A, Kalleberg KT, Dahl JA, et al. Persisting symptoms three to eight months after non-hospitalized COVID-19, a prospective cohort study. *PLoS One*. 2021 Aug 26;16(8):e0256142.

Table 1. Demographics, pulmonary function and symptoms in 43 patients with COVID-19 and 39 controls with other respiratory infection at the time of acute infection (visit 1) and after two months follow-up (visit 2).

Demographics	COVID-19 positive cases			Covid-19 negative controls			p-value** (pos to neg)	
	Visit 1 (n=43)	Visit 2(n=39)	p-value* (visit 1to2)	Visit 1(n=39)	Visit 2(n=35)	p-value* (visit 1to2)	Visit 1	Visit 2
Females	24 (56)			26 (67)			0.315	
Age (years)	41.0 [28.0–48.0]			39.0 [34.0–46.0]			0.658	
BMI (kg/m ²)	27.0 [25.1–31.1]			27.5 [23.3–29.7]			1	
Duration of symptoms (days)	4.6 [4.4–6.5]			5.4 [4.4–8.4]			0.514	
Comorbidities, lung disease ¹	1 (2)			1 (3)			1	
Comorbidities, other	14 (33)			14 (36)			0.752	
Current smoker	10 (23)			9 (23)			1	
Airway allergy	17 (40)			9 (23)			0.110	
Pulmonary function								
FVC (% pred)		94 [88–102]			92 [85–103]			0.623
FEV ₁ (% pred)		93 [86–99]			91 [86–105]			0.641
FEV ₁ /FVC (% pred)		99 [92–102]			100 [93–104]			1
R5 (cmH ₂ O.s/L)	2.88 [2.29–3.61]	3.31 [2.34–3.82]	0.062	2.59 [2.14–3.26]	2.86 [2.20–3.79]	0.011		
(Z-score)	-0.10 [-0.90–0.70]	-0.10 [-0.70–0.73]	0.059	-0.50 [-1.10–0.10]	-0.50 [-0.90–0.40]	0.132	0.113	0.188
R20 (cmH ₂ O.s/L)	2.67 [2.33–3.39]	2.72 [2.32–3.48]	0.152	2.65 [2.31–3.07]	2.63 [2.18–3.15]	0.255		
(Z-score)	-0.20 [-0.90–0.10]	-0.60 [-0.75–0.20]	0.102	-0.50 [-1.00– -0.10]	-0.50 [-1.00– -0.10]	0.113	0.500	0.880
R5-20 (cmH ₂ O.s/L)	0.09 [-0.05–0.28]	0.32 [0.09–0.70]	< 0.001	-0.10 [-0.23–0.18]	0.17 [-0.04–0.64]	< 0.001		
(Z-score)	0.40 [-0.40–0.70]	0.70 [-0.23–1.50]	< 0.001	-0.20 [-0.60–0.30]	0.20 [-0.10–1.10]	0.003	0.025	0.188
X5 (cmH ₂ O.s/L)	-0.92 [-1.24– -0.79]	-0.99 [-1.30– -0.81]	0.052	-0.94 [-1.05– -0.72]	-1.04 [-1.20– -0.77]	0.048		
(Z-score)	0.20 [-0.60–0.70]	-0.05 [-0.90–0.83]	0.075	0.40 [-0.10–0.90]	0.40 [-0.50–0.80]	0.224	0.282	0.129
AX (cmH ₂ O.s/L)	4.28 [2.40–7.17]	4.90 [3.02–8.21]	< 0.001	2.95 [2.20–4.77]	4.13 [2.43–9.37]	0.005		
(Z-score)	0.70 [0.10–1.50]	1.05 [0.25–1.60]	0.005	0.30 [-0.20–0.70]	0.50 [0.00–1.40]	0.018	0.045	0.398
Symptom VAS 0-100								
Need to blow nose	30 [15–57]	9 [0–24]	< 0.001	18 [6–33]	8 [0–14] [§]	0.004	0.002	0.718
Sneezing	22 [9–53] [#]	6 [0–17]	< 0.001	10 [0–23]	7 [0–15] [§]	> 0.05	0.057	0.904
Runny nose	31 [8–60]	2 [0–13]	< 0.001	13 [7–41]	3 [0–12] [§]	< 0.001	0.119	0.919
Decreased Sense of Smell/Taste	52 [13–87]	1 [0–12]	< 0.001	0 [0–11]	0 [0–1] [§]	0.009	<0.001	0.066
Cough	33 [9–59] [#]	0 [0–12]	< 0.001	12 [0–27]	1 [0–5] [§]	< 0.001	0.010	0.431
Post-nasal discharge	15 [5–42] [#]	0 [0–15]	< 0.001	18 [5–40]	5 [0–11] [§]	0.001	0.915	0.205
Thick nasal discharge	24 [5–41] [#]	0 [0–9]	< 0.001	5 [0–27]	1 [0–3] [§]	< 0.001	0.095	0.584
Wet cough	17 [5–39]	0 [0–14]	< 0.001	4 [0–20]	0 [0–1] [§]	0.001	0.185	0.653
Dyspnoea (shortness of breath)	8 [0–22] [†]	0 [0–9]	0.006	0 [0–18]	0 [0–3] [§]	> 0.05	0.043	0.842
Fatigue	50 [20–68]	13 [0–50] [†]	0.001	24 [7–51]	11 [2–22] [§]	0.001	0.027	0.816

Data are presented as median [Interquartile range] or n (%). FVC: forced vital capacity; FEV₁: forced expiratory volume in one second; R5: total airway resistance; R20: large airways resistance; R5-20: small airways resistance; X5: airway reactance; AX: area of reactance; Z-score: ratio of the difference between the measured value and that predicted with the residual standard deviation; * Related-Samples Wilcoxon Signed Rank Test and; ** Independent-Samples Median Test or Fishers exact test or Chi-square (Pearson); ¹ Asthma; [#] n=42; [†] N=41; [§] n=38; [§] n=36; The level of significance was set at p <0.05 (indicated with bold font).