

On-line supplement

Interaction between asthma and smoking increases risk of adult airway obstruction

Aanerud M et al

List of investigators in the Early life Working Group in ECRHS

Chair: Cecilie Svanes. Members: Shyamali Dharmage, Melbourne, Australia; Thorarinn Gislason, Reykjavik; Maria Gunnbjørnsdottir, Uppsala; Joachim Heinrich, Neuherberg; Deborah Jarvis, London; Benedicte Leynaert, Paris; Dora Ludvigsdottir, Reykjavik; Roberto de Marco, Verona; Francoise Neukirch, Paris; Chantal Raherison, Paris; Jordi Sunyer, Barcelona; Simona Villani, Pavia, Matthias Wjst, Neuherberg.

List of Principal Investigators and Senior Scientific Teams in ECRHS

Belgium: South Antwerp & Antwerp City (P Vermeire, J Weyler, M Van Sprundel, V Nelen). Estonia: Tartu (R Jogi, A Soon). France: Paris (F Neukirch, B Leynaert, R Liard, M Zureik), Grenoble (I Pin, J Ferran-Quentin). Germany: Erfurt (J Heinrich, M Wjst, C Frye, I Meyer). Iceland: Reykjavik (T Gislason, E Bjornsson, D Gislason, T Blondal, KB Jorundsdottir). Italy: Turin (M Bugiani, P Piccioni, E Caria, A Carosso, E Migliore, G Castiglioni), Verona (R de Marco, G Verlato, E Zanolin, S Accordini, A Poli, V Lo Cascio, M Ferrari), Pavia (A Marinoni, S Villani, M Ponzio, F Frigerio, M Comelli, M Grassi, I Cerveri, A Corsico). Netherlands: Groningen & Geleen (J Schouten, M Kerkhof). Norway: Bergen (A Gulsvik, E Omeneas, C Svanes, B Laerum). Spain: Barcelona (JM Antó, J Sunyer, M Kogevinas, JP Zock, X Basagana, A Jaen, F Burgos), Huelva (J Maldonado, A Pereira, JL Sanchez), Albacete (J Martinez-Moratalla Rovira, E Almar), Galdakao (N Muniozguren, I Urritia), Oviedo (F Payo). Sweden: Uppsala (C Janson, G Boman, D Norback, M Gunnbjørnsdottir), Goteborg (K Toren, L Lillienberg, AC Olin, B Balder, A Pfeifer-Nilsson, R Sundberg), Umea (E Norrman, M Soderberg, K Franklin, B Lundback, B Forsberg, L Nystrom). Switzerland: Basel (N Künzli, B Dibbert, M Hazenkamp, M Brutsche, U Ackermann-Liebrich). United Kingdom: Norwich (D Jarvis, B Harrison), Ipswich (D Jarvis, R Hall, D Seaton)

Centres taking part at their own expense: Australia: Melbourne (M Abramson, S Dharmage, J Raven, EH Walters). France: Bordeaux (A Taylard, C Raherison), Montpellier (J Bousquet, P Demoly). Germany: Hamburg (K Richter). USA: Portland (M Osborne, S Buist, W Vollmer, L Johnson).

Funding

None of the study sponsors/funders had any role on study design, data collection, data analysis, data interpretation or writing of the report. The corresponding author had full access to all the data in the study and had full responsibility for the decision to submit for publication.

The first author, Marianne Aanerud, was supported by a grant from the HelseVest and a travelling grant from The Norwegian Association for Physicians.

The coordination of ECRHS II was supported by the European Commission, as part of their Quality of Life programme. The following bodies funded the local studies in ECRHS II in this article.

Albacete—Fondo de Investigaciones Santarias (grant code: 97/0035-01, 99/0034-01, and 99/0034-02), Hospital Universitario de Albacete, Consejeria de Sanidad.

Antwerp—FWO (Fund for Scientific Research)- Flanders Belgium (grant code: G.0402.00), University of Antwerp, Flemish Health Ministry.

Barcelona—Fondo de Investigaciones Sanitarias (grant code: 99/0034-01, and 99/0034-02), Red Respira (RTIC 03/11 ISC IIF).

Basel—Swiss National Science Foundation, Swiss Federal Office for Education & Science, Swiss National Accident Insurance Fund (SUVA).

Bergen—Norwegian Research Council; Norwegian Asthma & Allergy Association (NAAF); Glaxo Wellcome AS, Norway Research Fund.

Bordeaux—Institut Pneumologique d’Aquitaine.

Erfurt—GSF-National Research Centre for Environment & Health, Deutsche Forschungsgemeinschaft (DFG) (grant code FR 1526/1-1).

Galdakao—Basque Health Department.

Gothenburg—Swedish Heart Lung Foundation, Swedish Foundation for Health Care Sciences & Allergy Research, Swedish Asthma & Allergy Foundation, Swedish Cancer & Allergy Foundation.

Grenoble—Programme Hospitalier de Recherche Clinique-DRC de Grenoble 2000 no. 2610, Ministry of Health, Direction de la Recherche Clinique, Ministere de l’Emploi et de la Solidarite, Direction Generale de la Sante, CHU de Grenoble, Comite des Maladies Respiratoires de l’Isere.

Hamburg—GSF-National Research Centre for Environment & Health, Deutsche Forschungsgemeinschaft (DFG) (grant code MA 711/4-1).

Ipswich and Norwich—National Asthma Campaign (UK).

Huelva—Fondo de Investigaciones Sanitarias (FIS) (grant code: 97/0035-01, 99/0034-01, and 99/0034-02).

Montpellier—Programme Hospitalier de Recherche Clinique-DRC de Grenoble 2000 no. 2610, Ministry of Health, Direction de la Recherche Clinique, CHU de Grenoble, Ministere de l’Emploi et de la Solidarite, Direction Generale de la Sante, Aventis (France), Direction Régionale des Affaires Sanitaires et Sociales Languedoc-Roussillon. *Oviedo*—Fondo de Investigaciones Santarias (FIS) (grant code: 97/0035-01, 99/0034-01, and 99/0034-02).

Paris—Ministere de l’Emploi et de la Solidarite, Direction Generale de la Sante, UCBPharma (France), Aventis (France), Glaxo France, Programme Hospitalier de Recherche Clinique-DRC de Grenoble 2000 no. 2610, Ministry of Health, Direction de la Recherche Clinique, CHU de Grenoble.

Pavia—Glaxo, Smith & Kline Italy, Italian Ministry of University and Scientific and Technological Research (MURST), Local University Funding for Research 1998 & 1999 (Pavia, Italy).

Portland—American Lung Association of Oregon, Northwest Health Foundation, Collins Foundation, Merck Pharmaceutical.

Reykjavik—Icelandic Research Council, Icelandic University Hospital Fund.

Tartu—Estonian Science Foundation.

Turin—ASL 4 Regione Piemonte (Italy), AO CTO/ICORMA Regione Piemonte (Italy), Ministero dell’Università e della Ricerca Scientifica (Italy), Glaxo Wellcome spa (Verona, Italy).

Umeå—Swedish Heart Lung Foundation, Swedish Foundation for Health Care Sciences & Allergy Research, Swedish Asthma & Allergy Foundation, Swedish Cancer & Allergy Foundation.

Uppsala—Swedish Heart Lung Foundation, Swedish Foundation for Health Care Sciences & Allergy Research, Swedish Asthma & Allergy Foundation, Swedish Cancer & Allergy Foundation.

Verona—University of Verona; Italian Ministry of University and Scientific and Technological Research (MURST); Glaxo, Smith & Kline Italy.

The following bodies funded ECRHS I for centres in ECRHS II:

Asthma Foundation of Victoria, Allen+Hanburys (Australia); Belgian Science Policy Office, National Fund for Scientific Research; Ministère de la Santé, Glaxo France, Institut Pneumologique d’Aquitaine, Contrat de Plan Etat-Région Languedoc-Rousillon, CNMATS, CNMRT (90MR/10, 91AF/6), Ministre délégué de la santé, RNSP, France; GSF, and the Bundesminister für Forschung und Technologie, Bonn, Germany; Ministero dell’Università e della Ricerca Scientifica e Tecnologica, CNR, Regione Veneto grant RSF n. 381/05.93, Italy; Norwegian Research Council project no. 101422/310; Dutch Ministry of Wellbeing, Public Health and Culture, Netherlands; Ministerio de Sanidad y Consumo FIS (grants #91/0016060/00E-05E and #93/0393), and grants from Hospital General de Albacete, Hospital General Juan Ramón Jiménez, Consejería de Sanidad, Principado de Asturias, Spain; The Swedish Medical Research Council, the Swedish Heart Lung Foundation, the Swedish Association against Asthma and Allergy; Swiss National Science Foundation grant 4026-28099; National Asthma Campaign, British Lung Foundation, Department of Health, South Thames Regional Health Authority, UK; United States Department of Health, Education and Welfare Public Health Service (grant #2 S07 RR05521-28).

Online repository

Table E1. Sensitivity analysis. Association of smoking status and asthma with adult airway obstruction. Excluding individuals with airway obstruction at baseline but not at follow-up (n=18508)

	<i>Adult airway obstruction, Adj</i>			
	%	OR‡	95% CI†	
No asthma, never smoker		0.3	1.00	-
No asthma, current smoker		1.7	4.89	3.07 7.79
Early-onset asthma, never smoker		8.0	23.48	13.49 40.88
Late-onset asthma, current smoker		10.1	28.25	15.76 50.63
RERI		-0.21		
Late-onset asthma, never smoker		4.9	10.82	6.20 18.88
Late-onset asthma, current smoker		11.0	27.78	16.31 47.32
RERI		15.4	*	

‡ Odds ratio adjusted for education, age, height, sex, sample and country. † 95%CI= 95% confidence interval. RERI= Relative Excess due to Interaction.

Early-onset defined as asthma onset before 10 yrs of age, late-onset defined as asthma onset after 10 yrs of age

Table E2. Association of smoking history and asthma with adult airway obstruction

Asthma	Smoking	<i>N_{adult airway obstruction/total}</i>		OR [‡]	95% CI [†]
No asthma	Never smoker	29/8213	0.3%	1.00	-
	Current smoker	127/7020	1.8%	4.43	(2.88,6.83)
Early-onset asthma	Never smoker	59/679	8.7%	20.34	(12.22,33.87)
	Current smoker	42/387	10.8%	24.16	(14.13,41.33)
Late-onset asthma	Never smoker	56/921	6.1%	11.09	(6.67,18.42)
	Current smoker, started after asthma onset	18/141	12.8%	26.95	(13.74,52.86)
	Current smoker, started before asthma onset	49/410	12.0%	24.36	(14.40,41.21)

n=17771. [‡]Odds ratio adjusted for education, age height, sex, sample and centre. [†] 95%CI= 95% confidence interval. Early onset asthma before 10 years of age, late onset after 10 years of age.

Table E3. Association of smoking history and asthma with adult airway obstruction, former smokers included.

Asthma	Smoking	<i>N_{adult airway obstruction/total}</i>		OR [‡]	95% CI
No asthma	Never smoker	29/8213	0.3%	1.00	-
	Current smoker	127/7020	1.8%	4.21	(2.74,6.46)
	Former smoker	37/4555	0.8%	1.85	(1.11,3.07)
Early-onset asthma	Never smoker	59/679	8.7%	19.60	(11.86,32.39)
	Current smoker	40/380	10.5%	22.52	(13.20,38.52)
	Former smoker	34/287	11.8%	25.11	(14.36,43.89)
Late-onset asthma	Never smoker	56/921	6.1%	10.85	(6.56,17.96)
	Current smoker, started after asthma onset	17/126	13.5%	28.33	(14.25,56.31)
	Smoker before asthma	50/412	12.1%	23.06	(13.70,38.82)
	Former smoker after asthma	6/91	6.6%	10.58	(5.80,19.28)
	Former smoker before asthma	27/374	7.2%	12.09	(4.41,33.15)

n=23058. Odds ratio adjusted for education, age height, sex, sample, and country. 95% CI = 95% Confidence interval. Early-onset asthma before 10 years of age, late-onset asthma after 10 years of age.

Table E4. Risk of adult airway obstruction, odds ratio (confidence interval), by sex, smoking, and early-onset asthma, stratified by age of asthma onset.

		Odds ratio males	Confidence interval		Odds ratio females	Confidence interval	
No asthma	Non smoker	1			1		
	Current smoker	2.5	(0.8, 1.2)		4.3	(1.7, 2.8)	
Asthma onset age 0-4	Non smoker	15.8	(6.6, 11.3)		22.9	(11.4, 22.7)	
	Current smoker	21.2	(10.6, 21.2)		32.3	(19.0, 45.9)	
Asthma onset age 5-9	Non smoker	16.4	(6.8, 11.7)		19.8	(11.3, 26.4)	
	Current smoker	12.7	(6.7, 14.4)		30.5	(21.3, 70.8)	
Asthma onset age 10-14	Non smoker	12.5	(6.6, 13.9)		11.0	(6.9, 18.2)	
	Current smoker	26.7	(14.0, 29.6)		51.7	(30.6, 74.8)	
Asthma onset age 15-19	Non smoker	7.1	(4.3, 11.0)		24.1	(13.3, 29.6)	
	Current smoker	10.4	(7.7, 29.1)		31.5	(18.9, 47.3)	
Asthma onset age > 20	Non smoker	10.8	(6.0, 13.4)		22.2	(11.6, 24.1)	
	Current smoker	21.6	(14.0, 39.9)		15.1	(10.5, 34.0)	

Table E5. Population attributable fractions of adult airway obstruction from asthma and smoking

	N total (%)	N adult airway obstruction (%)	OR	95%CI	AF
Asthma					
No asthma	13884 (91.1)	119 (58.9)	1		
Early-onset asthma	594 (3.9)	34 (16.8)	10.23	(7.77-13.49)	15.2%
Late-onset asthma	761 (5.0)	49 (24.3)	5.83	(4.46-7.61)	20.1%
Early+late	1355 (8.9)	83 (41.1)	7.23	(5.74-9.11)	35.3%
Total	15239 (100)	202 (100)			
Smoking					
Never smoker	6669 (43.4)	51 (24.1)	1		
Current smoker	3784 (24.6)	56 (26.4)	2.45	(1.92-3.12)	29.3%
Former smoker	4908 (32.0)	105 (49.5)	1.34	(1.01-1.78)	6.7%
Current and former smoker	8692 (56.6)	161 (75.9)	1.93	(1.54-2.42)	36.0%
Total	15361 (100)	212 (100)			
Interaction asthma*smoking					
No asthma, never smoker	5892(38.9)	20 (9.9)	1		
No asthma, current smoker	4513 (29.8)	71 (35.1)	4.27	(2.78-6.54)	26.9%
No asthma, former smoker	3403 (22.5)	28 (13.9)	1.85	(1.11-3.06)	6.3%
Early-onset asthma, never smoker	286 (1.9)	11 (5.4)	19.82	(12.01-32.69)	5.2%
Late-onset asthma, current smoker	149 (1.0)	9 (4.5)	22.48	(13.23-38.20)	4.3%
Early-onset asthma, former smoker	154 (1.0)	14 (6.9)	26.80	(15.58-46.09)	6.7%
Late-onset asthma, never smoker	405 (2.7)	19 (9.4)	10.73	(6.50-17.74)	8.5%
Late-onset asthma, current smoker	189 (1.3)	20 (9.9)	23.63	(14.50-38.52)	9.5%
Late-onset asthma, former smoker	166 (1.1)	10 (5.0)	10.57	(6.01-18.61)	4.5%
Any versus none*	9265 (61.1)	182 (90.1)	6.23	(4.20-9.23)	71.9%

OR: Odds ratios estimated from GEE random effects model adjusted for age, sex, sample, country, height and education. 95% CI: 95% confidence interval. AF: Attributable fraction. * Having any risk factor (asthma, either early- or late onset, or smoking, either as current or former smoker, compared to subjects who never smoked and never have had asthma).

Table E6. Association of adult airway obstruction with atopy and smoking, stratified by asthma (n=16108)

		Adult airway obstruction ¹				
		Atopy	Smoking status [†]	% <i>Adult airway obstruction</i>	Odds ratio ²	95% confidence interval
All	No atopy		Never smoker	0.7%	1.00	-
			Current smoker	2.6%	3.57	2.38 5.35
	Atopy		Never smoker	3.1%	3.65	2.47 5.41
			Current smoker	3.9%	4.76	3.13 7.24
			<i>RERI</i> ⁴	-1.46	-3.15	0.24
No asthma ³	No atopy		Never smoker	0.4%	1.00	-
			Current smoker	0.5%	1.38	0.62 3.05
	Atopy		Never smoker	1.8%	4.42	2.61 7.51
			Current smoker	1.7%	4.51	2.44 8.35
			<i>RERI</i> ⁴	-0.28	-2.53	1.96
Asthma ³	No atopy		Never smoker	4.7%	1.00	
			Current smoker	8.8%	1.80	1.02 3.18
	Atopy		Never smoker	12.9%	2.99	1.59 5.62
			Current smoker	11.3%	2.29	1.25 4.20
			<i>RERI</i> ⁴	-1.50	-3.33	0.33

[†]Excluding former smokers. ¹ Adult airway obstruction defined as pre-bronchodilator FEV1/FVC < 70 % and FEV1 < 80 % predicted. ² Odds ratio adjusted for education, age height, sex, sample and centre ³Asthma = early or late onset asthma ⁴ RERI= Relative Excess due to Interaction.

Table E7. Number and percentage of asthma according to adult airway obstruction at baseline and follow-up

Adult airway obstruction†	Total	Ever asthma at baseline		p-value‡
		n	%	
No adult airway obstruction	8,049	976	12.1%	
Adult airway obstruction at baseline <u>not</u> at follow-up (1)	67	42	62.7%	
Adult airway obstruction at baseline <u>and</u> at follow-up (2)	97	59	60.8%	0.810
Adult airway obstruction only at follow-up	101	42	41.6%	
Total	8,314	1,119	13.5%	

† Adult airway obstruction defined as pre-bronchodilator FEV1/FVC < 70 % and FEV1 < 80 % predicted ‡p-value from chi² test (comparing distribution of asthmatic in (1) and (2), Pearson chi²(1) = 0.0581)

Figure E1. Smoking intensity for current smokers by asthma status (no asthma, early onset, and late onset)



