



HIV and multidrug-resistant tuberculosis: overlapping epidemics

To the Editor:

People infected with *Mycobacterium tuberculosis* and HIV are much more likely to develop active tuberculosis (TB) than people with *M. tuberculosis* but without HIV [1]. Patients infected with multidrug-resistant (MDR)-TB (defined as resistance to at least rifampicin and isoniazid, the two most powerful anti-TB drugs) require longer, more expensive treatment regimens than drug-susceptible TB, with poorer treatment success [2, 3]. Therefore, MDR-TB poses a major challenge to the control of TB, with an estimated global disease incidence in 2012 of ~450 000 cases (95% CI 300 000–600 000) [4]. Although HIV is a powerful risk factor for all forms of TB and institutional outbreaks of MDR-TB among people living with HIV have been reported [5], population-level data on the association between HIV infection and MDR-TB are limited.

We explored the relationship between HIV infection and MDR-TB disease using data reported by member states to the World Health Organization (WHO) within the context of the Global Project on Anti-TB Drug Resistance Surveillance. The data were aggregated numbers of cases reported from either drug resistance surveys or continuous surveillance systems. Such surveys are epidemiological studies designed to measure drug resistance among a representative sample of notified pulmonary TB patients. Continuous surveillance is based on routine drug susceptibility testing of all bacteriologically confirmed TB patients. Subnational level data that were not representative of the entire country were excluded from the analysis, except for the Russian Federation and Ukraine, which are high MDR-TB burden countries for which high quality national level data were not available. The data included in the analysis met the criteria for data quality and national representativeness provided in detail elsewhere [4, 6]. The laboratory methods used for diagnosis were endorsed by WHO.

For each country, the relationship between HIV infection and MDR-TB disease was investigated by logistic regression to calculate odds ratios and 95% confidence intervals, using Stata (version 12; StataCorp, College Station, TX, USA). For countries with data from multiple years, robust standard errors accounted for within-country time dependencies. In order to minimise bias, data for a given year were excluded if $\leq 25\%$ of reported TB patients had a documented HIV test result. 41 countries, accounting for 25% of the estimated global MDR-TB burden in 2012, met the inclusion criteria for ≥ 1 year. Most were high-income countries and/or in the European Region. Only four countries were in the African region, which accounted for 75% of the global number of HIV-positive TB cases in 2012 [4]. Of these 41 countries, odds ratios could be calculated for 24 countries; the other 17 countries reported no HIV-positive MDR-TB cases (table 1). Complete HIV and MDR-TB data were available for 104 781 TB patients from 1997–2012.

For 11 of the 24 countries for which the analysis was performed, HIV-positive TB patients had a significantly higher odds ($p < 0.05$) of MDR-TB disease than HIV-negative TB patients (table 1). Seven of these countries were in eastern Europe and central Asia: Estonia, Kazakhstan, Latvia, the Republic of Moldova, the Russian Federation, the Ukraine and Uzbekistan. For almost all of these 11 countries, the prevalence of MDR-TB among newly diagnosed TB cases (table 1) was higher than the estimated global average of 3.6% (95% CI 2.1–5.1%) in 2012 [4]. Although the odds ratio was highest in Kuwait, only low numbers of HIV-positive cases were reported. HIV-positive TB patients in the USA had a lower odds of MDR-TB disease than HIV-negative patients.

As data were included if $\geq 75\%$ of reported TB patients had a documented HIV test, the possibility of bias due to missing data cannot be ruled out. Although a positive association between HIV infection and MDR-TB disease was demonstrated in less than half of the countries, a recent systematic review and meta-analysis found an odds of MDR-TB in HIV-positive patients that was 1.24 times (95% CI 1.04–1.43) higher than in HIV-negative patients [7]. A subgroup analysis showed that this association was stronger for primary MDR-TB disease than acquired MDR-TB disease, which supports previous findings [8]. However, most of these studies were from institutional settings, where outbreaks are known to occur more commonly in HIV-positive patients [9]. At the population level, a detailed analysis of surveillance data from the Republic of Moldova showed that, even after adjustment for potential confounders, a positive association HIV and MDR-TB existed [10].

TABLE 1 Odds of multidrug-resistant (MDR) tuberculosis (TB) disease in HIV-positive patients compared with HIV-negative patients

Country	Patients with data on HIV and MDR-TB included in analysis n		OR (95% CI)	p-value	Period from which data were available years	Estimated HIV prevalence in incident TB cases in 2012 %	MDR-TB prevalence in new/previously treated TB cases % (year)*
	Total	HIV-positive					
Australia	1456	27	47	0.827	2	2.5	1.9/6.5 (2012)
Bahamas	36	11	2	0.653	2	23.0	3.7/0 (2012)
Belarus	4668	219	2170	0.081	2	4.2	34.8/68.6 (2012)
Belgium	2723	165	62	0.375	4	3.8	1.3/11.4 (2012)
Benin	420	58	8	0.914	1	14.0	0.5/13.2 (2010)
Cuba	1828	47	30	0.170	7	3.6	0.7/11.8 (2012)
Estonia	1328	165	348	0.037	5	13.0	19.7/50.0 (2012)
France	2426	235	54	0.000	2	6.9	0.5/13.2 (2009)
Israel	1321	72	55	0.005	5	3.8	4.7/33.3 (2012)
Kazakhstan	40 975	577	7861	0.000	2	1.2	22.9/55.0 (2012)
Kuwait	865	3	14	0.000	2	0	0/0 (2011)
Latvia	828	61	129	0.047	1	9.3	11.1/32.0 (2011)
Malawi	1761	820	32	0.454	1	62.0	0.4/4.8 (2012)
Mexico	1600	58	35	0.806	1	5.8	2.4/6.5 (2009)
Namibia	1432	672	98	0.060	1	49.0	3.8/16.4 (2008)
Nigeria	1248	175	66	0.129	1	25.0	2.9/14.3 (2010)
Republic of Moldova	8395	565	3553	0.000	4	6.0	23.7/62.3 (2012)
Russian Federation [†]	2047	15	469	0.001	3	7.1	23.1/48.6 (2011)
Singapore	2700	117	30	0.728	3	3.7	1.6/3.2 (2012)
Swaziland	565	451	114	0.005	1	77.0	7.7/33.9 (2009)
Uganda	1313	399	31	0.533	1	53.0	1.4/12.1 (2011)
Ukraine ⁺	1450	307	369	0.006	1	11.0	14.4/32.2 (2012)
USA	12 190	935	181	0.000	2	9.6	1.0/3.0 (2012)
Uzbekistan	4722	87	1806	0.001	1	2.0	23.3/62.0 (2011)

Odds ratios are rounded to the nearest decimal point for display purposes. Significance was assessed prior to rounding. The 17 countries that did not report any HIV-positive MDR-TB cases in the years eligible to be included in the study were: Albania, Bahrain, Bulgaria, Costa Rica, Iceland, Jordan, Mauritius, Micronesia, New Zealand, Oman, Palau, Qatar, the Republic of Macedonia, Slovakia, Uruguay, the Marshall Islands and Montenegro. #: the most recent high quality MDR-TB prevalence data for new and previously treated TB cases are given; †: odds ratio calculated from data from Tomsk Oblast only; ‡: odds ratio calculated from data from Donetsk Oblast only.

Among those 11 countries with a positive association between HIV infection and MDR-TB disease, there was wide variation in the estimated prevalence of HIV among incident TB cases in 2012 (table 1) [4]. Given that the analysis was performed using aggregated rather than individual patient data, there are likely to be risk factors common to HIV-positive and MDR-TB patients that could not be explored in our analysis, such as shared behaviours or population characteristics. Further investigation is needed for specific high risk groups, such as prisoners or people living in congregate settings, miners, or injecting drug users. Additionally, the relationship between HIV and MDR-TB probably depends on the epidemiological setting. In the USA, where a negative association was observed between HIV infection and MDR-TB disease, investigation into the demographics of these patients is required in order to identify potential confounders. However, a cross-sectional study in eight states did not demonstrate a significant association between HIV infection and MDR-TB disease [11]. HIV has been shown to be an independent risk factor for drug resistance acquired during treatment in California [12].

Data were only available for three countries listed among both the 27 high MDR-TB burden countries and 41 high TB/HIV burden countries: Nigeria, the Russian Federation and the Ukraine. In order to better understand the relationship between HIV infection and MDR-TB, more high quality data are needed from other high burden countries. This includes the Democratic Republic of Congo, Ethiopia, Myanmar, South Africa and Vietnam. In these five countries, the percentage of estimated cases of MDR-TB that were detected and notified in 2012 ranged from just 2.2% to 30% and the percentage of TB patients with a known HIV status in 2012 ranged from 13 to 66%, excluding South Africa [4]. This is probably a reflection of low levels of testing for MDR-TB and HIV among TB patients as well as underreporting. Priorities in these countries should include integrated TB and HIV services and programmes, with universal access to HIV testing and routine and prompt TB drug susceptibility testing for patients suspected to have MDR-TB. This must be underpinned by greater awareness of healthcare workers, increased laboratory capacity, and improved data management and reporting [13]. The roll-out of rapid diagnostic tests and the use of MDR-TB treatment models that include community-based care could rapidly increase MDR-TB case notification and enrolment on second-line treatment, ultimately limiting spread [14]. Development of region- and country-specific comprehensive responses, including research for better diagnosis, treatment and scale-up of quality services, is essential [15].



@ERSpublications

Development of region- and country-specific comprehensive responses to HIV/MDR-TB co-infection is essential <http://ow.ly/tgkvt>

Anna S. Dean, Matteo Zignol, Dennis Falzon, Haileyesus Getahun and Katherine Floyd
Global Tuberculosis Programme, World Health Organization, Geneva, Switzerland.

Correspondence: K. Floyd, World Health Organization, 20 Avenue Appia, 1211 Geneva, Switzerland. E-mail: floydk@who.int

Received: Dec 13 2013 | Accepted after revision: Jan 25 2014

Conflict of interest: None declared.

Acknowledgements: All authors are staff members of WHO. The authors alone are responsible for the views expressed in this publication and they do not necessarily represent the decisions or policies of WHO.

References

- 1 Selwyn PA, Hartel D, Lewis VA, *et al.* A prospective study of the risk of tuberculosis among intravenous drug users with human immunodeficiency virus infection. *N Engl J Med* 1989; 320: 545–550.
- 2 Migliori GB, Sotgiu G, Gandhi NR, *et al.* Drug resistance beyond extensively drug-resistant tuberculosis: individual patient data meta-analysis. *Eur Respir J* 2013; 42: 169–179.
- 3 Falzon D, Gandhi N, Migliori GB, *et al.* Resistance to fluoroquinolones and second-line injectable drugs: impact on multidrug-resistant TB outcomes. *Eur Respir J* 2013; 42: 156–168.
- 4 World Health Organization. Global Tuberculosis Report 2013. www.who.int/tb/publications/global_report/en/index.html Date last accessed: November 1, 2013. Date last updated: 2013.
- 5 Frieden TR, Sherman LF, Maw KL, *et al.* A multi-institutional outbreak of highly drug-resistant tuberculosis: epidemiology and clinical outcomes. *JAMA* 1996; 276: 1229–1235.
- 6 World Health Organization. Guidelines for surveillance of drug resistance in tuberculosis. www.who.int/tb/publications/mdr_surveillance/en/# Date last accessed: August 6, 2013. Date last updated: 2009.
- 7 Mesfin YM, Hailemariam D, Biadgign S, *et al.* Association between HIV/AIDS and multi-drug resistance tuberculosis: a systematic review and meta-analysis. *PLoS ONE* 2014; 9: e82235.
- 8 Suchindran S, Brouwer ES, Van Rie A. Is HIV infection a risk factor for multi-drug resistant tuberculosis? A systematic review. *PLoS ONE* 2009; 4: e5561.
- 9 Wells CD, Cegielski JP, Nelson LJ, *et al.* HIV infection and multidrug-resistant tuberculosis: the perfect storm. *J Infect Dis* 2007; 196: Suppl. 1, S86–S107.

- 10 Jenkins HE, Plesca V, Ciobanu A, *et al.* Assessing spatial heterogeneity of multidrug-resistant tuberculosis in a high-burden country. *Eur Respir J* 2012; 42: 1291–1301.
- 11 Moonan PK, Teeter LD, Salcedo K, *et al.* Transmission of multidrug-resistant tuberculosis in the USA: a cross-sectional study. *Lancet Infect Dis* 2013; 13: 777–784.
- 12 Porco TC, Oh P, Flood JM. Antituberculosis drug resistance acquired during treatment: an analysis of cases reported in California, 1994–2006. *Clin Infect Dis* 2013; 56: 761–769.
- 13 Falzon D, Jaramillo E, Wares F, *et al.* Universal access to care for multidrug-resistant tuberculosis: an analysis of surveillance data. *Lancet Infect Dis* 2013; 13: 690–697.
- 14 Nardell E, Dharmadhikari A. Turning off the spigot: reducing drug-resistant tuberculosis transmission in resource-limited settings. *Int J Tuberc Lung Dis* 2010; 14: 1233–1243.
- 15 Getahun H, Gunneberg C, Granich R, *et al.* HIV infection–associated tuberculosis: the epidemiology and the response. *Clin Infect Dis* 2010; 50: S201–S207.

Eur Respir J 2014; In Press | DOI: 10.1183/09031936.00205413