



## ERRATUM

“PHYSICAL ACTIVITY IN PATIENTS WITH COPD”. H. WATZ, B. WASCHKI, T. MEYER AND H. MAGNUSSEN. *EUR RESPIR J* 2009; 33: 262–272.

Unfortunately, in table 4 of the above manuscript, the cut-off point of the 6-min walk distance (6MWD) was reported incorrectly. It should have been given as 445 m and not 44 m. The corrected table is presented below.

<b>TABLE 4</b> Receiver operating characteristic curves for prediction of very inactive <sup>#</sup> patients with chronic obstructive pulmonary disease											
Predictors	Area	SE	p-value	95% CI of area	Cut-off point <sup>†</sup>	Sensitivity	Specificity	Positive predictive value	Negative predictive value	Likelihood ratio of a positive test result	Likelihood ratio of a negative test result
<b>GOLD stages</b>	0.823	0.035	<0.001	0.76–0.89	III	0.80	0.75	0.69	0.85	3.25	0.26
<b>BODE score</b>	0.796	0.036	<0.001	0.73–0.87	3	0.67	0.79	0.69	0.78	3.23	0.42
<b>FEV<sub>1</sub> % pred</b>	0.795	0.036	<0.001	0.73–0.87	54	0.80	0.71	0.65	0.84	2.78	0.28
<b>6MWD m</b>	0.773	0.037	<0.001	0.70–0.85	445	0.79	0.74	0.68	0.84	3.06	0.29
<b>MMRC grade</b>	0.768	0.037	<0.001	0.70–0.84	2	0.65	0.78	0.67	0.77	3.01	0.44

CI: confidence interval; GOLD: Global Initiative for Chronic Obstructive Lung Disease; BODE: body mass index, airway obstruction, dyspnoea, exercise capacity; FEV<sub>1</sub>: forced expiratory volume in one second; 6MWD: 6-min walk distance; MMRC: modified Medical Research Council dyspnoea scale. <sup>#</sup>: defined as a physical activity level <1.40, including 66 patients with COPD (40.5% of all COPD patients); <sup>†</sup>: equal or greater than.